# MAT 347 A proof of Sylow's Theorems October 22, 2018

Given a finite group G and a subgroup  $H \leq G$ , recall that H is a p-subgroup if  $|H| = p^k$  for some k, and H is a  $Sylow\ p$ -subgroup if |H| is the highest power of p dividing G. The goal for today is to prove Sylow's Theorem:

**Theorem** Let G be a group of order  $p^{\alpha}m$ , where p is a prime not dividing m. Let  $n_p$  denote the number of Sylow p-subgroups of G.

- 1. G has a Sylow p-subgroup  $(n_p > 0)$ .
- 2. If P is a Sylow p-subgroup of G and Q is any p-subgroup of G, then there exists  $g \in G$  such that  $Q \leq gPg^{-1}$ . In particular, all Sylow p-subgroups are conjugate.
- 3.  $n_p \equiv 1 \pmod{p}$ , and  $n_p = |G: N_G(P)| \mid m$  for any Sylow *p*-subgroup P of G.

## Part 1

- 1. Prove that  $\binom{p^{\alpha}m}{p^{\alpha}} \equiv m \pmod{p}$ .
  - *Hint:* What is  $(1+x)^p$  when you reduce the cofficients modulo p? What about  $(1+x)^{p^2} = ((1+x)^p)^p$ ?  $(1+x)^{p^{\alpha}}$ ? And finally,  $(1+x)^{p^{\alpha}m} = ((1+x)^p)^m$ ?
- 2. Let S denote the collection of all subsets of G with cardinality  $p^{\alpha}$ . G acts on S by left multiplication. Prove that there exists an orbit  $\mathcal{O}$  of this action such that  $p \nmid |\mathcal{O}|$ . Hint: What does the quantity in Question 1 count?
- 3. Given  $X \in \mathcal{O}$ , prove that  $p^{\alpha} \mid |\operatorname{Stab}(X)|$ . (Here,  $\mathcal{O}$  is as in Question 2.)
- 4. With X as above, let  $x \in X$ . What is the relationship between the sets  $\operatorname{Stab}(X)x$  and X? Use this to prove  $|\operatorname{Stab}(X)| \leq p^{\alpha}$ , and conclude that  $\operatorname{Stab}(X)$  is a Sylow p-subgroup.

#### Lemma

5. Let H be a p-group ( $|H| = p^n$  for some n) acting on a set T, and let  $Fix(H) = \{t \in T : \forall h \in H, h \cdot t = t\}$  denote the set of elements of T which are fixed by the action. Prove that  $|Fix(H)| \equiv |T| \pmod{p}$ .

## Part 2

6. Let P be a Sylow p-subgroup, and Q any p-subgroup of G. Then Q acts on G/P (the set of left cosets) by left multiplication. Prove that there exists a coset of P which is fixed by this action.

*Hint:* Use the lemma.

7. Take  $g \in G$  so that gP is fixed by the action of Q. Prove that  $Q \leq gPg^{-1}$ .

## Part 3

8. Use the Second Sylow Theorem to prove that  $n_p = |G: N_G(P)|$ . Deduce that  $n_p$  divides m.

Hint: What group action does  $N_G(P)$  have anything to do with?

- 9. Given a Sylow p-subgroup P, P acts on the set of all Sylow p-subgroups by conjugation. Show that there is a fixed point P'. For any fixed point P' deduce that  $P \leq N_G(P')$  and  $P' \triangleleft N_G(P')$ .
- 10. Show that P' = P, and conclude that  $n_p \equiv 1 \pmod{p}$ .

Hint: Apply the Second Sylow Theorem to  $N_G(P')$  and remember the lemma.

Fun problem: Generalising your proof to Question 1, show that  $\binom{n}{k} \equiv \binom{n_0}{k_0} \binom{n_1}{k_1} \cdots \pmod{p}$ , where  $n = n_0 + pn_1 + \cdots$  and  $k = k_0 + pk_1 + \cdots$  are the base p expansions of the integers n, k (meaning  $n_i, k_i \in \{0, 1, \dots, p-1\}$ ). For example,  $\binom{100}{50} \equiv \binom{2}{1} \binom{0}{0} \binom{2}{1} \equiv 4 \pmod{7}$ .