

Putnam Problems: Algebra (F.H., Fall 2015)

October 8, 2015

A1-1999 Find polynomials $f(x), g(x)$, and $h(x)$, if they exist, such that for all x ,

$$|f(x)| - |g(x)| + h(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0. \end{cases}$$

B1-2005 Find a nonzero polynomial $P(x, y)$ such that $P(\lfloor a \rfloor, \lfloor 2a \rfloor) = 0$ for all real numbers a . (Note: $\lfloor \nu \rfloor$ is the greatest integer less than or equal to ν .)

- Do there exist $n \times n$ matrices with $AB - BA = 1$ (identity)?
- Find the zeroes of the polynomial $x^4 - 6x^3 + 18x^2 - 30x + 25$, knowing that the sum of two of its roots is 4.

B1-2004 Let $P(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_0$ be a polynomial with integer coefficients. Suppose that r is a rational number such that $P(r) = 0$. Show that the n numbers

$$c_n r, c_n r^2 + c_{n-1} r, c_n r^3 + c_{n-1} r^2 + c_{n-2} r, \\ \dots, c_n r^n + c_{n-1} r^{n-1} + \cdots + c_1 r$$

are integers.

- Find all polynomials $P(x)$ such that $(x + 1)P(x) = (x - 10)P(x + 1)$.

A1-2001 Consider a set S and a binary operation $*$, i.e., for each $a, b \in S$, $a * b \in S$. Assume $(a * b) * a = b$ for all $a, b \in S$. Prove that $a * (b * a) = b$ for all $a, b \in S$.

B1-2003 Do there exist polynomials $a(x), b(x), c(y), d(y)$ such that

$$1 + xy + x^2 y^2 = a(x)c(y) + b(x)d(y)$$

holds identically?

B2-1993 For nonnegative integers n and k , define $Q(n, k)$ to be the coefficient of x^k in the expansion of $(1 + x + x^2 + x^3)^n$. Prove that

$$Q(n, k) = \sum_{j=0}^k \binom{n}{j} \binom{n}{k-2j},$$

where $\binom{a}{b}$ is the standard binomial coefficient. (Reminder: For integers a and b with $a \geq 0$, $\binom{a}{b} = \frac{a!}{b!(a-b)!}$ for $0 \leq b \leq a$, with $\binom{a}{b} = 0$ otherwise.)

A2-1999 Let $p(x)$ be a polynomial that is nonnegative for all real x . Prove that for some k , there are polynomials $f_1(x), \dots, f_k(x)$ such that

$$p(x) = \sum_{j=1}^k (f_j(x))^2.$$

B2-1999 Let $P(x)$ be a polynomial of degree n such that $P(x) = Q(x)P''(x)$, where $Q(x)$ is a quadratic polynomial and $P''(x)$ is the second derivative of $P(x)$. Show that if $P(x)$ has at least two distinct roots then it must have n distinct roots.

B2-2001 Find all pairs of real numbers (x, y) satisfying the system of equations

$$\begin{aligned} \frac{1}{x} + \frac{1}{2y} &= (x^2 + 3y^2)(3x^2 + y^2) \\ \frac{1}{x} - \frac{1}{2y} &= 2(y^4 - x^4). \end{aligned}$$

A4-1997 Let G be a group with identity e and $\phi : G \rightarrow G$ a function such that

$$\phi(g_1)\phi(g_2)\phi(g_3) = \phi(h_1)\phi(h_2)\phi(h_3)$$

whenever $g_1g_2g_3 = e = h_1h_2h_3$. Prove that there exists an element $a \in G$ such that $\psi(x) = a\phi(x)$ is a homomorphism (i.e. $\psi(xy) = \psi(x)\psi(y)$ for all $x, y \in G$).

B-4 Let n be a positive integer. Find the number of pairs P, Q of polynomials with real coefficients such that

$$(P(X))^2 + (Q(X))^2 = X^{2n} + 1$$

and $\deg P > \deg Q$.

You can find some more algebra problems at

<http://www.math.utoronto.ca/barbeau/putnamalg.pdf>.