

# MAT 351: Partial Differential Equations

## Assignment 17, due March 13, 2017

### Summary:

Consider the eigenvalue problem for the Laplacian on a domain  $D \subset \mathbb{R}^d$  with Dirichlet boundary conditions

$$-\Delta u = \lambda u \text{ on } D, \quad u = 0 \text{ on } \partial D.$$

We assume that  $D$  is bounded and that its boundary is smooth (e.g., the domain could be defined by an inequality  $D = \{x \in \mathbb{R}^d \mid g(x) > 0\}$ , where  $g$  is a smooth function that satisfies the hypotheses of the Implicit Function Theorem at every point where  $g(x) = 0$ .) Our goal is to prove that there is an infinite sequence of positive eigenvalues  $\lambda_1 < \lambda_2 \leq \dots$ , whose growth is governed by **Weyl's law**:  $\lambda_n \sim (4\pi^2) \left(\frac{n}{\text{Vol } D}\right)^{\frac{2}{d}}$ . Furthermore, we have completeness, i.e.,  $L^2(\mathbb{R}^d)$  has an orthonormal basis consisting of the corresponding eigenvectors  $\{v_n\}$ .

The main tool for the proof is the **variational characterization of eigenvalues**:

- **max-min**:  $\lambda_n = \max_{w_1, \dots, w_{n-1}} \left\{ \min_{w \perp w_1, \dots, w_{n-1}} \frac{\int_D |\nabla w|^2 dx}{\|w\|^2} \right\};$
- **min-max**:  $\lambda_n = \min_{w_1, \dots, w_n} \left\{ \max_{w \in \text{span}\{w_1, \dots, w_n\}} \frac{\int_D |\nabla w|^2 dx}{\|w\|^2} \right\}.$

In these variational problems, the eigenvalues play the role of Lagrange multipliers. The objective function is called the **Rayleigh quotient**. It is minimized by the lowest eigenvalue

$$\lambda_1 = \min_{\|w\|=1} \int_D |\nabla w|^2 dx.$$

In these formulas, it is understood that  $w_1, \dots, w_n$  and  $w$  should all satisfy the Dirichlet boundary conditions. For both the max-min and the min-max principle, the functions  $w_i$  must be linearly independent (but they need not be orthonormal). The min-max principle is widely used to obtain upper bounds on eigenvalues. The max-min principle can provide lower bounds, but it is difficult to apply, since it requires to solve two infinite-dimensional problems. The following finite-dimensional approximation method is surprisingly powerful.

- **Rayleigh-Ritz principle**: Choose  $n$  orthonormal "*trial functions*"  $w_1, \dots, w_n$  that satisfy the Dirichlet boundary conditions. Define a symmetric matrix  $A$  by

$$A_{ij} = \int_D \nabla w_i \cdot \nabla w_j dx,$$

and let  $\mu_1 \leq \dots \leq \mu_n$  be its eigenvalues. Then  $\lambda_i \leq \mu_i$  for each  $i = 1, \dots, n$ .

(There are more complicated versions of this that do not require orthogonality.)

The proof of Weyl's law proceeds by comparing  $D$  with a finite union of rectangles. Once we have Weyl's law, we will obtain completeness of the eigenfunctions from the min-max principle.

### Midterm 3: Wednesday, March 15, 5-7pm

MS 2173 (Medical Sciences Building, 1 King's College Circle)

Please let me know if you have a conflict.

#### Assignments:

Start reading Chapter 11.

69. Let  $f(x)$  be a function on the interval  $[0, 3]$  such that

$$f(0) = f(3) = 0, \quad \int_0^3 |f(x)|^2 dx = 1, \quad \int_0^3 |f'(x)|^2 dx = 1.$$

Find such a function if you can. If it cannot be found, explain why not.

70. Estimate the first eigenvalue of  $-\Delta$  with Dirichlet boundary conditions in the triangle

$$D = \{(x, y) \mid x + y < 1, x > 0, y > 0\},$$

using the Rayleigh quotient with trial function  $xy(1 - x - y)$ .

71. Let  $D$  be a smooth, bounded domain in  $\mathbb{R}^d$ .

- (a) Show that the smallest Neumann eigenvalue for  $-\Delta$  on  $D$  is given by  $\mu_0 = 0$ . What is the corresponding eigenfunction?
- (b) If, moreover  $D$  is connected, show that  $\mu_0$  is simple, by proving that the next eigenvalue  $\mu_1$  is strictly positive.

72. Let  $D$  be a smooth bounded domain in  $\mathbb{R}^d$ . Denote by  $(\lambda_n)_{n \geq 1}$  the sequence of eigenvalues of the negative Dirichlet Laplacian  $-\Delta$ , and by  $(\phi_n)_{n \geq 1}$  an orthonormal basis of corresponding eigenfunctions.

(a) Find

$$\sum_{n=1}^{\infty} \int_D \phi_n(y) dy.$$

In what sense does the series converge?

- (b) Assume that  $u$  solves the heat equation  $-\Delta u = 0$  on  $D$ , with Dirichlet boundary conditions  $u|_{\partial D} = 0$  and initial values  $u(x, 0) = f(x)$ . Express  $u$  in terms of the Dirichlet eigenvalues and eigenfunctions defined above.