

# MAT 351: Partial Differential Equations

## Assignment 7, due November 7, 2016

### Summary

**Separation of variables** is a method for finding special solutions of a partial differential equation. Sometimes we can generate all solutions of the PDE from these special solutions, using the

- **Superposition principle:** If the PDE is linear and homogeneous, then any linear combination of solutions is again a solution.

If, moreover, its coefficients are constant, then translates and derivatives of solutions are again solutions; if the equation has additional symmetries (such as rotations and dilations), they can be used to generate yet more solutions.

To explain the method, consider the wave equation on an interval

$$u_{tt} = c^2 u_{xx}, \quad (0 < x < \ell),$$

with Dirichlet boundary conditions  $u(0, t) = u(\ell, t) = 0$ . We seek solutions that can be written as a product

$$u(x, t) = X(x)T(t)$$

for some unknown functions  $X$  and  $T$ . Inserting this into the PDE and collecting terms, we see that

$$\frac{T''(t)}{c^2 T(t)} = \frac{X''(x)}{X(x)} =: -\lambda,$$

and deduce that  $\lambda$  can depend neither on  $x$  nor on  $t$ . We obtain a system of two ODE:

$$-X''(x) = \lambda X(x), \quad -T''(t) = c^2 \lambda T(t).$$

Enforcing the Dirichlet boundary conditions  $X(0) = X(\ell) = 0$ , we conclude that

$$X(x) = \sin \beta x, \quad \lambda = \beta^2, \quad T(t) = A \cos(c\beta t) + B \sin(c\beta t),$$

where  $\beta = n\pi/\ell$  for some integer  $n \geq 1$ , and  $A$  and  $B$  are constants. We have found the special solutions

$$u_n(x, t) = \cos\left(\frac{N\pi ct}{\ell}\right) \sin\left(\frac{n\pi x}{\ell}\right), \quad v_n(x, t) = \sin\left(\frac{N\pi ct}{\ell}\right) \sin\left(\frac{n\pi x}{\ell}\right).$$

Two questions remain:

- Can we construct the **general solution** of the wave equation from the  $u_n$  and  $v_n$  by superposition?
- How can we determine the coefficients in the superposition from **initial values**?

## Hand-in (due Monday, November 7):

Read Chapter 4.

26. A metal rod ( $0 < x < \ell$ ) insulated along its sides but not its ends is initially at a constant temperature  $u_0$ . Suddenly both ends are plunged into a bath of temperature zero.
- Write the initial-value problem for the temperature.
  - Use the formula

$$1 = \frac{4}{\pi} \sum_{k=1}^{\infty} \frac{1}{2k+1} \sin\left(\frac{2\pi(k+1)x}{\ell}\right), \quad (0 < x < \ell)$$

to represent the solution  $u(x, t)$  as a series.

27. Apply separation of variables to the Schrödinger equation  $iu_t = u_{xx}$  with Dirichlet boundary conditions on  $0 < x < 1$ . Here,  $i$  is the imaginary unit, satisfying  $i^2 = -1$ . You may find Euler's formula useful:  $e^{a+ib} = e^a(\cos b + i \sin b)$ .

28. Consider the PDE

$$u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta}, \quad (r > 0, \theta \in \mathbb{R})$$

with periodic boundary conditions in  $\theta$ , i.e.,  $u(r\theta + 2\pi) = u(r, \theta)$  for all  $\theta$ .

- Set  $u(r, \theta) = f(r)g(\theta)$  and separate variables to obtain a pair of ODE for  $f$  and  $g$ .
- Solve these ODE to obtain special solutions for the PDE. (*Hint:* Try  $f(r) = r^\alpha$ .)

*Remark:* We will see later this year that this is Laplace's equation in polar coordinates.

29. Find all eigenvalues of the linear operator  $Au = 2u_{xxxx}$  on the space of  $2\pi$ -periodic functions.
30. Use separation of variables to find solutions of the diffusion problem  $u_t = ku_{xx}$  with mixed boundary conditions  $u(0, t) = u_x(\ell, t) = 0$ .

## Please remember:

Our first midterm test is scheduled for **Wednesday November 4**, in-class.