

MAT 351: Partial Differential Equations

Assignment 8, due November 21, 2016

Summary

A **Hilbert space** is a vector space \mathcal{H} over \mathbb{C} with an **inner product** $\langle f, g \rangle$ that is

- **linear** in the first slot: $\langle a_1 f_1 + a_2 f_2, g \rangle = a_1 \langle f_1, g \rangle + a_2 \langle f_2, g \rangle$ for $a_1, a_2 \in \mathbb{C}$
- **Hermitian:** $\langle f, g \rangle = \overline{\langle g, f \rangle}$
- **positive definite:** $\langle f, f \rangle \geq 0$, with equality only for $f = 0$

such that \mathcal{H} is **complete** under the norm $\|f\| = (\langle f, f \rangle)^{\frac{1}{2}}$, in the sense that every Cauchy sequence in \mathcal{H} converges to a limit in \mathcal{H} .

Hilbert spaces share many geometric properties of Euclidean space, such as the **Schwarz inequality** $|\langle f, g \rangle| \leq \|f\| \|g\|$ and the **parallelogram identity** $\|f + g\|^2 + \|f - g\|^2 = 2(\|f\|^2 + \|g\|^2)$. The most important examples are the finite-dimensional complex vector spaces \mathbb{C}^m with inner product $u \cdot v$, and the function space $L^2(a, b)$ with inner product $\int_a^b f(x) \bar{g}(x) dx$.

Two vectors $f, g \in \mathcal{H}$ are **orthogonal**, if $\langle f, g \rangle = 0$. In that case, we write $f \perp g$. We have

- **Pythagoras:** If $f \perp g$, then $\|f + g\|^2 = \|f\|^2 + \|g\|^2$,

just as in \mathbb{R}^m . If X_1, X_2, \dots is a (finite or countable) sequence of orthogonal vectors in \mathcal{H} ,

- **Bessel's inequality** $\|f\|^2 \geq \sum_n |a_n|^2 \|X_n\|^2$, where $a_n = \frac{\langle f, X_n \rangle}{\|X_n\|^2}$

follows from the fact that $f - \sum_n a_n X_n$ is orthogonal to $\sum_n a_n X_n$. For the partial sums

$$S_N = \sum_{n=1}^N a_n X_n, \quad \|S_N\|^2 = \sum_{n=1}^N |a_n|^2 \|X_n\|^2,$$

Bessel's inequality implies that $\|S_N\|^2$ converges to $\sum_{n=1}^{\infty} |a_n|^2 \|X_n\|^2$, and by the Cauchy criterion S_N converges to $S = \sum_{n=1}^{\infty} a_n X_n$.

We say that the $\{X_n\}$ is an **orthogonal basis** for \mathcal{H} , if every $f \in \mathcal{H}$ can be written as a convergent series

$$f = \sum_n a_n X_n.$$

In that case, we also say that the orthogonal sequence is **complete**. We will prove that $\{\sin nx\}_{n \geq 1}$ and $\{\cos nx\}_{n \geq 0}$ are orthogonal bases for $L^2(0, \pi)$, and that $\{e^{inx}\}_{n \in \mathbb{Z}}$ is an orthogonal basis for $L^2(-\pi, \pi)$. In particular, each of the classical Fourier series of an L^2 -function f converges in L^2 to f . (We say that the Fourier series **represents** the function.) A more subtle question is under what conditions a Fourier series converges pointwise or even uniformly to f . There are examples of continuous 2π -periodic functions whose Fourier series diverges for every x !

Theorem Let X_1, X_2, \dots be a sequence of orthogonal vectors. The following are equivalent:

- (1) Finite linear combinations $\sum_{n=1}^N b_n X_n$ are **dense** in \mathcal{H} ;
- (2) If $\langle f, X_n \rangle = 0$ for all n then $f = 0$;
- (3) **Parseval's identity:** For each $f \in \mathcal{H}$, $\|f\|^2 = \sum_n |a_n|^2 \|X_n\|^2$, where $a_n = \frac{\langle f, X_n \rangle}{\|X_n\|^2}$;
- (4) $\{X_n\}_{n \geq 1}$ is an orthogonal basis.

Assignments:

Read Chapter 5 of Strauss.

31. (a) Find a pair a pair of ODE for X and Y such that $u(x, y) = X(x)Y(y)$ solves the PDE

$$u\Delta u = |\nabla u|^2, \quad (x, y \in \mathbb{R}).$$

(Do not try to solve these equations).

(b) Can you find the general solution by superposition of such product form solutions? Please explain!

32. (a) On the interval $[-1, 1]$, show that the function x is orthogonal to the constant functions.
 (b) Find a quadratic polynomial that is orthogonal to both 1 and x .
 (c) Find a cubic polynomial that is orthogonal to all quadratics.
 (These are the first three Legendre polynomials.)
33. Let ϕ be a 2π -periodic function with Fourier series $\phi(x) = \sum_n A_n e^{inx}$.
 (a) If ϕ is real-valued, show that $A_{-n} = \bar{A}_n$.
 (b) If, additionally, ϕ is even, what can you say about the Fourier coefficients? Use this to represent ϕ as a cosine series.
 (c) What if ϕ is odd?
34. (a) Find the Fourier sine series of the function $f(x) = x$ on $[0, \pi]$.
 (b) Apply Parseval's identity to compute $\sum_{n=1}^{\infty} \frac{1}{n^2}$.
 (c) Integrate the sine series term by term to obtain a Fourier cosine series for the function $\frac{1}{2}x^2$. Note that the constant of integration appears as the $n = 0$ term in the series.
 (d) Then by setting $x = 0$, find the sum of the series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2}$.
35. Let γ_n be a sequence of constants with $\lim_{n \rightarrow \infty} \gamma_n = \infty$. Define a sequence of functions on $[0, 1]$ by $f_n(x) = \gamma_n \sin(n\pi x)$ for $0 \leq x \leq \frac{1}{n}$, and $f_n(x) = 0$ otherwise.
 (a) Show that $f_n \rightarrow 0$ pointwise, but not uniformly.
 (b) If $\gamma_n = n^{1/3}$, prove that $f_n \rightarrow 0$ in L^2 .
 (c) If $\gamma_n = n^{2/3}$, show that f_n does not converge in L^2 .
36. Let f be a smooth 2π -periodic function with $\int_{-\pi}^{\pi} f(x) dx = 0$. Use the Fourier series representation and Parseval's identity to show that $\|f\| \leq \|f'\|$.