

# APM 351: Differential Equations in Mathematical Physics

## Assignment 1, due Sept. 22, 2011)

### Summary

A **partial differential equation** (PDE) is an equation of the form

$$F(x, u, Du, \dots, D^k u) = 0, \quad (1)$$

where

- $u = u(x_1, \dots, x_n)$  is the unknown function, also called the **dependent variable**; ( $u$  and  $F$  could be vector-valued; in that case, Eq. (1) is called a **system** of PDE);
- the integer  $k \geq 1$  is called the **order** of the PDE;
- $D^k u$  means all the  $k$ -th order partial derivatives of  $u$ ; and
- the **independent variables**  $x = (x_1, \dots, x_n)$  range over some open set  $D \subset \mathbb{R}^n$ .

A **solution** of the PDE is a function  $u(x_1, \dots, x_n)$  that satisfies Eq. (1) for all  $x \in D$ .

We will mostly consider first- and second order equations in two, three, or four variables. Ideally, we would like to represent the solution of a given PDE explicitly in terms of its boundary values. It turns out that this is possible only for a small number of classical equations. Among these are the **transport equation**  $u_t + bu_x = 0$ , **Poisson's equation**  $u_{xx} + u_{yy} = f(x, y)$ , the **heat equation** or **diffusion equation**  $u_t = u_{xx}$ , the **wave equation**  $u_{tt} - u_{xx} = 0$ , and the **Schrödinger equation**  $iu_t + u_{xx} - V(x)u = 0$ , all of which are fundamental in Physics. These equations are all **linear**, i.e., they can be written as

$$Lu = f,$$

where  $L$  is a linear differential operator and  $f$  is a given function.

Almost nothing can be said about a general non-linear PDE. Fundamental questions are:

- Does there **exist** a solution for a given PDE?
- Under what additional boundary conditions is the solution **unique**?
- Does the solution **depend continuously on the data**?

If the answer is “Yes!” to all three questions, then the boundary-value problem is **well-posed**. Well-posedness is crucial, if one wants to evaluate the solution of a PDE numerically, because in an ill-posed problem, even small discretization errors can have a devastating effect. But there is no reason for a general PDE to be well-posed. There even is a famous example of a linear PDE that has no solutions!

## Assignments:

Read Chapter 1 of Strauss.

1. (*Classification of PDE*)

For each of the following PDE, state the order and whether it is nonlinear, linear inhomogeneous, or linear homogeneous; provide reasons.

(a)  $u_t - u_{xx} + 1 = 0$ ;

(b)  $u_x - u_y^2 = 0$ ;

(c)  $u_x + e^y u_y = 0$ ;

(d)  $u_t + uu_x + u_{xxx} = 0$ ;  
(Kortevég-de-Vries equation)

(e)  $u_{xx} + u_{yy} + u_{zz} = \lambda u$ , where  $\lambda$  is a constant;  
(Eigenvalue problem for the Laplacian)

(f)  $u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} = 0$ ;  
(Laplace's equation in polar coordinates)

(g)  $u_x u_y = 0$ ;

(h)  $u_t + (u_x)^2 = 0$ ;  
(a Hamilton-Jacobi equation)

(h)  $u_t + (u^2)_x = 0$ ;  
(a conservation law)

2. (a) Show that there exists a unique solution for the system

$$\begin{aligned}u_x &= 3x^2y + y \\u_y &= x^3 + x\end{aligned}$$

together with the initial value  $u(0, 0) = 0$ .

(b) What is the general solution of this system of PDE?

(c) Prove that the system

$$\begin{aligned}u_x &= 2.99x^2y + y \\u_y &= x^3 + x\end{aligned}$$

has no solution at all.

3. (*Method of characteristics*)

Solve the equation  $yu_x + xu_y = 0$  with  $u(0, y) = \cos y$ . Please sketch some characteristics! In what region of the plane is the solution uniquely determined? If you enlarge the region, what fails – existence or uniqueness?