

PUTNAM PROBLEMS

ALGEBRA

Putnam problems

2007-A-1. Find all values of α for which the curves $y = \alpha x^2 + \alpha x + 1/24$ and $x = \alpha y^2 + \alpha y + 1/24$ are tangent to each other.

2007-A-2. A *repunit* is a positive integer whose digits in base 10 are all ones. Find all polynomials f with real coefficients such that if n is a repunit, then so is $f(n)$.

2007-B-1. Let f be a polynomial with positive integer coefficients. Prove that if n is a positive integer, then $f(n)$ divides $f(f(n) + 1)$ if and only if $n = 1$.

2007-B-4. Let n be a positive integer. Find the number of pairs P, Q of polynomials with real coefficients such that

$$(P(X))^2 + (Q(X))^2 = X^{2n} + 1$$

and $\deg P > \deg Q$.

2007-B-5. Let k be a positive integer. Prove that there exist polynomials $P_0(n), P_1(n), \dots, P_{k-1}(n)$ (which may depend on k) such that, for any integer n ,

$$\left\lfloor \frac{n}{k} \right\rfloor = P_0(n) + P_1(n) \left\lfloor \frac{n}{k} \right\rfloor + \dots + P_{k-1}(n) \left\lfloor \frac{n}{k} \right\rfloor^{k-1}.$$

2006-B-1. Show that the curve $x^3 + 3xy + y^3 = 1$ contains only one set of three distinct points A, B , and C , which are the vertices of an equilateral triangle, and find its area.

2005-A-3. Let $p(z)$ be a polynomial of degree n , all of whose zeros have absolute value 1 in the complex plane. Put $g(z) = p(z)/z^{n/2}$. Show that all zeros of $g'(z) = 0$ have absolute value 1.

2005-B-1. Find a nonzero polynomial $P(x, y)$ such that $P([a], [2a]) = 0$ for all real numbers a . (Note: $[\nu]$ is the greatest integer less than or equal to ν .)

2005-B-5. Let $P(x_1, \dots, x_n)$ denote a polynomial with real coefficients in the variables x_1, \dots, x_n , and suppose that

$$(a) \quad \left(\frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_n^2} \right) P(x_1, \dots, x_n) = 0 \quad (\text{identically})$$

and that

$$(b) \quad x_1^2 + \dots + x_n^2 \text{ divides } P(x_1, \dots, x_n).$$

Show that $P = 0$ identically.

2004-A-4. Show that for any positive integer n there is an integer N such that the product $x_1 x_2 \cdots x_n$ can be expressed identically in the form

$$x_1 x_2 \cdots x_n = \sum_{i=1}^N c_i (a_{i1} x_1 + a_{i2} x_2 + \dots + a_{in} x_n)^n$$

where the c_i are rational numbers and each a_{ij} is one of the numbers, $-1, 0, 1$.

2004-B-1. Let $P(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_0$ be a polynomial with integer coefficients. Suppose that r is a rational number such that $P(r) = 0$. Show that the n numbers

$$c_n r, c_n r^2 + c_{n-1} r, c_n r^3 + c_{n-1} r^2 + c_{n-2} r, \dots, c_n r^n + c_{n-1} r^{n-1} + \cdots + c_1 r$$

are integers.

2003-A-4. Suppose that a, b, c, A, B, C are real numbers, $a \neq 0$ and $A \neq 0$, such that

$$|ax^2 + bx + c| \leq |Ax^2 + Bx + C|$$

for all real numbers x . Show that

$$|b^2 - 4ac| \leq |B^2 - 4AC|.$$

2003-B-1. Do there exist polynomials $a(x), b(x), c(y), d(y)$ such that

$$1 + xy + x^2 y^2 = a(x)c(y) + b(x)d(y)$$

holds identically?

2003-B-4. Let

$$f(z) = az^4 + bz^3 + cz^2 + dz + e = a(z - r_1)(z - r_2)(z - r_3)(z - r_4)$$

where a, b, c, d, e are integers, $a \neq 0$. Show that if $r_1 + r_2$ is a rational number, and if $r_1 + r_2 \neq r_3 + r_4 - 4$, then $r_1 r_2$ is a rational number.

2002-A-1. Let k be a positive integer. The n th derivative of $1/(x^k - 1)$ has the form $P_n(x)/(x^k - 1)^{n+1}$ where $P_n(x)$ is a polynomial. Find $P_n(1)$.

2002-B-6. Let p be a prime number. Prove that the determinant of the matrix

$$\begin{pmatrix} x & y & z \\ x^p & y^p & z^p \\ x^{p^2} & y^{p^2} & z^{p^2} \end{pmatrix}$$

is congruent modulo p to a product of polynomials in the form $ax + by + cz$, where a, b, c are integers. (We say two integer polynomials are congruent modulo p if corresponding coefficients are congruent modulo p .)

2001-A-3. For each integer m , consider the polynomial

$$P_m(x) = x^4 - (2m + 4)x^2 + (m - 2)^2.$$

For what values of m is $P_m(x)$ the product of two nonconstant polynomials with integer coefficients?

2001-B-2. Find all pairs of real numbers (x, y) satisfying the system of equations

$$\frac{1}{x} + \frac{1}{2y} = (x^2 + 3y^2)(3x^2 + y^2)$$

$$\frac{1}{x} - \frac{1}{2y} = 2(y^4 - x^4).$$

2000-A-6. Let $f(x)$ be a polynomial with integer coefficients. Define a sequence a_0, a_1, \dots of integers such that $a_0 = 0$ and $a_{n+1} = f(a_n)$ for $n \geq 0$. Prove that if there exists a positive integer m for which $a_m = 0$, then either $a_1 = 0$ or $a_2 = 0$.

1999-A-1. Find polynomials $f(x)$, $g(x)$ and $h(x)$, if they exist, such that, for all x ,

$$|f(x)| - |g(x)| + |h(x)| = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0. \end{cases}$$

1999-A-2. Let $p(x)$ be a polynomial that is non-negative for all x . Prove that, for some k , there are polynomials $f_1(x), \dots, f_k(x)$ such that

$$p(x) = \sum_{j=1}^k (f_j(x))^2 .$$

1999-B-2. Let $P(x)$ be a polynomial of degree n such that $P(x) = Q(x)P''(x)$, where $Q(x)$ is a quadratic polynomial and $P''(x)$ is the second derivative of $P(x)$. Show that if $P(x)$ has at least two distinct roots then it must have n distinct roots. [The roots may be either real or complex.]

1997-B-4. Let $a_{m,n}$ denote the coefficient of x^n in the expansion of $(1 + x + x^2)^m$. Prove that for all $k \geq 0$,

$$0 \leq \sum_{i=0}^{\lfloor 2k/3 \rfloor} (-1)^i a_{k-i,i} \leq 1 >$$

1995-B-4. Evaluate

$$\sqrt[8]{2207 - \frac{1}{2207 - \frac{1}{2207 - \dots}}} .$$

Express your answer in the form $(a + b\sqrt{c})/d$, where a, b, c, d are integers.

1993-B-2. For nonnegative integers n and k , define $Q(n, k)$ to be the coefficient of x^k in the expansion of $(1 + x + x^2 + x^3)^n$. Prove that

$$Q(n, k) = \sum_{j=0}^n \binom{n}{j} \binom{n}{k-2j} ,$$

where $\binom{a}{b}$ is the standard binomial coefficient. (Reminder: For integers a and b with $a \geq 0$, $\binom{a}{b} = \frac{a!}{b!(a-b)!}$ for $0 \leq b \leq a$ and $\binom{a}{b} = 0$ otherwise.)