

TOPOLOGICAL FLATNESS OF ORTHOGONAL LOCAL MODELS IN THE SPLIT, EVEN CASE. I

BRIAN D. SMITHLING

ABSTRACT. Local models are schemes, defined in terms of linear algebra, that were introduced by Rapoport and Zink to study the étale-local structure of integral models of certain PEL Shimura varieties over p -adic fields. A basic requirement for the integral models, or equivalently for the local models, is that they be flat. In the case of local models for even orthogonal groups, Genestier observed that the original definition of the local model does not yield a flat scheme. In a recent article, Pappas and Rapoport introduced a new condition to the moduli problem defining the local model, the so-called spin condition, and conjectured that the resulting “spin” local model is flat. We prove a weak form of their conjecture in the split, Iwahori case, namely that the spin local model is topologically flat. An essential combinatorial ingredient is the equivalence of μ -admissibility and μ -permissibility for two minuscule cocharacters μ in root systems of type D .

1. INTRODUCTION

An important problem in the arithmetic theory of Shimura varieties is the definition and subsequent study of reasonable integral models. For certain PEL Shimura varieties with parahoric level structure at p , Rapoport and Zink [RZ] have constructed natural models over the ring of integers in the completion of the reflex field at any place lying over p . One of the most basic requirements for the models is that they be flat. The essential tool to investigate this and other questions of a local nature, also introduced in [RZ], is the *local model*: this is a scheme étale-locally isomorphic to the original model, but defined in terms of a purely linear-algebraic moduli problem, and thus — at least in principle — more amenable to direct study.

Local models for groups involving only types A and C have received much study in the past decade; see, for example, work of Pappas [P], Görtz [G1, G2, G3, G4], Haines and Ngô [HN1], Pappas and Rapoport [PR1, PR2, PR3, PR4], Krämer [K], and Arzdorf [A]. By contrast, the subject of this paper is the essential case of type D : local models for the split orthogonal similitude group GO_{2n} with Iwahori level structure.

Unfortunately, as observed by Genestier (see [PR4, §8.3, p. 560]), the local model defined in [RZ] *fails to be flat* in the orthogonal case, even when the group is split; subsequently, this scheme has come to be renamed the *naive local model* M^{naive} . Failure of flatness has also been observed, first by Pappas [P], for local models in type A and C cases for groups that split only after a ramified field extension. As in these cases, there is a “brute force” correction available to non-flatness of M^{naive} :

Date: March 12, 2010.

2010 Mathematics Subject Classification. Primary 14G35; Secondary 05E15; 11G18; 17B22.

Key words and phrases. Shimura variety; local model; orthogonal group; Iwahori-Weyl group; admissible set.

one simply *defines* the true local model M^{loc} to be the scheme-theoretic closure in M^{naive} of the generic fiber M_η^{naive} . A priori, this definition of M^{loc} carries the disadvantage of not admitting a ready moduli-theoretic description. Thus it is of interest when such a description can be found. In [PR4] Pappas and Rapoport propose to describe M^{loc} by adding a new condition, the so-called *spin condition* (see §2.3), to the moduli problem defining M^{naive} . We denote by M^{spin} the subscheme of M^{naive} representing Pappas's and Rapoport's strengthened moduli problem. One obtains $M^{\text{spin}} \subset M^{\text{naive}}$ as a closed subscheme, and Pappas and Rapoport show that the generic fibers of the two schemes agree. They conjecture the following.

Conjecture (Pappas-Rapoport [PR4, Conj. 8.1]). $M^{\text{spin}} = M^{\text{loc}}$, that is, M^{spin} is the scheme-theoretic closure in M^{naive} of the generic fiber.

Although the conjecture remains open in general, Pappas and Rapoport have obtained a considerable amount of computer evidence in support of it, and they explicitly work out the case $n = 1$ and part of the case $n = 2$ in [PR4]. Hand calculations in the case $n = 3$ show that M^{spin} is indeed flat with reduced special fiber. The main result of this paper is the following weak form of the conjecture.

Theorem (7.6.1). M^{spin} is topologically flat, that is, it has dense generic fiber.

In other words, the theorem asserts that the underlying topological spaces of M^{spin} and M^{loc} are the same. The strategy to prove the theorem is the same as that pioneered in Görtz's original paper [G1]: we

- (1) embed the special fiber M_k^{naive} in an appropriate affine flag variety \mathcal{F} , this time attached to GO_{2n} , over the residue field k ;
- (2) identify the set-theoretic images of M_k^{naive} and M_k^{spin} as unions of certain Schubert cells in \mathcal{F} , and obtain a good description of the Schubert cells occurring in the image of M_k^{spin} ; and
- (3) show that the Schubert cells in the image of M_k^{spin} are all in the closure of the generic fiber.

By far, (2) is the most nontrivial. The problem of obtaining a good description of the Schubert cells occurring in M_k^{spin} is essentially that of identifying the Schubert cells of maximal dimension in M_k^{spin} , since these parametrize the irreducible components of M_k^{spin} ; and this translates to a purely combinatorial problem in the Iwahori-Weyl group \widetilde{W} of GO_{2n} , which indexes the Schubert cells in \mathcal{F} . In this form, the problem becomes essentially that of μ -*permissibility vs. μ -admissibility* considered by Kottwitz-Rapoport [KR] and subsequently by Haines-Ngô [HN2]. More precisely, consider the dominant minuscule cocharacters

$$(1.1) \quad \mu_1 := (1^{(n)}, 0^{(n)}) \quad \text{and} \quad \mu_2 := (1^{(n-1)}, 0, 1, 0^{(n-1)})$$

for GO_{2n} (expressed as cocharacters for the standard diagonal torus in the ambient GL_{2n}), and regard them as translation elements in \widetilde{W} . Let W° denote the finite Weyl group of the identity component GO_{2n}° . The special fiber M_k^{spin} has two connected components, and it is easy to see that the Schubert cells corresponding to the W° -conjugates of μ_1 (resp., μ_2) are all contained in one component (resp., the other). Of course, the closures of the Schubert cells obtained in this way are again contained in M_k^{spin} . For $\mu \in \{\mu_1, \mu_2\}$, the μ -*admissible set* $\text{Adm}^\circ(\mu)$ consists of the $w \in \widetilde{W}$ whose corresponding Schubert cell C_w is contained in the closure of

$C_{\mu'}$ for some $\mu' \in \widetilde{W}^\circ \mu$. On the other hand, the condition for a given Schubert cell C_w to be contained in M_k^{spin} admits a combinatorial formulation in terms of w , and we define the μ -spin-permissible set $\text{Perm}^{\text{sp}}(\mu)$ to consist of the $w \in \widetilde{W}$ for which C_w is contained in the connected component of M_k^{spin} marked by μ . There is also a third set to consider, the μ -permissible set $\text{Perm}(\mu)$ defined in [KR].

Theorem (7.6.1, 8.8.1). *For $\mu \in \{\mu_1, \mu_2\}$, we have equalities of subsets of \widetilde{W}*

$$\text{Adm}^\circ(\mu) = \text{Perm}^{\text{sp}}(\mu) = \text{Perm}(\mu).$$

The theorem is an analog of theorems for GL_n and GSp_{2n} obtained by Kottwitz and Rapoport [KR, 3.5, 4.5, 12.4]. It is especially worth comparing with the symplectic case. Indeed, denote by $\widetilde{W}_{GL_{2n}}$ (resp., $\widetilde{W}_{GSp_{2n}}$) the Iwahori-Weyl group for GL_{2n} (resp., GSp_{2n}). Then \widetilde{W} and $\widetilde{W}_{GSp_{2n}}$ become identified under these groups' respective natural embeddings into $\widetilde{W}_{GL_{2n}}$. However, the relevant admissible and permissible sets in \widetilde{W} and $\widetilde{W}_{GSp_{2n}}$ do not agree. In the symplectic case, Kottwitz and Rapoport show these sets are obtained by intersecting $\widetilde{W}_{GSp_{2n}}$ with the relevant sets in $\widetilde{W}_{GL_{2n}}$, so that the theorem for GSp_{2n} follows from the theorem for GL_{2n} . But there seems to be no such royal road in the orthogonal case. To prove our theorem, we go back to Kottwitz's and Rapoport's original argument for GL_n and adapt it to the orthogonal setting, where some new subtleties arise.

Kottwitz and Rapoport define μ -admissibility and μ -permissibility for any cocharacter μ in any extended affine Weyl group attached to a root datum, and they show that μ -admissibility always implies μ -permissibility. However, Haines and Ngô [HN2, 7.2] have shown that the reverse implication does not hold in general. On the other hand, motivated by considerations arising from Shimura varieties, Rapoport [R, §3, p. 283] has raised the question of whether μ -admissibility and μ -permissibility agree for *minuscule* cocharacters μ , or even for sums of dominant minuscule cocharacters. In the particular setting of this paper, \widetilde{W} contains exactly three dominant minuscule cocharacters modulo the subgroup $\mathbb{Z} \cdot (1, \dots, 1)$: μ_1 , μ_2 , and

$$\mu_3 := (1, 0^{(2n-2)}, -1).$$

We give a proof of the equality $\text{Adm}^\circ(\mu_3) = \text{Perm}(\mu_3)$ in [S1]. Results of Kottwitz-Rapoport [KR], this paper, and [S1] combine to answer Rapoport's question in the affirmative for all minuscule μ in root data involving only types A , B , C , and D . By contrast, we shall show in [S3] that the answer to the more optimistic question, namely whether μ -admissibility and μ -permissibility are equivalent for μ a sum of dominant minuscule cocharacters, can be negative.

Somewhat surprisingly, Pappas and Rapoport have discovered that a version of the spin condition also turns up in their study of local models for ramified, quasi-split GU_n [PR4]. We shall show that the "spin" local models they define are topologically flat in [S2, S3].

For simplicity, in this paper we focus solely on the case of Iwahori level structure, and we ignore the explicit connection between local models and Shimura varieties — although this is certainly our main source of motivation to study local models. We intend to take up the case of general parahoric level structure, as well as the connection to Shimura varieties, in a subsequent paper.

Then $\widehat{\Lambda}$ is an \mathcal{O} -lattice in V , and h restricts to a perfect \mathcal{O} -bilinear pairing

$$(2.1.2) \quad \Lambda \times \widehat{\Lambda} \longrightarrow \mathcal{O}.$$

Given a nonempty collection \mathcal{L} of lattices in V , we say that \mathcal{L} is

- *periodic* if $a\Lambda \in \mathcal{L}$ for all $\Lambda \in \mathcal{L}$ and $a \in F^\times$;
- *self-dual* if $\widehat{\Lambda} \in \mathcal{L}$ for all $\Lambda \in \mathcal{L}$; and
- a *chain* if the lattices in \mathcal{L} are totally ordered under inclusion.

We say that a periodic lattice chain is *complete* if all successive quotients are k -vector spaces of dimension 1.

For $i = 2nk + r$ with $0 \leq r \leq 2n - 1$, we define the \mathcal{O} -lattice

$$(2.1.3) \quad \Lambda_i := \sum_{j=1}^r \pi^{-k-1} \mathcal{O}e_j + \sum_{j=r+1}^{2n} \pi^{-k} \mathcal{O}e_j \subset V.$$

Then $\widehat{\Lambda}_i = \Lambda_{-i}$ for all i , and the Λ_i 's form a complete, periodic, self-dual lattice chain Λ_\bullet , which we call the *standard chain*,

$$\cdots \subset \Lambda_{-2} \subset \Lambda_{-1} \subset \Lambda_0 \subset \Lambda_1 \subset \Lambda_2 \subset \cdots.$$

Let $f_i: \mathcal{O}^{2n} \rightarrow \mathcal{O}^{2n}$ multiply the i th standard basis element by π and send all other standard basis elements to themselves. Then there is a unique isomorphism of chains of \mathcal{O} -modules

$$(2.1.4) \quad \begin{array}{ccccccc} \cdots & \hookrightarrow & \Lambda_0 & \hookrightarrow & \Lambda_1 & \hookrightarrow & \cdots & \hookrightarrow & \Lambda_{2n} & \hookrightarrow & \cdots \\ & & \uparrow \sim & & \uparrow \sim & & & & \uparrow \sim & & \\ \cdots & \xrightarrow{f_{2n}} & \mathcal{O}^{2n} & \xrightarrow{f_1} & \mathcal{O}^{2n} & \xrightarrow{f_2} & \cdots & \xrightarrow{f_{2n}} & \mathcal{O}^{2n} & \xrightarrow{f_1} & \cdots \end{array}$$

such that the leftmost vertical arrow identifies the standard ordered basis of \mathcal{O}^{2n} with the ordered \mathcal{O} -basis e_1, \dots, e_{2n} of Λ_0 .

2.2. Naive local models. In this subsection we recall the definition of naive local models from Rapoport's and Zink's book [RZ] in the orthogonal case. Given an \mathcal{O} -module M and an \mathcal{O} -scheme S , we write M_S for the quasi-coherent \mathcal{O}_S -module $M \otimes_{\mathcal{O}} \mathcal{O}_S$.

Let \mathcal{L} be a periodic self-dual lattice chain in V . The *naive local model* $M_{\mathcal{L}}^{\text{naive}}$ attached to \mathcal{L} is the following contravariant functor on the category of \mathcal{O} -schemes. Given an \mathcal{O} -scheme S , an S -point in $M_{\mathcal{L}}^{\text{naive}}$ consists of, up to an obvious notion of isomorphism,

- a functor

$$\begin{array}{ccc} \mathcal{L} & \longrightarrow & (\mathcal{O}_S\text{-modules}) \\ \Lambda & \longmapsto & \mathcal{F}_\Lambda, \end{array}$$

where \mathcal{L} is regarded as a category in the obvious way; together with

- an injection $\mathcal{F}_\Lambda \hookrightarrow \Lambda_S$ for each $\Lambda \in \mathcal{L}$, functorial in Λ ;

satisfying, for all $\Lambda \in \mathcal{L}$,

- (LM1) \mathcal{F}_Λ embeds in Λ_S as an \mathcal{O}_S -locally direct summand of rank n ;
- (LM2) the isomorphism $\Lambda_S \xrightarrow{\sim} (\pi\Lambda)_S$ obtained by tensoring $\Lambda \xrightarrow{\sim} \pi\Lambda$ identifies \mathcal{F}_Λ with $\mathcal{F}_{\pi\Lambda}$; and

(LM3) the perfect \mathcal{O}_S -bilinear pairing $\Lambda_S \times \widehat{\Lambda}_S \rightarrow \mathcal{O}_S$ obtained by tensoring (2.1.2) identifies $\mathcal{F}_\Lambda^\perp \subset \widehat{\Lambda}_S$ with $\widetilde{\mathcal{F}}_\Lambda$, where for any \mathcal{O}_S -submodule $M \subset \Lambda_S$, $M^\perp \subset \widehat{\Lambda}_S$ is the subsheaf of sections that pair to 0 with all sections of M .

The functor $M_{\mathcal{L}}^{\text{naive}}$ is plainly represented by a closed subscheme, which we again denote $M_{\mathcal{L}}^{\text{naive}}$, of a finite product of Grassmannians over $\text{Spec } \mathcal{O}$.

If π is invertible on S , then any inclusion $\Lambda \subset \Lambda'$ of \mathcal{O} -lattices becomes an isomorphism after tensoring with \mathcal{O}_S . Hence, for such S , any S -point of $M_{\mathcal{L}}^{\text{naive}}$ is determined by $\mathcal{F}_\Lambda \hookrightarrow \Lambda_S$ for any single $\Lambda \in \mathcal{L}$. Hence $M_{\mathcal{L}}^{\text{naive}}$ has generic fiber $\text{OGr}(n, 2n)_F$, the orthogonal Grassmannian of totally isotropic n -planes in $2n$ -space; this is a smooth $\binom{n}{2}$ -dimensional scheme with two components, each isomorphic to $SO(h)/P$, where $P \subset SO(h)$ is a parabolic subgroup stabilizing some totally isotropic n -plane.

In this paper we restrict to the Iwahori case, that is, to local models attached to *complete* lattice chains. It is not hard to verify directly that the special orthogonal group $SO(h)(F)$ acts transitively on the complete periodic self-dual lattice chains in V . Hence the local models attached to any two complete lattice chains are isomorphic. We shall work with respect to the standard chain Λ_\bullet , and we abbreviate $M^{\text{naive}} := M_{\Lambda_\bullet}^{\text{naive}}$.

The chain isomorphism (2.1.4) permits a very concrete description of the points of M^{naive} : an R -point consists of R -submodules $\mathcal{F}_0, \mathcal{F}_1, \dots, \mathcal{F}_{2n} \subset R^{2n}$, each a locally direct summand of rank n , such that $(f_i \otimes R)(\mathcal{F}_{i-1}) \subset \mathcal{F}_i$ for all $i = 1, \dots, 2n$; $\mathcal{F}_0 = \mathcal{F}_{2n}$; and $\mathcal{F}_i^\perp = \mathcal{F}_{2n-i}$ for all $i = 1, \dots, 2n$, where R^{2n} carries the split symmetric form having matrix (2.1.1) with respect to its standard basis.

2.3. The spin condition of Pappas and Rapoport. In [PR4], Pappas and Rapoport introduce a conjectural correction to the non-flatness of $M_{\mathcal{L}}^{\text{naive}}$ by adding a new constraint, the *spin condition*, to the moduli problem. They define the spin condition in the case of an arbitrary nondegenerate symmetric bilinear form h on V . We are only concerned in this paper with the case that h is *split*. The formulation of the spin condition simplifies a bit in the split case: namely, we can get by without explicit use of the *discriminant algebra* of [PR4, §7.1]. It is a simple exercise to check that the formulation of the spin condition we're about to give is equivalent to the spin condition in [PR4] in the split case.

To formulate the spin condition, we shall recall only the bare minimum of linear algebra we need from [PR4, §7]. In particular, we refer to [PR4] for a more expansive and satisfying version of the following discussion.

For a subset $E \subset \{1, \dots, 2n\}$ of cardinality n , set

$$(2.3.1) \quad e_E := e_{j_1} \wedge \cdots \wedge e_{j_n} \in \bigwedge_F^n V,$$

where $E = \{j_1, \dots, j_n\}$ with $j_1 < \cdots < j_n$. Given such E , we also set

$$(2.3.2) \quad E^* := 2n + 1 - E \quad \text{and} \quad E^\perp := (E^*)^c = (E^c)^*,$$

where the set complements are taken in $\{1, \dots, 2n\}$. Then E^* specifies the indices j' such that $h(e_j, e_{j'}) = 1$ for some $j \in E$, and E^\perp specifies the j' such that $h(e_j, e_{j'}) = 0$ for all $j \in E$.

We define an operator a on $\bigwedge^n V$ by its action on the standard basis elements e_E for varying E ,

$$a(e_E) := \text{sgn}(\sigma_E) e_{E^\perp},$$

where σ_E is the permutation on $\{1, \dots, 2n\}$ sending $\{1, \dots, n\}$ to the elements of E^* in decreasing order, and sending $\{n+1, \dots, 2n\}$ to the elements of E^\perp in increasing order. Then a satisfies $a^2 = \text{id}_{\bigwedge^n V}$ [PR4, Prop. 7.1]. Hence $\bigwedge^n V$ decomposes as

$$\bigwedge^n V = \left(\bigwedge^n V \right)_+ \oplus \left(\bigwedge^n V \right)_-,$$

where $\left(\bigwedge^n V \right)_\pm$ denotes the ± 1 eigenspace for a . Using that a^2 is the identity, we see that

$$(2.3.3) \quad \left(\bigwedge^n V \right)_\pm = \text{span}_F \{e_E \pm \text{sgn}(\sigma_E) e_{E^\perp}\},$$

where E ranges through the subsets of $\{1, \dots, 2n\}$ of cardinality n .

Now let $\Lambda \subset V$ be an \mathcal{O} -lattice. Then $\bigwedge_{\mathcal{O}}^n \Lambda$ is naturally an \mathcal{O} -submodule of $\bigwedge_F^n V$, and we set

$$\left(\bigwedge_{\mathcal{O}}^n \Lambda \right)_\pm := \left(\bigwedge_{\mathcal{O}}^n \Lambda \right) \cap \left(\bigwedge_F^n V \right)_\pm.$$

We are now ready to state the spin condition. Let \mathcal{L} be a periodic self-dual lattice chain. We say that an S -point $\{\mathcal{F}_\Lambda \hookrightarrow \Lambda_S\}_{\Lambda \in \mathcal{L}}$ of $M_{\mathcal{L}}^{\text{naive}}$ satisfies the *spin condition* if

(LM4) Zariski locally on S , either $\bigwedge_{\mathcal{O}_S}^n \mathcal{F}_\Lambda$ is contained in

$$\text{im} \left[\left(\bigwedge_{\mathcal{O}}^n \Lambda \right)_+ \otimes_{\mathcal{O}} \mathcal{O}_S \longrightarrow \bigwedge_{\mathcal{O}_S}^n \Lambda_S \right]$$

for all Λ in \mathcal{L} , or in

$$\text{im} \left[\left(\bigwedge_{\mathcal{O}}^n \Lambda \right)_- \otimes_{\mathcal{O}} \mathcal{O}_S \longrightarrow \bigwedge_{\mathcal{O}_S}^n \Lambda_S \right]$$

for all Λ in \mathcal{L} .

The *spin local model attached to \mathcal{L}* , which we denote $M_{\mathcal{L}}^{\text{spin}}$, is the closed subscheme of $M_{\mathcal{L}}^{\text{naive}}$ whose points satisfy the spin condition. Pappas and Rapoport show in [PR4, §8.2.1] that the arrow $(M_{\mathcal{L}}^{\text{spin}})_F \hookrightarrow (M_{\mathcal{L}}^{\text{naive}})_F$ on generic fibers is an isomorphism.

As in the previous subsection, when working with complete periodic self-dual chains \mathcal{L} , $M_{\mathcal{L}}^{\text{spin}}$ is independent of \mathcal{L} up to isomorphism, and we put $M^{\text{spin}} := M_{\Lambda_\bullet}^{\text{spin}}$.

3. ORTHOGONAL SIMILITUDE GROUP

In this section we review some basic facts about split GO_{2n} . We switch to working over the field K . Except in §3.4, K may be an arbitrary field of characteristic not 2; in §4 we'll return to our blanket assumptions on K stated in the introduction.

3.1. Orthogonal similitudes. Abusing notation, we denote again by h the symmetric bilinear form on K^{2n} whose matrix with respect to the standard ordered basis is (2.1.1). We denote by $G := GO_{2n} := GO(h)$ the algebraic group over K of *orthogonal similitudes* of h : for any K -algebra R , $G(R)$ is the set of elements $g \in GL_{2n}(R)$ satisfying $h_R(gx, gy) = c(g)h_R(x, y)$ for some $c(g) \in R^\times$ and all $x, y \in R^{2n}$, where h_R is the induced form on R^{2n} . As the form h is nonzero, the scalar $c(g)$ is uniquely determined, and c defines an exact sequence of K -groups

$$1 \longrightarrow O \longrightarrow G \xrightarrow{c} \mathbb{G}_m \longrightarrow 1$$

with evident kernel $O := O_{2n} := O(h)$ the orthogonal group of h . The displayed sequence splits (noncanonically), so that the choice of a splitting presents G as a semidirect product $O \rtimes \mathbb{G}_m$.

3.2. Center. The center $Z := Z_G$ of G consists of the scalar matrices; on R -valued points,

$$Z(R) = \{ r \cdot \text{id} \in GL_{2n}(R) \mid r \in R^\times \},$$

where id denotes the identity matrix, so that $Z \cong \mathbb{G}_m$.

We write $G_{\text{ad}} := PGO_{2n} := PGO(h) := G/Z$ for the adjoint group.

3.3. Connected components. The group G possesses two connected components. For $g \in G(R)$ with $\text{Spec } R$ connected, the corresponding morphism $\text{Spec } R \rightarrow G$ factors through the identity component or the non-identity component according as $c(g)^{-n} \det(g)$ is $+1$ or -1 , respectively. The identity component G° is split reductive.

3.4. Standard maximal torus. Let T denote the standard split maximal torus of diagonal matrices in G ; on R -points,

$$T(R) := \{ \text{diag}(a_1, \dots, a_{2n}) \in GL_{2n}(R) \mid a_1 a_{2n} = a_2 a_{2n-1} = \dots = a_n a_{n+1} \},$$

so that $T \simeq \mathbb{G}_m^{n+1}$.

Now let us assume K is as in the introduction, so that it is discretely valued with valuation ring \mathcal{O}_K and uniformizer t . Then we identify the cocharacter lattice $X_*(T)$ with $T(K)/T(\mathcal{O}_K)$ via the rule $\lambda \mapsto \lambda(t) \bmod T(\mathcal{O}_K)$, and we identify $T(K)/T(\mathcal{O}_K)$ with

$$(*) \quad \{ (r_1, \dots, r_{2n}) \in \mathbb{Z}^{2n} \mid r_1 + r_{2n} = \dots = r_n + r_{n+1} \}$$

via ord . These identifications in turn identify

- $X_*(T_{\text{der}})$ with the subgroup of $(*)$ of elements (r_1, \dots, r_{2n}) such that $r_1 + r_{2n} = \dots = r_n + r_{n+1} = 0$, where $G_{\text{der}} := SO := SO_{2n} := SO(h)$ is the derived group of G and T_{der} is its split maximal torus $T \cap G_{\text{der}}$; and
- $X_*(T_{\text{ad}})$ with the quotient of $(*)$ by the subgroup $\mathbb{Z} \cdot (1, \dots, 1)$, where $T_{\text{ad}} := T/Z$ is the split maximal torus in G_{ad} obtained as the image of T .

3.5. Roots, coroots. Let χ_i denote the character on T sending

$$\text{diag}(a_1, \dots, a_{2n}) \mapsto a_i.$$

The roots of the pair (G, T) are the set

$$\begin{aligned} \Phi_G &:= \Phi_{G,T} := \{ \pm(\chi_i - \chi_j) \mid 1 \leq i < j < i^* \} \\ &= \{ \pm(\chi_i - \chi_j), \pm(\chi_i + \chi_j - c) \mid 1 \leq i < j \leq n \}, \end{aligned}$$

where we use the same symbol c to denote the composite $T \hookrightarrow G \xrightarrow{c} \mathbb{G}_m$. Of course, the roots of G descend to the roots $\Phi_{G_{\text{ad}}} := \Phi_{G_{\text{ad}}, T_{\text{ad}}}$ of the pair $(G_{\text{ad}}, T_{\text{ad}})$. When $n = 1$, $G^\circ \simeq \mathbb{G}_m^2$ is abelian and $\Phi_G = \Phi_{G_{\text{ad}}} = \emptyset$. Otherwise, the root system $\Phi_{G_{\text{ad}}}$ is of type $A_1 \times A_1$ for $n = 2$, A_3 for $n = 3$, and D_n for $n \geq 4$.

For $n > 1$, we take the n roots

$$(3.5.1) \quad \chi_1 - \chi_2, \dots, \chi_{n-1} - \chi_n, \chi_{n-1} + \chi_n - c$$

as simple roots.

For $1 \leq i \leq n$, let $\lambda_i \in X_*(T)$ denote the cocharacter

$$x \mapsto \text{diag}(1, \dots, 1, x, 1, \dots, 1, x^{-1}, 1, \dots, 1),$$

where x is in the i th slot and x^{-1} is in the i^* th slot. Then for $n > 1$, the *coroots* consist of the cocharacters

$$(\chi_i - \chi_j)^\vee = \lambda_i - \lambda_j \quad \text{and} \quad (\chi_i + \chi_j - c)^\vee = \lambda_i + \lambda_j$$

for $1 \leq i < j \leq n$.

3.6. Weyl group. The torus T has *normalizer* $N := N_G T$ in G the algebraic group of monomial matrices contained in G , and *finite Weyl group*

$$W := W_{G,T} := N(K)/T(K).$$

The Weyl group W acts naturally on the set of lines in K^{2n} spanned by the standard ordered basis vectors, and this canonically identifies W with the group S_{2n}^h of permutations σ of $\{1, \dots, 2n\}$ satisfying

$$\sigma(i^*) = \sigma(i)^* \quad \text{for all } i.$$

The group S_{2n}^h decomposes as a semidirect product $\{\mathbb{Z}/2\mathbb{Z}\}^n \rtimes S_n$, where the non-trivial element in the i th copy of $\mathbb{Z}/2\mathbb{Z}$ acts as the transposition (i, i^*) , and where the symmetric group S_n acts on $\{1, \dots, n\}$ in the standard way and on $\{n+1, \dots, 2n\}$ in the way compatible with the display.

Note that W is not the Weyl group attached to the root system Φ_G . Rather, let

$$(3.6.1) \quad W^\circ := W_{G^\circ, T} := N_{G^\circ(F)} T(K)/T(K)$$

denote the finite Weyl group of T in G° . Then $W^\circ \cong W(\Phi_G)$ is naturally contained in W as a subgroup of index 2. In terms of permutations, W° corresponds to the elements of S_{2n}^h which are *even* as elements of the symmetric group S_{2n} .

3.7. Fundamental group. In terms of the identifications in §3.4, the *coroot lattice*

$$(3.7.1) \quad Q^\vee := Q^\vee(G, T) \subset X_*(T_{\text{der}}) \subset X_*(T)$$

consists of all $(r_1, \dots, r_{2n}) \in \mathbb{Z}^{2n}$ such that $r_1 + r_{2n} = \dots = r_n + r_{n+1} = 0$ and $r_1 + \dots + r_n$ is even. The *fundamental group* of G is the fundamental group of the identity component G° ,

$$\pi_1(G) := \pi_1(G^\circ) := X_*(T)/Q^\vee \simeq \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}.$$

Note that the derived group $G_{\text{der}} = G_{\text{der}}^\circ = SO$ is not simply connected, as its fundamental group $X_*(T_{\text{der}})/Q^\vee \cong \mathbb{Z}/2\mathbb{Z}$.

4. IWAHORI SUBGROUP

We return to our assumptions on K stated in the introduction. In this section we discuss the standard Iwahori subgroup of $G(K)$. In particular, we realize it as a lattice chain stabilizer.

4.1. Standard apartment. Let $\mathcal{B} := \mathcal{B}(G_{\text{ad}})$ denote the building of G_{ad} . We call the apartment in \mathcal{B} associated with T_{ad} the *standard apartment*, and we denote it by $\mathfrak{a} := \mathfrak{a}_{T_{\text{ad}}}$. In terms of the identifications in §3.4,

$$\mathfrak{a} = X_*(T_{\text{ad}}) \otimes_{\mathbb{Z}} \mathbb{R} \cong \frac{\{(r_1, \dots, r_{2n}) \in \mathbb{R}^{2n} \mid r_1 + r_{2n} = \dots = r_n + r_{n+1}\}}{\mathbb{R} \cdot (1, \dots, 1)}.$$

4.2. Base alcove. We take as our base alcove the alcove A in \mathfrak{a} containing the origin and contained in the *negative* Weyl chamber relative to our choice of simple roots (3.5.1). The alcove A has $n + 1$ vertices

$$\begin{aligned} a_0 &:= (0, \dots, 0), \\ a_{0'} &:= (-1, 0^{(2n-2)}, 1), \\ a_i &:= \left(\left(-\frac{1}{2}\right)^{(i)}, 0^{(2n-2i)}, \left(\frac{1}{2}\right)^{(i)} \right), \quad 2 \leq i \leq n-2, \\ a_n &:= \left(\left(-\frac{1}{2}\right)^{(n)}, \left(\frac{1}{2}\right)^{(n)} \right), \\ a_{n'} &:= \left(\left(-\frac{1}{2}\right)^{(n-1)}, \frac{1}{2}, -\frac{1}{2}, \left(\frac{1}{2}\right)^{(n-1)} \right), \end{aligned}$$

all taken mod $\mathbb{R} \cdot (1, \dots, 1)$. The vertices a_0 , $a_{0'}$, a_n , and $a_{n'}$ are hyperspecial; the other vertices are nonspecial.

4.3. Standard Iwahori subgroup. Let us say that an *Iwahori subgroup* of $G(K)$ is just an Iwahori subgroup of $G^\circ(K)$ in the usual sense for any connected reductive group. We denote by B the Iwahori subgroup of $G(K)$ attached to our base alcove A , and we call it the *standard Iwahori subgroup*.

To realize B as a lattice chain stabilizer, let λ_\bullet denote the \mathcal{O}_K -lattice chain in K^{2n} defined as the obvious analog of the \mathcal{O} -lattice chain Λ_\bullet (2.1.3), where \mathcal{O}_K replaces \mathcal{O} and t replaces π . Let

$$P_{\lambda_\bullet} := \{ g \in G(K) \mid g\lambda_i = \lambda_i \text{ for all } i \}.$$

Then P_{λ_\bullet} is the intersection of $G(K)$ with the standard Iwahori subgroup

$$\begin{pmatrix} \mathcal{O}_K^\times & & & \mathcal{O}_K \\ & \ddots & & \\ & & \ddots & \\ t\mathcal{O}_K & & & \mathcal{O}_K^\times \end{pmatrix}$$

of $GL_{2n}(K)$.

Proposition 4.3.1. $B = P_{\lambda_\bullet}$.

To prepare for the proof, recall [HR, 3; BT, remark after 5.2.8] that for any facet F in \mathcal{B} , the associated parahoric subgroup P_F is precisely the set of all $g \in G^\circ(K)$ with trivial Kottwitz invariant such that $ga = a$ for all vertices a of F . The Kottwitz homomorphism admits a simple description for any split connected reductive group H with split maximal torus S : it is a functorial surjective homomorphism

$$\kappa_H: H(K) \twoheadrightarrow \pi_1(H)$$

which is characterized in terms of the Cartan decomposition

$$H(K) = H(\mathcal{O}_K)S(K)H(\mathcal{O}_K)$$

as being trivial on $H(\mathcal{O}_K)$ and as restricting on $S(K)$ to the composite

$$S(K) \twoheadrightarrow S(K)/S(\mathcal{O}_K) \cong X_*(S) \twoheadrightarrow X_*(S)/Q_{H,S}^\vee = \pi_1(H),$$

where $Q_{H,S}^\vee$ denotes the coroot lattice for S in H . In the case of our group G° , upon choosing a splitting $G^\circ \simeq SO \rtimes \mathbb{G}_m$, we identify its fundamental group with $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}$ in the way that κ_{G° sends $(g, x) \mapsto (\kappa_{SO}(g), \text{ord}(x))$.

Lemma 4.3.2. $P_{\lambda_\bullet} \subset G^\circ(K)$.

Proof. Given $g \in P_{\lambda_\bullet}$, we must show that $c(g)^n = \det(g)$. Since $\text{char}(k) \neq 2$ and the only other possibility is $c(g)^n = -\det(g)$, it suffices to show $c(g)^n \equiv \det(g) \pmod{t}$.

Write g as a matrix (g_{ij}) . Since g preserves the form h up to the scalar $c(g)$, the i th and i^* th columns of g pair to $c(g)$ for $1 \leq i \leq 2n$. Hence

$$c(g) \equiv g_{i,i} g_{i^*,i^*} \pmod{t}.$$

Hence

$$c(g)^n \equiv \prod_{i=1}^{2n} g_{i,i} \equiv \det(g) \pmod{t}. \quad \square$$

Proof of (4.3.1). Since plainly $P_{\lambda_\bullet} \subset G(\mathcal{O}_K)$, the lemma implies $P_{\lambda_\bullet} \subset \ker \kappa_{G^\circ}$. On the other hand, we see from the explicit form of κ_{G° that any $g \in B \subset \ker \kappa_{G^\circ}$ has determinant of valuation 0. The equality $B = P_{\lambda_\bullet}$ now follows easily from the explicit expressions for the vertices of A and from the usual identification of \mathcal{B} with homothety classes of certain norms on K^{2n} . \square

5. IWAHORI-WEYL GROUP

In this section we discuss a few matters related to the Iwahori-Weyl group of G . Once we specialize to the function field case later on, we'll use the Iwahori-Weyl group to index Schubert cells in the affine flag variety attached to G .

5.1. Iwahori-Weyl group. The *Iwahori-Weyl group* \widetilde{W} of G is the group

$$\widetilde{W} := \widetilde{W}_G := \widetilde{W}_{G,T} := N(K)/T(\mathcal{O}_K).$$

We shall also need the Iwahori-Weyl group \widetilde{W}° of the identity component G° ,

$$\widetilde{W}^\circ := \widetilde{W}_{G^\circ} := \widetilde{W}_{G^\circ,T} := N_{G^\circ(K)}T(K)/T(\mathcal{O}_K).$$

It will be convenient for us to single out the permutation matrix $\tau \in G(K)$ corresponding to the transposition $(n, n+1)$. Then τ is contained in the non-identity component of $O(K)$ and normalizes T , so that there is a decomposition $\widetilde{W} = \widetilde{W}^\circ \amalg \tau \widetilde{W}^\circ$.

5.2. Affine Bruhat decomposition. Let H be a split connected reductive K -group with split maximal torus S , and let $I \subset H(K)$ be the Iwahori subgroup corresponding to an alcove in the apartment associated with S . The *affine Bruhat decomposition* asserts that the natural map $\widetilde{W}_{H,S} := N_{H(K)}S(K)/S(\mathcal{O}_K) \rightarrow I \backslash H(K)/I$ sending $n \pmod{S(\mathcal{O}_K)} \mapsto InI$ is a bijection; see Haines and Rapoport [HR, 8]. In this subsection we show that the analogous result still holds for our disconnected group G .

Proposition 5.2.1. *The natural map $\widetilde{W} \rightarrow B \backslash G(K)/B$ is a bijection of sets.*

Proof. This follows from the affine Bruhat decomposition for G° . Indeed, we have decompositions

$$\widetilde{W} = W^\circ \amalg \tau \widetilde{W}^\circ$$

and

$$B \backslash G(K)/B = (B \backslash G^\circ(K)/B) \amalg (B \backslash \tau G^\circ(K)/B),$$

and we at least obtain $W^\circ \xrightarrow{\sim} B \backslash G^\circ(K)/B$. So it remains to show that the map $\tau \widetilde{W}^\circ \rightarrow B \backslash \tau G^\circ(K)/B$ is a bijection. Since τ plainly stabilizes the base alcove A ,

τ normalizes B . Hence $B \backslash \tau G^\circ(K) / B = \tau(B \backslash G^\circ(K) / B)$. So we get what we need again from the affine Bruhat decomposition for G° . \square

5.3. Semidirect product decompositions. As usual, \widetilde{W} admits two standard semidirect product decompositions, which we now describe.

The first decomposition is

$$\widetilde{W} \cong (T(K)/T(\mathcal{O}_K)) \rtimes W \cong X_*(T) \rtimes W,$$

where we lift the finite Weyl group W to $N(K)$ by choosing permutation matrices as representatives, and where we identify $T(K)/T(\mathcal{O}_K) \cong X_*(T)$ as in §3.4. In this way, we refer to $X_*(T)$ as the *translation subgroup* of \widetilde{W} , and we denote the image of $\mu \in X_*(T)$ in \widetilde{W} by t_μ . Concretely, in terms of our identifications for $X_*(T)$ and W in §3.4 and §3.6, respectively, we have

$$(5.3.1) \quad \widetilde{W} \cong \{ (r_1, \dots, r_{2n}) \in \mathbb{Z}^{2n} \mid r_1 + r_{2n} = \dots = r_n + r_{n+1} \} \rtimes S_{2n}^h.$$

The second decomposition involves the *affine Weyl group* W_a of G . In terms of our first semidirect product decomposition, we have $W_a := Q^\vee \rtimes W^\circ \subset \widetilde{W}$, where we recall $Q^\vee \subset X_*(T)$ is the coroot lattice (3.7.1) and $W^\circ \subset W$ is the finite Weyl group of G° (3.6.1). Then

- W_a is a normal subgroup of \widetilde{W} ; and
- W_a is canonically identified with the affine Weyl group of the root system $(\Phi_{G_{\text{ad}}}, X_*(T_{\text{ad}}) \otimes \mathbb{R})$, so that W_a acts simply transitively on the set of alcoves in the standard apartment.

Hence \widetilde{W} is the semidirect product of W_a and the stabilizer Ω of the base alcove A ,

$$\widetilde{W} \cong W_a \rtimes \Omega.$$

We remark that, in contrast with the analogous situation for a connected reductive group, the quotient $\widetilde{W}/W_a \cong \Omega$ is *nonabelian*. Indeed, we have an identification $\widetilde{W}/W_a \cong X_*(T)/Q^\vee \rtimes W/W^\circ$; and the point is that W/W° is nontrivial and acts nontrivially on $X_*(T)/Q^\vee$. To see this, recall the cocharacters $\mu_1, \mu_2 \in X_*(T)$ from (1.1); these yield distinct dominant minuscule coweights for G_{ad} . Hence μ_1 and μ_2 have distinct images in $X_*(T)/Q^\vee$. But $W/W^\circ \cong \mathbb{Z}/2\mathbb{Z}$ is generated by the image of τ , and the action of τ on $X_*(T)$ interchanges μ_1 and μ_2 .

5.4. Length, Bruhat order. The decomposition $\widetilde{W} \cong W_a \rtimes \Omega$ furnishes a *length function* and *Bruhat order* on \widetilde{W} in the standard way, which we briefly recall. The reflections through the walls of the base alcove form a generating set for the Coxeter group W_a . Hence we get a length function ℓ and Bruhat order \leq on W_a . These then extend to \widetilde{W} as usual: for $x\omega, x'\omega' \in \widetilde{W}$ with $x, x' \in W_a$ and $\omega, \omega' \in \Omega$, we have $\ell(x\omega) := \ell(x)$ and $x\omega \leq x'\omega'$ exactly when $\omega = \omega'$ and $x \leq x'$ in W_a .

We remark now that, in the function field case, the Bruhat order gives the correct closure relations for Schubert varieties in the affine flag variety; see (6.4.1) below.

5.5. μ -admissible set. Let $\mu \in X_*(T)$ be a cocharacter. Then we define the *μ -admissible set* $\text{Adm}(\mu) \subset \widetilde{W}$ in the most obvious way based on the usual definition for connected groups,

$$\text{Adm}(\mu) := \{ w \in \widetilde{W} \mid w \leq \sigma t_\mu \sigma^{-1} \text{ for some } \sigma \in W \}.$$

Of course, we in fact have $\text{Adm}(\mu) \subset \widetilde{W}^\circ$.

In the case of a connected group, all elements of the μ -admissible set are congruent mod \widetilde{W}_a since $\widetilde{W}/\widetilde{W}_a$ is abelian; in fact, as shown by Rapoport [R, 3.1], this common element in $\widetilde{W}/\widetilde{W}_a$ depends only on the geometric conjugacy class of μ . In the case of our disconnected group G , we have already seen that $\widetilde{W}/\widetilde{W}_a$ is nonabelian. And indeed, it can happen that $\text{Adm}(\mu)$ possesses elements that are *distinct* mod \widetilde{W}_a . For example, this is the case for $\mu = \mu_1$ (1.1), since $\tau\mu_1\tau^{-1} = \mu_2$.

To make this a bit more precise, consider

$$\text{Adm}^\circ(\mu) := \{ w \in \widetilde{W} \mid w \leq \sigma t_\mu \sigma^{-1} \text{ for some } \sigma \in W^\circ \},$$

the *admissible set of μ in G°* . Then for any μ ,

$$\text{Adm}(\mu) = \text{Adm}^\circ(\mu) \cup \text{Adm}^\circ(\tau\mu\tau^{-1}).$$

Hence the study of admissible sets for G reduces to the study of admissible sets for G° . We see from this last display that $\text{Adm}(\mu)$ contains either 1 or 2 elements mod \widetilde{W}_a ; the union is disjoint precisely in the latter case.

5.6. Extended alcoves. We conclude the section by giving a combinatorial description of \widetilde{W} in terms of *extended alcoves* that will be convenient later on when we consider Schubert cells in the affine flag variety. Identifying $\widetilde{W} \cong \widetilde{W}_{GS_{p_{2n}}}$ as in §5.4, our description will be the same as that for $\widetilde{W}_{GS_{p_{2n}}}$ given by Kottwitz and Rapoport in [KR, 4.2], except we shall adopt some slightly different conventions to make the relation with the affine flag variety clearer. Following the notation of [KR], given $v \in \mathbb{Z}^{2n}$, we write $v(i)$ for the i th entry of v , and we write Σv for the sum of the entries of v . We write $v \geq w$ if $v(i) \geq w(i)$ for all i .

An *extended alcove* for G is a sequence v_0, \dots, v_{2n-1} of elements in \mathbb{Z}^{2n} such that, putting $v_{2n} := v_0 - (1, \dots, 1)$,

- (A1) $v_0 \geq v_1 \geq \dots \geq v_{2n}$;
- (A2) $\Sigma v_i = \Sigma v_{i-1} - 1$ for all $1 \leq i \leq 2n$; and
- (A3) there exists $d \in \mathbb{Z}$ such that $v_i(j) + v_{2n-i}(j^*) = d$ for all $1 \leq i, j \leq 2n$.

We frequently refer to (A3) as the *duality condition*. The sequence of elements $\omega_i := ((-1)^{(i)}, 0^{(2n-i)})$ is an extended alcove, with $d = -1$, which we call the *standard extended alcove*. The group \widetilde{W} acts naturally on extended alcoves via its expression in (5.3.1). Just as in [KR, 4.2], this action is simply transitive, and we identify \widetilde{W} with the set of extended alcoves by taking the standard extended alcove as base point.

6. AFFINE FLAG VARIETY

In this section we discuss a few basic aspects of the affine flag variety attached to G in the function field case. We take $K = k((t))$ and $\mathcal{O}_K = k[[t]]$ from now on. We follow closely [PR4, §§3.1–3.2].

6.1. Affine flag variety. We recall the construction of the affine flag variety over k .

To begin, the *loop group* LG is the functor on k -algebras

$$LG: R \longmapsto G(R((t))),$$

where $R((t))$ is the ring of Laurent series with coefficients in R , regarded as a K -algebra in the obvious way.

Next recall the standard Iwahori subgroup $B \subset G(K)$. Abusing notation, we denote again by B the associated Bruhat-Tits scheme over \mathcal{O}_K ; this is a smooth affine group scheme with generic fiber G° and with connected special fiber. We denote by L^+B the functor on k -algebras

$$L^+B: R \longmapsto B(R[[t]]),$$

where $R[[t]]$ is regarded as an \mathcal{O}_K -algebra in the obvious way.

Finally, the *affine flag variety* \mathcal{F} is the fpqc quotient LG/L^+B of sheaves on the category of k -algebras. It is an ind- k -scheme of ind-finite type [PR3, 1.4]. Note that \mathcal{F} is a disjoint union of two copies of the affine flag variety $\mathcal{F}^\circ := LG^\circ/L^+B$ for G° ,

$$\mathcal{F} = \mathcal{F}^\circ \amalg \tau \mathcal{F}^\circ,$$

with $\tau \in G(K)$ the element of §5.1.

6.2. Lattice-theoretic description. In this subsection we describe points on the affine flag variety in terms of certain lattice chains in K^{2n} . Let R be a k -algebra. Recall that an $R[[t]]$ -lattice in $R((t))^{2n}$ is an $R[[t]]$ -submodule $L \subset R((t))^{2n}$ which is free as an $R[[t]]$ -module Zariski-locally on $\text{Spec } R$, and such that the natural arrow $L \otimes_{R[[t]]} R((t)) \rightarrow R((t))^{2n}$ is an isomorphism. Borrowing our earlier notation, given an $R[[t]]$ -lattice L , we write \widehat{L} for the dual lattice

$$\widehat{L} := \{x \in R((t))^{2n} \mid h_{R((t))}(L, x) \subset R[[t]]\},$$

where $h_{R((t))} := h \otimes_K R((t))$ is the induced form on $R((t))^{2n}$. We say that an indexed sequence

$$\cdots \subset L_{-1} \subset L_0 \subset L_1 \subset \cdots$$

of lattices in $R((t))^{2n}$ is an *indexed chain* if all successive quotients are locally free R -modules. We say that an indexed chain L_\bullet is *periodic* if $tL_i = L_{i-2n}$ for all i , and *complete* if all successive quotients are locally free R -modules of rank 1.

We define \mathcal{F}' to be the functor on the category of k -algebras that sends each algebra R to the set of all complete periodic indexed lattice chains L_\bullet in $R((t))^{2n}$ with the property that Zariski-locally on $\text{Spec } R$, there exists a scalar $\alpha \in R((t))^\times$ such that $\widehat{L}_i = \alpha L_{-i}$ for all i . The natural action of $G(R((t)))$ on $R((t))^{2n}$ yields an action of LG on \mathcal{F}' . Taking the standard chain $\lambda_\bullet \in \mathcal{F}'(k)$ as base point, we obtain a map $LG \rightarrow \mathcal{F}'$ which induces, quite as in [PR4, §3.2],¹ an LG -equivariant isomorphism

$$\mathcal{F} \xrightarrow{\sim} \mathcal{F}'.$$

We shall always identify \mathcal{F} and \mathcal{F}' in this way.

6.3. Schubert cells and varieties. In this subsection we discuss Schubert cells and varieties in the affine flag variety. For $w \in \widetilde{W}$, the associated *Schubert cell* C_w is the reduced k -subscheme

$$C_w := L^+B \cdot \dot{w} \subset \mathcal{F},$$

where \dot{w} is any representative of w in $G(K)$. The associated *Schubert variety* S_w is the reduced closure of C_w in \mathcal{F} . Since $L^+B \subset LG^\circ$, every Schubert cell and variety is contained entirely in \mathcal{F}° or entirely in $\tau \mathcal{F}^\circ$. By (5.2.1), \widetilde{W} is in bijective

¹Though note that the scalar α in the definition of \mathcal{F}' in [PR4] should only be required to exist Zariski-locally, so that \mathcal{F}' satisfies the sheaf property.

correspondence with the set of Schubert cells in \mathcal{F} . We have $\dim C_w = \dim S_w = \ell(w)$.

6.4. Closure relations between Schubert cells. We now discuss closure relations between Schubert cells in \mathcal{F} . In the case of a connected reductive group over K , closure relations between Schubert cells correspond exactly to the Bruhat order in the Iwahori-Weyl group. Our aim here is to show that this statement carries over to our disconnected group G .

Proposition 6.4.1. *Let $w, w' \in \widetilde{W}$. Then $w \leq w'$ in the Bruhat order $\iff S_w \subset S_{w'}$ in \mathcal{F} .*

Proof. We reduce to the analogous statement for G° , using the decompositions $\widetilde{W} = \widetilde{W}^\circ \amalg \tau \widetilde{W}^\circ$ and $\mathcal{F} = \mathcal{F}^\circ \amalg \tau \mathcal{F}^\circ$. Let $w, w' \in \widetilde{W}$. Then for w and w' to be related in the Bruhat order on the one hand, and for S_w and $S_{w'}$ to be contained both in \mathcal{F}° or both in $\tau \mathcal{F}^\circ$ on the other hand, we must at least have $w \equiv w' \pmod{\widetilde{W}^\circ}$. So we suppose this is the case.

If $w, w' \in \widetilde{W}^\circ$, then the conclusion follows at once from the lemma for G° . If $w, w' \in \tau \widetilde{W}^\circ$, then we observe that

- the left-multiplication-by- τ map $\widetilde{W}^\circ \xrightarrow{\sim} \tau \widetilde{W}^\circ$ respects the Bruhat order, since τ stabilizes A ; and
- the left-multiplication-by- τ map $\mathcal{F}^\circ \xrightarrow{\sim} \tau \mathcal{F}^\circ$ respects Schubert cells, since τ normalizes L^+B in LG .

So the conclusion in this case follows again from the statement for G° . \square

7. EMBEDDING THE SPECIAL FIBER IN THE AFFINE FLAG VARIETY

In this section we embed the special fiber of M^{naive} into the affine flag variety \mathcal{F} .

7.1. The map. We write $M_k^{\text{naive}} := M^{\text{naive}} \otimes_{\mathcal{O}} k$. The embedding $M_k^{\text{naive}} \hookrightarrow \mathcal{F}$ we wish to construct will make use of the lattice-theoretic description of \mathcal{F} from §6.2. We first note that the \mathcal{O}_K -lattice chain λ_\bullet admits a “trivialization” in obvious analogy with (2.1.4), where λ_i replaces Λ_i , \mathcal{O}_K replaces \mathcal{O} , and t replaces π . Then this trivialization together with (2.1.4) itself and the canonical identifications $\mathcal{O}/\pi\mathcal{O} \cong k \cong \mathcal{O}_K/t\mathcal{O}_K$ yields an identification of chains of k -vector spaces

$$(*) \quad \Lambda_\bullet \otimes_{\mathcal{O}} k \cong \lambda_\bullet \otimes_{\mathcal{O}_K} k.$$

To define $M_k^{\text{naive}} \hookrightarrow \mathcal{F}$, suppose we have an R -point $\{\mathcal{F}_i \hookrightarrow \Lambda_i \otimes_{\mathcal{O}} R\}$ of M_k^{naive} for some k -algebra R . Let $L_i \subset \lambda_i \otimes_{\mathcal{O}_K} R[[t]]$ be the submodule rendering the diagram

$$\begin{array}{ccc} L_i \subset & \longrightarrow & \lambda_i \otimes_{\mathcal{O}_K} R[[t]] \\ \downarrow & & \downarrow \\ \mathcal{F}_i \subset & \longrightarrow & \Lambda_i \otimes_{\mathcal{O}} R \cong (\lambda_i \otimes_{\mathcal{O}_K} k) \otimes_k R \end{array}$$

Cartesian, where the identification in the bottom right corner is made via (*). Then the L_i 's form an indexed $R[[t]]$ -lattice chain

$$L_0 \subset L_1 \subset \cdots \subset L_{2n-1} \subset t^{-1}L_0$$

in $R((t))^{2n}$. The chain extends periodically to an R -point of \mathcal{F} (we may globally take the scalar α discussed in §6.2 to equal t^{-1}), which we take to be the image of our original R -point of M_k^{naive} . It is clear that $M_k^{\text{naive}} \hookrightarrow \mathcal{F}$ is then a monomorphism, and, as M_k^{naive} is proper, the map is a closed immersion. From now on, we frequently identify M_k^{naive} with its image in \mathcal{F} .

7.2. The image of the special fiber. Let R be a k -algebra. It is clear from the definition of the map $M_k^{\text{naive}} \hookrightarrow \mathcal{F}$ that the image of $M_k^{\text{naive}}(R)$ in $\mathcal{F}(R)$ consists precisely of all complete periodic self-dual chains L_\bullet in $\mathcal{F}(R)$ such that, for all i ,

- $\lambda_{i,R[[t]]} \supset L_i \supset t\lambda_{i,R[[t]]}$, where $\lambda_{i,R[[t]]} := \lambda_i \otimes_{\mathcal{O}_K} R[[t]]$; and
- the R -module $\lambda_{i,R[[t]]}/L_i$ is locally free of rank n for all i .

It is clear from this that the action of L^+B on \mathcal{F} preserves the closed subschemes M_k^{naive} and M_k^{spin} . We deduce that *the underlying topological spaces of M_k^{naive} and M_k^{spin} are unions of Schubert varieties in \mathcal{F}* . One of our essential goals for the rest of the paper is to obtain a good description of the Schubert varieties that occur in M_k^{spin} .

7.3. Schubert varieties in M_k^{naive} . As a preliminary step towards describing the Schubert varieties S_w that occur in M_k^{spin} , in this subsection we translate the condition that S_w be contained in the image of M_k^{naive} in \mathcal{F} into a condition on the extended alcove v_0, \dots, v_{2n-1} attached to $w \in \widetilde{W}$ (§5.6).

Upon inspecting definitions, the previous subsection makes plain that S_w is contained in $M_k^{\text{naive}} \iff$

- (P1) $\omega_i \leq v_i \leq \omega_i + (1, \dots, 1)$ for all $0 \leq i \leq 2n - 1$; and
- (P2) $\Sigma v_0 = n$.

We say that such a w is *GL-permissible*. If w is *GL-permissible*, then necessarily $d = 0$ in the duality condition (A3), and it follows from the duality condition that the inequalities in (P1) hold for all i as soon as they hold for all $0 \leq i \leq n$. The condition that w be *GL-permissible* is exactly the condition that it be permissible in $\widetilde{W}_{GL_{2n}}$ relative to the cocharacter $(1^{(n)}, 0^{(n)})$, or that, modulo conventions, its associated extended alcove be minuscule of size n in the terminology of [KR].

Given a *GL-permissible* w , the point $w \cdot \lambda_\bullet$ in $\mathcal{F}(k)$ corresponds to a point $\{\mathcal{F}_i \subset \Lambda_i \otimes_{\mathcal{O}} k\}$ in $M_k^{\text{naive}}(k)$ of a rather special sort: namely, identifying $\Lambda_i \otimes_{\mathcal{O}} k$ with k^{2n} via (2.1.4), we have

- (T) \mathcal{F}_i is spanned by standard basis vectors in k^{2n} for all i .

On the other hand, for any point $\{\mathcal{F}_i\}$ in $M_k^{\text{naive}}(k)$, let us say that $\{\mathcal{F}_i\}$ is a *T-fixed point* if it satisfies (T); it is easy to check that the *T-fixed points* are exactly the points in $M_k^{\text{naive}}(k)$ fixed by $L^+T(k)$. In this way, we get a bijection between the *GL-permissible* $w \in \widetilde{W}$ and the *T-fixed points* in $M_k^{\text{naive}}(k)$.

The *T-fixed point* $\{\mathcal{F}_i^w\}$ associated with a *GL-permissible* w is easy to describe in terms of the extended alcove v_0, \dots, v_{2n-1} . Indeed, let

$$(7.3.1) \quad \mu_i^w := v_i - \omega_i, \quad 0 \leq i \leq 2n - 1.$$

Then μ_i^w is a vector in \mathbb{Z}^{2n} having n entries equal to 0 and n entries equal to 1, and

$$(7.3.2) \quad \mathcal{F}_i^w = \sum_{\mu_i^w(j)=0} k\epsilon_j \subset k^{2n},$$

where $\epsilon_1, \dots, \epsilon_{2n}$ is the standard ordered basis in k^{2n} .

7.4. *T*-fixed points in M_k^{spin} . By the previous subsection, every Schubert cell in \mathcal{F} contained in M_k^{naive} contains a unique *T*-fixed point in M_k^{naive} . So to understand which Schubert cells are contained in M_k^{spin} , we need to understand which *T*-fixed points satisfy the spin condition. This is the object of this subsection.

We begin by fixing some notation. We continue to write e_1, \dots, e_{2n} for the standard basis in V and $\epsilon_1, \dots, \epsilon_{2n}$ for the standard basis in k^{2n} , and we identify Λ_i with \mathcal{O}^{2n} , and hence $\Lambda_i \otimes k$ with k^{2n} , via (2.1.4). Quite generally, for any subset $E \subset \{1, \dots, 2n\}$, we define

$$kE := \sum_{j \in E} k\epsilon_j \subset k^{2n}.$$

When E has cardinality n , consider the wedge product, in increasing index order, of the n standard basis vectors in \mathcal{O}^{2n} indexed by the elements of E ; we denote by $e_E^i \in \bigwedge_F^n V$ the image of this element under the map $\bigwedge_{\mathcal{O}}^n \mathcal{O}^{2n} \xrightarrow{\sim} \bigwedge_{\mathcal{O}}^n \Lambda_i \subset \bigwedge_F^n V$. When $i = 0$, we have $e_E^0 = e_E$ (2.3.1).

Now let $\{\mathcal{F}_i \subset k^{2n}\}$ be a *T*-fixed point in $M_k^{\text{naive}}(k)$. For each i , let $E_i \subset \{1, \dots, 2n\}$ be the subset of indices j such that $\epsilon_j \in \mathcal{F}_i$, so that $\mathcal{F}_i = kE_i$ and $\mathcal{F}_i^\perp = kE_i^\perp$ (2.3.2). To understand the spin condition for the \mathcal{F}_i 's, we need to get a good handle on the elements $e_{E_i}^i$ and $e_{E_i^\perp}^i$. More precisely, let

$$d_i := \#(E_i \cap \{1, \dots, i\}) \quad \text{and} \quad d_i^\perp := \#(E_i^\perp \cap \{1, \dots, i\}).$$

Then, referring again to (2.3.1),

$$e_{E_i}^i = \frac{1}{\pi^{d_i}} e_{E_i} \quad \text{and} \quad e_{E_i^\perp}^i = \frac{1}{\pi^{d_i^\perp}} e_{E_i^\perp},$$

and we need to understand the integer $d_i^\perp - d_i$.

To proceed, we'll consider pairs of the form $(i, 2n - i)$ simultaneously, so that we may assume $0 \leq i \leq n$. Let

$$A_i := \{1, \dots, i, i^*, \dots, 2n\} \quad \text{and} \quad B_i := \{i + 1, \dots, 2n - i\},$$

so that we get an orthogonal decomposition $k^{2n} = kA_i \oplus kB_i$. Since \mathcal{F}_n is totally isotropic, E_i cannot contain any pair of the form j, j^* with $j \leq i$. Hence we may write A_i as a disjoint union

$$A_i = R_i \amalg S_i,$$

where

$$R_i := \{j \in A_i \mid \text{exactly one of } j, j^* \text{ is in } E_i\} \quad \text{and} \quad S_i := \{j \in A_i \mid j, j^* \notin E_i\}.$$

Plainly, the sets R_i and S_i have even cardinalities, say equal to $2r_i$ and $2s_i$, respectively. We have

$$E_i \cap A_i = E_i \cap R_i \quad \text{and} \quad E_i^\perp \cap A_i = (E_i^\perp \cap R_i) \amalg S_i = (E_i \cap R_i) \amalg S_i.$$

Hence

$$\#(E_i \cap A_i) = r_i \quad \text{and} \quad \#(E_i^\perp \cap A_i) = r_i + 2s_i.$$

We now need a couple of lemmas.

Lemma 7.4.1. *For $1 \leq i \leq n$, the image of \mathcal{F}_i in \mathcal{F}_{2n-i} under the structure maps is totally isotropic.*

Proof. Since \mathcal{F}_i is spanned by standard basis vectors, the image in question is contained in $\mathcal{F}_i \cap \mathcal{F}_{2n-i} = \mathcal{F}_i \cap \mathcal{F}_i^\perp$. \square

The $i = 0$ version of (7.4.1) is simply the statement that $\mathcal{F}_0 = \mathcal{F}_0^\perp$.

Lemma 7.4.2. $\#(E_i \cap A_i) \leq i$.

Proof. The intersection $\mathcal{F}_i \cap kA_i$ is the precisely the image of \mathcal{F}_i in \mathcal{F}_{2n-i} under the structure maps, hence is totally isotropic by the previous lemma. Since the form on k^{2n} restricts to a nondegenerate form on kA_i , we conclude

$$\#(E_i \cap A_i) = \dim \mathcal{F}_i \cap kA_i \leq \frac{1}{2} \dim kA_i = i. \quad \square$$

The lemma leaves us with two cases to consider.

Case 1: $\#(E_i \cap A_i) < i$. Then $S_i \neq \emptyset$. Hence

$$d_i^\perp - d_i = s_i > 0.$$

Hence by (2.3.3), we have

$$e_{E_i}^i \pm \pi^{d_i^\perp - d_i} \operatorname{sgn}(\sigma_{E_i}) e_{E_i^\perp}^i \in \left(\bigwedge_{\emptyset}^n \Lambda_i \right)_\pm,$$

and the image of this element under the map $(\bigwedge_{\emptyset}^n \Lambda_i)_\pm \otimes k \rightarrow \bigwedge_k^n \Lambda_i \otimes k$ spans the line $\bigwedge_k^n \mathcal{F}_i$. Moreover, it is easy to check that $d_{2n-i}^\perp - d_{2n-i} = s_i$ as well, so that we similarly conclude $\bigwedge_k^n \mathcal{F}_{2n-i} \subset \operatorname{im}[(\bigwedge_{\emptyset}^n \Lambda_{2n-i})_\pm \otimes k \rightarrow \bigwedge_k^n \Lambda_{2n-i} \otimes k]$.

Case 2: $\#(E_i \cap A_i) = i$. We claim $E_i = E_i^\perp$, that is, \mathcal{F}_i is a (maximal) totally isotropic subspace of k^{2n} . Indeed, in this case $\mathcal{F}_i \cap kA_i$ is maximal totally isotropic in kA_i , and it suffices to show that $\mathcal{F}_i \cap kB_i$ is totally isotropic of dimension $n - i$. For this, consider the structure map $f: \mathcal{F}_{2n-i} \rightarrow \mathcal{F}_i$. Then $\operatorname{im} f$ is plainly contained in kB_i and is totally isotropic by the argument in (7.4.1). So it suffices, in turn, to show that $\ker f = \mathcal{F}_{2n-i} \cap kA_i$ has dimension i . But

$$\mathcal{F}_{2n-i} \cap kA_i \subset (\mathcal{F}_i \cap kA_i)^\perp \cap kA_i = \mathcal{F}_i \cap kA_i,$$

where the equality in the display follows from our case assumption, and the reverse inclusion $\mathcal{F}_i \cap kA_i \subset \mathcal{F}_{2n-i} \cap kA_i$ is trivial. The claim follows. We deduce that $e_{E_i}^i$ and $e_{E_{2n-i}}^{2n-i} = e_{E_i}^{2n-i}$ are scalar multiples of each other; and as in [PR4, §7.1.4], both are contained in the one of the submodules $(\bigwedge_{\emptyset}^n \Lambda_i)_\pm$.

We obtain the following.

Proposition 7.4.3. *Let $\{\mathcal{F}_i \subset \Lambda_i \otimes k\}$ be a T -fixed point in $M^{\text{naive}}(k)$. The following are equivalent.*

- (i) $\{\mathcal{F}_i \subset \Lambda_i \otimes k\}$ satisfies the spin condition.
- (ii) Upon identifying the $\Lambda_i \otimes k$'s with k^{2n} via (2.1.4), all the \mathcal{F}_i 's for $0 \leq i \leq n$ which are totally isotropic in k^{2n} specify points on the same connected component of the orthogonal Grassmannian $\operatorname{OGr}(n, 2n)$.
- (iii) Under the above identifications, whenever \mathcal{F}_i and $\mathcal{F}_{i'}$ for $0 \leq i, i' \leq n$ are totally isotropic in k^{2n} , $\mathcal{F}_i \cap \mathcal{F}_{i'}$ has even codimension in \mathcal{F}_i and $\mathcal{F}_{i'}$.
- (iv) The sets E_i for $0 \leq i \leq n$ for which $E_i = E_i^\perp$ are all W° -conjugate under the natural action of W° on $\{1, \dots, 2n\}$.

Proof. (i) \iff (ii) has already been explained. (ii) \iff (iii) is explained in [PR4, §7.1.4]. (ii) \iff (iv) is clear from the facts that the orthogonal group acts transitively on OGr , and that the element τ (§5.1) interchanges the two components. \square

7.5. Schubert varieties in M_k^{spin} . We now use the previous subsection to express the condition that the Schubert variety S_w attached to $w \in \widetilde{W}$ is contained in M_k^{spin} . Continuing from §7.3, we shall express this condition in terms of the extended alcove v_0, \dots, v_{2n-1} attached to w .

Let w be GL -permissible. Then the condition we wish to formulate can be essentially read off from (7.4.3). Recall the vector μ_i^w (7.3.1) and the subspace $\mathcal{F}_i^w \subset k^{2n}$ (7.3.2). We say μ_i^w is *totally isotropic* if $\mu_i(j) = 1 - \mu_i(j^*)$ for all j , or equivalently if \mathcal{F}_i^w is totally isotropic in k^{2n} . It is now immediate from our considerations of T -fixed points and from (7.4.3) that S_w is contained in M_k^{spin} $\iff w$ is GL -permissible and, in addition, satisfies

(P3) (spin condition) the vectors μ_i^w for $0 \leq i \leq n$ which are totally isotropic are all W° -conjugate.

The following trivial reformulation of (P3) is sometimes convenient. Borrowing our notation from the previous subsection, let $E_i^w \subset \{1, \dots, 2n\}$ be the subset

$$(7.5.1) \quad E_i^w := \{j \mid \mu_i(j) = 0\}.$$

We say E_i^w is *totally isotropic* if $E_i^w = (E_i^w)^\perp$, or equivalently if μ_i^w is totally isotropic. Then for GL -permissible w , condition (P3) is equivalent to

(P3') (spin condition') The sets E_i^w for $0 \leq i \leq n$ which are totally isotropic are all W° -conjugate.

We say that $w \in \widetilde{W}$ is *spin-permissible* if it satisfies (P1), (P2), and (P3), or equivalently (P1), (P2), and (P3'). It follows from the duality condition (A3) that for GL -permissible w , the vectors μ_0^w and μ_n^w are always totally isotropic; but in general, even for spin-permissible w , the possibilities can range from these two being the only totally isotropic vectors to all the all μ_i^w 's being totally isotropic.

It is useful to formulate a slight refinement of the notion of spin-permissible. There are exactly two orbits for the action of W° on the set of totally isotropic vectors with n entries equal to 0 and n entries equal to 1, namely

$$W^\circ \mu_1 \quad \text{and} \quad W^\circ \mu_2,$$

where $\mu_1 = (1^{(n)}, 0^{(n)})$ and $\mu_2 = (1^{(n-1)}, 0, 1, 0^{(n-1)})$ are the cocharacters of (1.1). For $j = 1, 2$, we say that w is μ_j -*spin-permissible* if w is GL -permissible and $\mu_i^w \in W^\circ \mu_j$ whenever μ_i^w is totally isotropic. We write $\text{Perm}^{\text{SP}}(\mu_j)$ for the set of μ_j -spin-permissible elements in \widetilde{W} . Thus the set of spin-permissible elements in \widetilde{W} is the disjoint union $\text{Perm}^{\text{SP}}(\mu_1) \amalg \text{Perm}^{\text{SP}}(\mu_2)$. We shall see in §8.8 that for $j = 1, 2$, $\text{Perm}^{\text{SP}}(\mu_j)$ is precisely the μ_j -*permissible set* defined by Kottwitz and Rapoport [KR].

7.6. Topological flatness of M^{spin} . We now come to the main result of the paper. We again recall the dominant minuscule cocharacters μ_1 and μ_2 for G from (1.1), and for any cocharacter $\mu \in X_*(T)$, we recall the μ -admissible sets $\text{Adm}(\mu)$ and $\text{Adm}^\circ(\mu)$ from §5.5. Let $\mathcal{A}(\mu)$ denote the reduced union of Schubert varieties $\bigcup_{w \in \text{Adm}^\circ \mu} S_w$ in the affine flag variety.

Theorem 7.6.1.

- (i) $\text{Adm}^\circ(\mu_1) = \text{Perm}^{\text{SP}}(\mu_1)$ and $\text{Adm}^\circ(\mu_2) = \text{Perm}^{\text{SP}}(\mu_2)$. In particular, the set $\text{Adm}(\mu_1) = \text{Adm}(\mu_2)$ equals $\text{Perm}^{\text{SP}}(\mu_1) \amalg \text{Perm}^{\text{SP}}(\mu_2)$.

- (ii) *The underlying topological space of the special fiber M_k^{spin} coincides with $\mathcal{A}(\mu_1) \amalg \mathcal{A}(\mu_2)$ in \mathcal{F} . In particular, M_k^{spin} has two isomorphic connected components, and the irreducible components of these are in respective bijective correspondence with $W^\circ \mu_1$ and $W^\circ \mu_2$.*
- (iii) *The underlying topological space of M^{spin} is the closure of the generic fiber in M^{naive} . In particular, M^{spin} is topologically flat.*

Proof. Assertion (ii) follows immediately from (i) and §7.5. To prove (ii), we must show that the irreducible components of M_k^{spin} are in the closure of the generic fiber $M_F^{\text{spin}} = M_F^{\text{naive}}$ in M^{naive} . This follows from (ii) by a more-or-less standard lifting argument. By a lemma of Görtz [G4, Lemma 2], it suffices to show that each irreducible component in M_k^{spin}

- (1) has dimension equal to the dimension of M_F^{spin} ; and
- (2) contains a closed point which is contained in no other irreducible component and which lifts to the generic fiber.

For (1), for $\mu \in W^\circ \mu_1 \cup W^\circ \mu_2$, one readily computes from the formula of Iwahori-Matsumoto [IM, Proposition 1.23]

$$\dim S_{t_\mu} = \ell(t_\mu) = \sum_{\substack{\text{positive} \\ \text{roots } \alpha}} |\langle \mu, \alpha \rangle| = \binom{n}{2}.$$

On the other hand, it is well-known that $M_F^{\text{spin}} \cong \text{OGr}(n, 2n)_F$ has dimension $\binom{n}{2}$. For (2), for each $\mu \in W^\circ \mu_1 \cup W^\circ \mu_2$, we just take the T -fixed point $\{\mathcal{F}_i^{t_\mu} \subset k^{2n}\} \in M^{\text{spin}}(k)$ attached to μ itself. Then the $\mathcal{F}_i^{t_\mu}$'s are all equal and spanned by the standard basis vectors ϵ_j for which $\mu(j) = 0$, and we have the obvious lift $\{\widetilde{\mathcal{F}}_i^{t_\mu} \subset \mathcal{O}^{2n}\} \in M^{\text{spin}}(\mathcal{O})$ where $\widetilde{\mathcal{F}}_i^{t_\mu}$ is the span of the corresponding standard basis vectors in \mathcal{O}^{2n} for all i .

It remains to prove (i). To prove the containments $\text{Adm}^\circ(\mu_j) \subset \text{Perm}^{\text{sp}}(\mu_j)$ for $j = 1, 2$, we first note that since M_k^{spin} is closed in \mathcal{F} and the Bruhat order reflects closure relations between Schubert varieties (6.4.1), $\text{Perm}^{\text{sp}}(\mu_j)$ is closed in the Bruhat order. Hence it suffices to show that $\text{Perm}^{\text{sp}}(\mu_j)$ contains the maximal elements of $\text{Adm}^\circ(\mu_j)$, that is, the W° -conjugates of t_{μ_j} in \widetilde{W} , which is obvious.

We are left to prove the containments $\text{Adm}^\circ(\mu_j) \supset \text{Perm}^{\text{sp}}(\mu_j)$ for $j = 1, 2$ in (i). This is the main object of §8. \square

8. ADMISSIBLE, PERMISSIBLE, AND SPIN-PERMISSIBLE SETS

Let $\mu \in \{\mu_1, \mu_2\}$. In this section we complete the proof of part (i) of (7.6.1) by showing that $\text{Perm}^{\text{sp}}(\mu) \subset \text{Adm}^\circ(\mu)$. In essence, this amounts to working through the argument of Kottwitz-Rapoport [KR, §5] in the case of the orthogonal similitude group. In the last subsection §8.8, we show that the notion of μ -spin-permissibility (§7.5) agrees with the notion of μ -permissibility from [KR].

8.1. Strategy. Our strategy for proving $\text{Perm}^{\text{sp}}(\mu) \subset \text{Adm}^\circ(\mu)$ is, in the large, the same strategy Kottwitz and Rapoport used to prove the analogous assertion for GL_r . Namely, let $w \in \text{Perm}^{\text{sp}}(\mu)$. Then the asserted containment holds $\iff w$ is a translation element, or w is not a translation element and we can find a reflection $s \in W_a$ such that $sw \in \text{Perm}^{\text{sp}}(\mu)$ and $sw > w$ in the Bruhat order. In the GL case, when w is not a translation element, Kottwitz and Rapoport found an explicit

affine root α such that the associated reflection had the desired properties. Since every affine root for G is the restriction of an affine root for GL_{2n} , we can approach the problem in our case in the following way: regarding w as an element in $\widetilde{W}_{GL_{2n}}$, we can take the affine root α prescribed by Kottwitz and Rapoport, attempt to restrict α to the maximal torus T in G , and then take the corresponding reflection in W_a . Two problems arise.

- (1) α may not restrict to an affine root of G .
- (2) Even when α does restrict to an affine root of G with associated reflection s_α , although one can show that $s_\alpha w > w$ and that $s_\alpha w$ satisfies (P1) and (P2), $s_\alpha w$ need not satisfy (P3).

It turns out that the first problem is quite easy to overcome. But the second is more serious and leads us to a more complicated case analysis than that encountered in [KR].

8.2. Reflections. Consider the affine linear function

$$\begin{array}{ccc} \alpha_{i,j;d}: & X_*(T) & \longrightarrow \mathbb{Z} \\ & (x_1, \dots, x_{2n}) & \longmapsto x_i - x_j - d \end{array}$$

for $i < j$ and $d \in \mathbb{Z}$. Then $\alpha := \alpha_{i,j;d}$ is an affine root of (G, T) precisely when $j \neq i^*$, and up to sign, all affine roots are obtained in this way. Plainly $\alpha_{i,j;d} = \alpha_{j^*,i^*;d}$. Attached to α is the reflection $s_\alpha = s_{i,j;d} \in W_a$ which acts on $X_*(T) \otimes \mathbb{R}$ by sending (x_1, \dots, x_n) to the tuple with $x_j + d$ in the i th slot, $x_i - d$ in the j th slot, $x_{i^*} + d$ in the j^* th slot, $x_{j^*} - d$ in the i^* th slot, and all other slots the same; visually, in the case $i < j < j^* < i^*$,

$$\begin{array}{ccc} (\dots, x_i, \dots, x_j, \dots, x_{j^*}, \dots, x_{i^*}, \dots) & \xrightarrow{s_\alpha} & (\dots, x_j + d, \dots, x_i - d, \dots, x_{i^*} + d, \dots, x_{j^*} - d, \dots). \end{array}$$

If $w \in \widetilde{W}$ has extended alcove v_0, \dots, v_{2n-1} , then $s_\alpha w$ has extended alcove $s_\alpha v_0, \dots, s_\alpha v_{2n-1}$.

8.3. The set K_m . Fix a GL -permissible $w \in \widetilde{W}$, and recall the vector μ_k^w for $0 \leq k \leq 2n-1$ from (7.3.1). As in [KR], for $1 \leq m \leq 2n$, we define $K_m \subset \{0, \dots, 2n-1\}$ to be the subset

$$K_m := \{k \mid \mu_k^w(m) = 0\}.$$

Just as in [KR, 5.4–5.5], and in the notation and terminology used there, the set K_m is either empty, all of $\{0, \dots, 2n-1\} \cong \mathbb{Z}/2n\mathbb{Z}$, or an *interval* in $\mathbb{Z}/2n\mathbb{Z}$ of the form $[\tilde{m}, m)$ for some $\tilde{m} \neq m$; in this last case, we say that m is *proper* and that K_m has *lower endpoint* \tilde{m} and *upper endpoint* m . For proper m , we always denote by \tilde{m} the lower endpoint of K_m . Of course, the lower endpoint $\tilde{m} \in \mathbb{Z}/2n\mathbb{Z}$ is characterized by the property

$$\tilde{m} \in K_m \quad \text{and} \quad \tilde{m} - 1 \notin K_m.$$

When m is proper, \tilde{m} is evidently proper too, and we have the simple formula

$$(8.3.1) \quad E_{\tilde{m}}^w = (m, \tilde{m}) \cdot E_{\tilde{m}-1}^w,$$

where (m, \tilde{m}) is the transposition interchanging m and \tilde{m} and the set $E_{\tilde{m}}^w$ is defined in (7.5.1). Plainly, the function $m \mapsto \tilde{m}$ defines a fixed-point-free bijection from the

set of proper elements in $\mathbb{Z}/2n\mathbb{Z}$ to itself. Note that, asymmetrically, we embed w into our notation for E_i^w but suppress w in our notation for K_m .

The duality condition (A3) may be expressed in terms of the E^w 's as

$$m \in E_k^w \iff m^* \notin E_{2n-k}^w$$

and in terms of the K 's as

$$k \in K_m \iff 2n - k \notin K_{m^*}.$$

Hence $K_{m^*} = -K_m^c$ for all m as subsets of $\mathbb{Z}/2n\mathbb{Z}$. Hence if K_m is an interval $[\tilde{m}, m)$, then K_{m^*} is the interval $[(\tilde{m})^*, m^*)$; in particular, $(m^*)^\sim = (\tilde{m})^*$. Moreover, m fails to be proper exactly when $K_m = \mathbb{Z}/2n\mathbb{Z}$ and $K_{m^*} = \emptyset$, or $K_m = \emptyset$ and $K_{m^*} = \mathbb{Z}/2n\mathbb{Z}$.

8.4. Reflections and GL -permissibility. Suppose $w \in \widetilde{W}$ is GL -permissible. In this subsection we determine the affine roots α such that $s_\alpha w$ is again GL -permissible. As usual, we denote by v_0, \dots, v_{2n-1} the extended alcove attached to w .

Recall from §7.3 that w is GL -permissible $\iff \Sigma v_0 = n$ and $\omega_k \leq v_k \leq \omega_k + (1^{(2n)})$ for all $0 \leq k \leq 2n-1$. Hence for $\alpha = \alpha_{i,j;d}$ with $i < j \neq i^*$, the element $s_\alpha w$ is GL -permissible \iff

$$(*) \quad \left. \begin{array}{l} v_k(j) + d - \omega_k(i) \in \{0, 1\} \\ v_k(i) - d - \omega_k(j) \in \{0, 1\} \\ v_k(i^*) + d - \omega_k(j^*) \in \{0, 1\} \\ v_k(j^*) - d - \omega_k(i^*) \in \{0, 1\} \end{array} \right\} \text{ for all } 0 \leq k \leq 2n-1.$$

By the duality condition, the last two containments in $(*)$ hold for all $k \iff$ the first two hold for all k .

It is convenient to express the conditions in $(*)$ in terms of the sets $[i, j)$, K_i , and K_j . For any subset $S \subset \{0, \dots, 2n-1\}$, let χ_S denote the characteristic function of S . Then for all k and m ,

$$\chi_{[i,j)}(k) = \omega_k(j) - \omega_k(i) \quad \text{and} \quad \chi_{K_m}(k) = 1 - \mu_k^w(m) = 1 - v_k(m) + \omega_k(m).$$

Hence we may rewrite the first two conditions in $(*)$ as

$$\chi_{K_j}(k) - \chi_{[i,j)}(k) - d \in \{0, 1\} \quad \text{and} \quad \chi_{K_i}(k) + \chi_{[i,j)}(k) + d \in \{0, 1\}$$

for all $0 \leq k \leq 2n-1$. Similarly to [KR, 5.2], either of these last two conditions implies that d equals 0 or -1 . We similarly conclude from the two conditions together that for $d = 0$,

$$s_{i,j;0}w \text{ is } GL\text{-permissible} \iff [i, j) \subset K_i^c \cap K_j,$$

and that for $d = -1$,

$$s_{i,j;-1}w \text{ is } GL\text{-permissible} \iff [i, j)^c \subset K_i \cap K_j^c.$$

The following is a convenient reformulation of the above discussion.

Lemma 8.4.1. *Let $i, j \in \{1, \dots, 2n\}$ with $j \neq i, i^*$. Then*

$$\begin{array}{l} \text{either } i < j \text{ and } s_{i,j;0}w \text{ is } GL\text{-permissible,} \\ \text{or } j < i \text{ and } s_{j,i;-1}w \text{ is } GL\text{-permissible} \end{array} \iff i \in K_j \text{ and } j-1 \notin K_i.$$

Proof. This is clear from the above discussion and the fact that K_i , resp. K_j , is either empty, all of $\mathbb{Z}/2n\mathbb{Z}$, or an interval with upper endpoint i , resp. j . \square

8.5. Reflections and the Bruhat order. We continue with our $w \in \widetilde{W}$ and affine root $\alpha = \alpha_{i,j;d}$ with $i < j \neq i^*$. The elements w and $s_\alpha w$ are related in the Bruhat order, and we have $w < s_\alpha w$ exactly when our base alcove A and the alcove wA lie on the same side of the hyperplane in \mathfrak{a} where α vanishes. We wish to understand this condition in terms of α and the extended alcove attached to w .

Actually, instead of working directly with A , it will be more convenient to use the analogous alcove A' for the symplectic group: this is the interior of the convex hull in \mathfrak{a} of the $n + 1$ points

$$a'_k := \frac{\omega_k + \omega_{2n-k}}{2} \pmod{\mathbb{R} \cdot (1, \dots, 1)} \quad \text{for } 0 \leq k \leq n.$$

Then $A' \subset A$, so that it suffices to use A' and wA' to detect the Bruhat relation between w and $s_\alpha w$. The vertices of wA' are

$$wa'_k = \frac{v_k + v_{2n-k}}{2} \pmod{\mathbb{R} \cdot (1, \dots, 1)} \quad \text{for } 0 \leq k \leq n.$$

Hence

$$(*) \quad \alpha(wa'_k) = \frac{\chi_{K_j}(k) - \chi_{K_i}(k) - \chi_{[i,j]}(k)}{2} + \frac{\chi_{K_j}(2n-k) - \chi_{K_i}(2n-k) - \chi_{[i,j]}(2n-k)}{2} - d$$

for $0 \leq k \leq n$.

When $d \geq 0$, the values of α on the vertices of A' are nonpositive. Hence, in this case,

$$w < s_\alpha w \iff \text{the value in } (*) \text{ is negative for some } k.$$

On the other hand, when $d \leq -1$, the values of α on the vertices of A' are nonnegative. Hence, in this case,

$$w < s_\alpha w \iff \text{the value in } (*) \text{ is positive for some } k.$$

The following lemma builds on (8.4.1) to give a useful characterization of when $s_\alpha w$ is GL -permissible and $w < s_\alpha w$.

Lemma 8.5.1. *Let $i, j \in \{1, \dots, 2n\}$ with $j \neq i, i^*$. Suppose that i is proper, so that K_i is an interval $[\tilde{i}, i)$ with $\tilde{i} \neq i$. Then*

$$\begin{aligned} & \text{either } i < j, s_{i,j;0}w \text{ is } GL\text{-permissible, and } w < s_{i,j;0}w; \\ & \text{or } j < i, s_{j,i;-1}w \text{ is } GL\text{-permissible, and } w < s_{j,i;-1}w \end{aligned} \iff i \in K_j \text{ and } \tilde{i} \notin K_j.$$

Proof. We'll only need to use the implication " \Leftarrow " later on, so we'll just prove that and leave the implication " \Rightarrow " to the reader. Let α denote the affine root $\alpha_{i,j;0}$ or $\alpha_{j,i;-1}$ according as $i < j$ or $j < i$.

We first address GL -permissibility. By (8.4.1), regardless of the ordering of i and j , we must show $j - 1 \notin K_i$. But our hypotheses $i \in K_j$ and $\tilde{i} \notin K_j$ clearly imply $j - 1 \in [i, \tilde{i}) = K_i^c$, where the superscript c denotes the complement in $\mathbb{Z}/2n\mathbb{Z}$, as desired.

So it remains to show $w < s_\alpha w$. We first suppose $i < j$, which leads us to look at the expression $\chi_{K_j} - \chi_{K_i} - \chi_{[i,j]}$. Since j is plainly proper by hypothesis, K_j is an interval $[\tilde{j}, j)$ for some $\tilde{j} \neq j$. Since $i \in K_j$, we have $[i, j) \subset K_j$, and

$$\chi_{K_j} - \chi_{[i,j]} = \chi_{[\tilde{j}, i)},$$

where we interpret $[\tilde{j}, i] = \emptyset$ if $\tilde{j} = i$. Moreover, since $\tilde{i} \notin K_j$ and $i \in K_j$, we have $\tilde{j} \in [\tilde{i} + 1, i + 1)$. Hence $\tilde{j} - 1 \in K_i$. Hence

$$\chi_{K_j} - \chi_{[i,j]} - \chi_{K_i} = -\chi_{[\tilde{i}, \tilde{j}]}.$$

Note that here $\tilde{i} \neq \tilde{j}$ by injectivity of the map $m \mapsto \tilde{m}$. Hence

$$\alpha(wa'_k) = \frac{-\chi_{[\tilde{i}, \tilde{j}]}(k) - \chi_{[\tilde{i}, \tilde{j}]}(2n - k)}{2}$$

is certainly negative for some $0 \leq k \leq n$, as desired.

In the case $j < i$ with $\alpha = \alpha_{j,i,-1}$, one must find a vertex of wA' on which α is positive. This time one considers the expression

$$\chi_{K_i} - \chi_{K_j} - \chi_{[j,i]} + 1 = \chi_{K_i} - \chi_{K_j} + \chi_{[i,j]},$$

which by the above reasoning equals $\chi_{[\tilde{i}, \tilde{j}]}$, and the rest of the proof goes through similarly. \square

As an important application, we obtain the following lemma.

Lemma 8.5.2. *Let $r \in \{1, \dots, 2n\}$, and suppose that K_r is an interval $[\tilde{r}, r)$ for some $\tilde{r} \neq r, r^*$.*

- (i) *If $[r, \tilde{r}) \subset K_{\tilde{r}}$ and $r < \tilde{r}$, let $\alpha := \alpha_{r, \tilde{r}; 0}$.*
- (ii) *If $[r, \tilde{r}) \subset K_{\tilde{r}}$ and $\tilde{r} < r$, let $\alpha := \alpha_{\tilde{r}, r; -1}$.*
- (iii) *If $K_{\tilde{r}} \subset [r, \tilde{r})$ and $r < \tilde{r}$, let $\alpha := \alpha_{r, \tilde{r}; -1}$.*
- (iv) *If $K_{\tilde{r}} \subset [r, \tilde{r})$ and $\tilde{r} < r$, let $\alpha := \alpha_{\tilde{r}, r; 0}$.*

Then in each case, $s_\alpha w$ is GL -permissible and $w < s_\alpha w$.

Note that, since $[r, \tilde{r})$ and $K_{\tilde{r}}$ are both intervals with upper endpoint \tilde{r} , the hypotheses in at least one of (i)–(iv) will always be satisfied. So the force of the lemma is that, provided r is proper and $\tilde{r} \neq r^*$, we always get an affine reflection that preserves GL -permissibility and increases length.

Proof of (8.5.2). We use (8.5.1). To handle (i) and (ii), we must show $r \in K_{\tilde{r}}$ and $\tilde{r} \notin K_{\tilde{r}}$, both of which are obvious. To handle (iii) and (iv), we must show $\tilde{r} \in K_r$ and $\tilde{r} \notin K_r$. The first of these is obvious, and the second follows from

$$\tilde{r} \in K_{\tilde{r}} \subset [r, \tilde{r}) = K_r^c. \quad \square$$

8.6. Reflections and the spin condition. We continue with our w and $\alpha = \alpha_{i,j,d}$ with $i < j \neq i^*$. We now suppose that w and $s_\alpha w$ are GL -permissible, and we wish to relate the spin condition on $s_\alpha w$ to the spin condition on w . By §8.4, we must have $d = 0$ or $d = -1$. Let $l_1 < l_2 < l_3 < l_4$ denote the elements of the set $\{i, i^*, j, j^*\}$ in increasing order, and consider the sets E_k^w and $E_k^{s_\alpha w}$ (7.5.1) for $0 \leq k \leq n$. It is easy to verify that for $1 \leq k < l_1$ and for $l_2 \leq k \leq n$, the sets E_k^w and $E_k^{s_\alpha w}$ are equal or conjugate by the permutation $(i, j)(i^*, j^*)$. Hence, if w is μ -spin-permissible, then we at least know that the totally isotropic $E_k^{s_\alpha w}$ for $k \in \{0, \dots, n\} \setminus [l_1, l_2)$ are W° -conjugate to E_0^w , and hence to $E_0^{t\mu}$.

It is a more subtle matter to handle the $E_k^{s_\alpha w}$'s for $k \in [l_1, l_2)$. Since $i < j \neq i^*$, there are four possibilities to consider:

$$i < j < j^* < i^*, \quad i < j^* < j < i^*, \quad j^* < i < i^* < j, \quad \text{or} \quad j^* < i^* < i < j.$$

In each case, one element $\text{ext}(i, j)$ of the pair i, j is extremal amongst the four elements, and the other element $\text{int}(i, j)$ of the pair is not; and ditto for the pair i^*, j^* . For fixed $k \in [l_1, l_2]$, one verifies that either

$$\text{ext}(i, j) \notin E_k^w, E_k^{s_\alpha w}, \quad \text{int}(i, j) \in E_k^w, E_k^{s_\alpha w}, \quad \text{and} \quad E_k^{s_\alpha w} = (i^*, j^*) \cdot E_k^w;$$

or

$$\text{ext}(i, j)^* \notin E_k^w, E_k^{s_\alpha w}, \quad \text{int}(i, j)^* \in E_k^w, E_k^{s_\alpha w}, \quad \text{and} \quad E_k^{s_\alpha w} = (i, j) \cdot E_k^w.$$

Example 8.6.1. The following illustration of our discussion will come up explicitly in §8.7. Assume that $i < j < j^* < i^*$ and that

$$i, j^* \in E_{i-1}^w, \quad i^*, j \notin E_{i-1}^w, \quad i, i^* \notin E_i^w, \quad \text{and} \quad j, j^* \in E_i^w.$$

Then, displaying the i th, j th, j^* th, and i^* th entries,

$$v_{i-1} = (\dots, 0, \dots, 1, \dots, 0, \dots, 1, \dots) \quad \text{and} \quad v_i = (\dots, 0, \dots, 0, \dots, 0, \dots, 1, \dots).$$

Hence

$$\begin{aligned} s_{i,j;0}v_{i-1} &= (\dots, 1, \dots, 0, \dots, 1, \dots, 0, \dots), \\ s_{i,j;0}v_i &= (\dots, 0, \dots, 0, \dots, 1, \dots, 0, \dots), \\ s_{i,j;-1}v_{i-1} &= (\dots, 0, \dots, 1, \dots, 0, \dots, 1, \dots), \quad \text{and} \\ s_{i,j;-1}v_i &= (\dots, -1, \dots, 1, \dots, 0, \dots, 1, \dots). \end{aligned}$$

Hence for $\alpha = \alpha_{i,j;0}$, provided $s_\alpha w$ is GL -permissible, we conclude

$$E_{i-1}^{s_\alpha w} = (i, j)(i^*, j^*)E_{i-1}^w \quad \text{and} \quad E_i^{s_\alpha w} = (i^*, j^*)E_i^w;$$

and for $\alpha = \alpha_{i,j;-1}$, provided $s_\alpha w$ is GL -permissible, we conclude

$$E_{i-1}^{s_\alpha w} = E_{i-1}^w \quad \text{and} \quad E_i^{s_\alpha w} = (i, j)E_i^w.$$

Either way, we conclude $E_{i-1}^{s_\alpha w} = E_{i-1}^{s_\alpha w}$. The same conclusions plainly hold if $i < j^* < j < i^*$.

Part (i) of the following lemma summarizes the first paragraph of this subsection, and part (ii) is an immediate consequence of the second paragraph.

Lemma 8.6.2. *Suppose that w is μ -spin-permissible and that $s_\alpha w$ is GL -permissible.*

- (i) $s_\alpha w$ fails to be μ -spin-permissible \iff there exists $k \in [l_1, l_2]$ such that $E_k^{s_\alpha w}$ is totally isotropic and not W° -conjugate to $E_0^{s_\alpha w}$.
- (ii) For $k \in [l_1, l_2]$, $E_k^{s_\alpha w}$ is totally isotropic $\iff \text{ext}(i, j), \text{ext}(i, j)^* \notin E_k^w; \text{int}(i, j), \text{int}(i, j)^* \in E_k^w$; and for every $r \in \{1, \dots, 2n\} \setminus \{i, j, i^*, j^*\}$, the set E_k^w contains exactly one element from the pair r, r^* . \square

8.7. Completion of the proof of (7.6.1). We now commence the proof proper that $\text{Perm}^{\text{sp}}(\mu) \subset \text{Adm}^\circ(\mu)$. We assume from now on that w is μ -spin-permissible and not a translation element in \widetilde{W} , and we must find an affine root α such that $s_\alpha w$ is μ -spin-permissible and $w < s_\alpha w$.

To say that w is not a translation element is precisely to say that some element in $\{1, \dots, 2n\}$ is *proper*; let us denote by a the minimal proper element in $\{1, \dots, 2n\}$. Then

$$E_0^w = E_1^w = \dots = E_{a-1}^w \neq E_a^w.$$

Since a is proper $\iff a^*$ is proper, we have $a \leq n$, and a^* is the maximal proper element in $\{1, \dots, 2n\}$. As usual, we have $K_a = [\tilde{a}, a)$ for some $\tilde{a} \neq a$; and

our minimality assumption implies $a < \tilde{a}$. We claim $\tilde{a} \neq a^*$. For suppose to the contrary that $\tilde{a} = a^*$. Since a^* is the maximal proper element, we have $E_{a^*}^w = E_0^w$. Hence E_{a^*} is totally isotropic. But $E_{a^*}^w = (a, a^*) \cdot E_{a^*-1}^w$ (8.3.1). Hence $E_{a^*-1}^w$ is totally isotropic too but not W° -conjugate to $E_{a^*}^w$, in violation of the spin condition.

Since $\tilde{a} \neq a^*$, (8.5.2), applied with $r = a$, immediately furnishes an affine root α such that $s_\alpha w$ at least is GL -permissible and $w < s_\alpha w$. Unfortunately, in general, $s_\alpha w$ need not satisfy the spin condition. To modify our choice of α if necessary, we shall need to set up a case analysis.

Since a is proper, the set $E_a^w \setminus E_{a-1}^w$ consists of a single proper element b , and $K_b = [a, b)$. Of course $b \neq a$; and it follows from the inequality $\tilde{a} \neq a^*$ that $b \neq a^*$. By minimality of a , we thus have $a < m < m^* < a^*$, where $m := \min\{b, b^*\}$. Since E_{a-1}^w is totally isotropic, it must contain b^* , and we conclude $b, b^* \in E_k^w$ for all $k \in [a, m)$. Note that by taking $r = b$ in (8.5.2), we again get an affine root α such that $s_\alpha w$ is GL -permissible and $w < s_\alpha w$, but we again have the problem that $s_\alpha w$ may not satisfy the spin condition.

We shall base our case analysis on the existence of proper elements in $[a, m)$ that satisfy certain conditions. Note that if $i \in [a, m)$ is proper, then \tilde{i} and \tilde{i}^* are proper too. Hence $a \leq \tilde{i}, \tilde{i}^* \leq a^*$.

8.7.A. *Case: There exists a proper $r \in [a, m)$ such that $\tilde{r} \neq r^*$ and $\min\{\tilde{r}, \tilde{r}^*\} < m$.* Then (8.5.2), applied to the element r , furnishes an affine root α such that $s_\alpha w$ is GL -permissible and $w < s_\alpha w$. To see that $s_\alpha w$ satisfies the spin condition, write $l_1 < l_2 < l_3 < l_4$ for the elements $r, r^*, \tilde{r}, \tilde{r}^*$ in increasing order. Our case assumption implies $[l_1, l_2) \subset [a, m)$. Hence $b, b^* \in E_k^w$ for all $k \in [l_1, l_2)$. Hence, by (8.6.2), $E_k^{s_\alpha w}$ is not totally isotropic for such k and $s_\alpha w$ satisfies the spin condition.

In the remaining two cases we shall assume there exists no proper $r \in [a, m)$ as in (8.7.A). Hence for every proper $i \in [a, m)$ with $\tilde{i} \neq i^*$, we have $m \leq \tilde{i}, \tilde{i}^* \leq m^*$. In particular, we have $m \leq \tilde{a}, \tilde{a}^* \leq m^*$, so that $a, a^* \notin E_k^w$ for all $k \in [a, m)$.

8.7.B. *Case: There exists no r as in (8.7.A), and there exists a proper $l \in [a, m)$ distinct from a .* In this case we have

$$a < l < m \leq \tilde{a}, \tilde{a}^* \leq m^* < l^* < a^*.$$

We consider the possibilities $\tilde{l} = l^*$ and $\tilde{l} \neq l^*$ separately.

If $\tilde{l} = l^*$, then $K_l = [l^*, l)$ and $K_{l^*} = [l, l^*)$. Plainly $a, a^* \in K_l$ and $\tilde{a}, \tilde{a}^* \notin K_l$. Hence $s_{a,l;0}w$ and $s_{l,a^*;-1}w$ are GL -permissible and $w < s_{a,l;0}w, s_{l,a^*;-1}w$ (8.5.1). Moreover, since K_l and K_{l^*} are disjoint, it is immediate from (8.6.2) that $s_{a,l;0}w$ and $s_{l,a^*;-1}w$ both satisfy the spin condition. We remark that similar reasoning reveals that one can also use either of the reflections $s_{b,l^*;0}$ or $s_{b^*,l^*;0}$.

If $\tilde{l} \neq l^*$, then $m \leq \tilde{l}, \tilde{l}^* \leq m^*$ by our case assumption, and we take $\alpha := \alpha_{l,m;0}$. Plainly

$$l \in [a, m) \subset K_m \quad \text{and} \quad \tilde{l}^* - 1 \in [a, m^*) \subset K_{m^*}.$$

By duality, the second displayed containment implies $\tilde{l} \notin K_m$. Hence $s_\alpha w$ is GL -permissible and $w < s_\alpha w$ (8.5.1). To check the spin condition, recall that $a, a^* \notin E_k^w$ for all $k \in [a, m) \supset [l, m)$. Hence, by (8.6.2), $E_k^{s_\alpha w}$ is not totally isotropic for such k , and $s_\alpha w$ is μ -spin-permissible.

Having dispensed with the above two cases, we are left with just the following case to consider.

8.7.C. *Case: a is the only proper element in $[a, m]$.* By taking $r = b$ in (8.5.2), we have that $s_\alpha w$ is GL -permissible and $w < s_\alpha w$ for $\alpha := \alpha_{a,b;0}$ or $\alpha := \alpha_{a,b;-1}$. Thus we reduce to proving the claim:

If $s_\alpha w$ is GL -permissible for $\alpha \in \{\alpha_{a,b;0}, \alpha_{a,b;-1}\}$, then $s_\alpha w$ is spin-permissible.

So suppose we have such an α . Our minimality assumption on a and our case assumption together imply

$$E_0^w = E_1^w = \cdots = E_{a-1}^w \neq E_a^w = E_{a+1}^w = \cdots = E_{m-1}^w.$$

But this places us exactly in the situation of (8.6.1), with $i = a$ and $j = b$. Hence for either possible α , we have equalities

$$E_0^{s_\alpha w} = E_1^{s_\alpha w} = \cdots = E_{m-1}^{s_\alpha w}.$$

Hence $s_\alpha w$ is μ -spin-permissible by (8.6.2).

This completes our case analysis, and with it the proof of (7.6.1). \square

Remark 8.7.1. Implicit in our proof is a slight simplification of part of the proof [KR, 5.8] of the main result for GL_n in Kottwitz's and Rapoport's paper. Indeed, our Lemma 8.5.2, formulated without the requirement that $\tilde{r} \neq r^*$, continues to hold in the GL_n setting. So, using the language of [KR], if $w \in \widetilde{W}_{GL_n}$ has minuscule associated alcove \mathbf{v} and is not a translation element, then there must exist a proper $r \in \{1, \dots, n\}$, and the lemma immediately furnishes an α such that $s_\alpha \mathbf{v}$ is minuscule and $w < s_\alpha w$. On the other hand, [KR] actually proves a little more: namely, that α can always be chosen to satisfy the additional constraint that the translation parts of w and $s_\alpha w$ are the same. We can find such an α by letting a denote the *minimal* proper element in $\{1, \dots, n\}$; then $\alpha_{a,\tilde{a};0}$ or $\alpha_{a,\tilde{a};-1}$ does the job.

As noted by Kottwitz and Rapoport, it follows that

$$(*) \quad \begin{array}{l} w \in \widetilde{W}_{GL_n} \text{ is } \mu\text{-admissible} \\ \text{for minuscule } \mu \end{array} \implies \begin{array}{l} w \text{ is less than or equal to its} \\ \text{translation part in the Bruhat order.} \end{array}$$

Much more generally, Haines [H, proof of 4.6], using Hecke algebra techniques, has shown that (*) continues to hold when \widetilde{W}_{GL_n} is replaced by the extended affine Weyl group attached to any root datum. Unfortunately, the arguments in this paper do not seem to yield a direct proof of (*) for \widetilde{W} and $\mu \in \{\mu_1, \mu_2\}$.

8.8. Permissibility and spin-permissibility. We conclude the paper by showing that Kottwitz's and Rapoport's notion of μ -permissibility [KR] agrees with our notion of μ -spin-permissibility for elements in \widetilde{W} . While we have only defined μ -spin-permissibility for $\mu \in \{\mu_1, \mu_2\}$, the notion of μ -permissibility makes sense for any cocharacter μ : quite generally, $w \in \widetilde{W}$ is μ -permissible if $w \equiv t_\mu \pmod{W_a}$ and $wx - x \in \text{Conv}(W^\circ \mu)$ for all x in \tilde{A} , where $\text{Conv}(W^\circ \mu)$ is the convex hull in $X_*(T) \otimes \mathbb{R}$ of the W° -conjugates of μ , and \tilde{A} is the alcove in $X_*(T) \otimes \mathbb{R}$ obtained as the inverse image of A . Of course, it is equivalent to require $wx - x \in \text{Conv}(W^\circ \mu)$ for all x in the closure of \tilde{A} . We denote by $\text{Perm}(\mu)$ the set of μ -permissible elements.

Proposition 8.8.1. $\text{Perm}^{\text{sp}}(\mu) = \text{Perm}(\mu)$ for $\mu \in \{\mu_1, \mu_2\}$.

Proof. The containment \subset follows from the equality $\text{Perm}^{\text{sp}}(\mu) = \text{Adm}^\circ(\mu)$ (7.6.1) and the general result [KR, 11.2] that μ -admissibility implies μ -permissibility for any cocharacter μ in any extended affine Weyl group attached to a root datum. (Note that while \widetilde{W} is not the extended affine Weyl group attached to a root datum, \widetilde{W}° is, and the sets in question are all contained in \widetilde{W}° .) To prove the reverse containment, suppose $w \in \text{Perm}(\mu)$. Since $\text{Conv}(W^\circ\mu)$ is contained in $\text{Conv}(W\mu)$ (this is the relevant convex hull that comes up for $GS_{p_{2n}}$, [KR, 12.4] shows, modulo conventions, that (P1) and (P2) hold for w . It remains to show that if the vector μ_k^w (7.3.1) is totally isotropic for $0 \leq k \leq n$, then $\mu_k^w \in W^\circ\mu$. For any k , since $\frac{\omega_k + \omega_{2n-k}}{2}$ is in the closure of \widetilde{A} , we have

$$\frac{\mu_k^w + \mu_{2n-k}^w}{2} \in \text{Conv}(W^\circ\mu).$$

But if μ_k^w is totally isotropic, then $\mu_k^w = \mu_{2n-k}^w$ and the displayed vector equals μ_k^w . Now use the obvious fact that $X_*(T) \cap \text{Conv}(W^\circ\mu) = W^\circ\mu$. \square

Although we didn't need it for the proof, it is not hard to give an explicit description of the convex hull $\text{Conv}(W^\circ\mu)$. We set $V := X_*(T) \otimes \mathbb{R}$, and we identify it with

$$\{(x_1, \dots, x_{2n}) \in \mathbb{R}^{2n} \mid x_1 + x_{2n} = x_2 + x_{2n-1} = \dots = x_n + x_{n+1}\}.$$

For $x = (x_1, \dots, x_{2n}) \in V$, we write $c(x)$ for the common value $x_1 + x_{2n} = \dots = x_n + x_{n+1}$. We write $x \cdot y$ for the usual dot product of vectors in \mathbb{R}^{2n} . Then it is readily verified that, when n is odd,

$$\text{Conv}(W^\circ\mu) = \left\{ x \in V \mid \begin{array}{l} (0, \dots, 0) \leq x \leq (1, \dots, 1), c(x) = 1, \\ \text{and } \mu' \cdot x \geq 1 \text{ for all } \mu' \in W^\circ\mu \end{array} \right\};$$

and when n is even,

$$\text{Conv}(W^\circ\mu) = \left\{ x \in V \mid \begin{array}{l} (0, \dots, 0) \leq x \leq (1, \dots, 1), c(x) = 1, \\ \text{and } \mu' \cdot x \geq 1 \text{ for all } \mu' \in \tau W^\circ\mu \end{array} \right\},$$

where τ is the usual element from §5.1.

Remark 8.8.2. For $i = 1, 2$, let Y_i denote the common set

$$\text{Adm}^\circ(\mu_i) = \text{Perm}^{\text{sp}}(\mu_i) = \text{Perm}(\mu_i).$$

Using a subscript GL_{2n} to denote the corresponding notions for elements in $\widetilde{W}_{GL_{2n}}$, let Z denote the common set

$$\text{Adm}_{GL_{2n}}(\mu_1) = \text{Adm}_{GL_{2n}}(\mu_2) = \text{Perm}_{GL_{2n}}(\mu_1) = \text{Perm}_{GL_{2n}}(\mu_2);$$

here we use the equivalence between admissibility and permissibility for minuscule cocharacters in GL_{2n} due to Kottwitz and Rapoport [KR, 3.5]. Then we have relations between Y_1 , Y_2 , and Z ,

$$Y_1 \amalg Y_2 \subsetneq Z \cap \widetilde{W}^\circ \subsetneq Z \cap \widetilde{W}.$$

(Recall that we always assume $n \geq 2$; here the first \subsetneq becomes an equality when $n = 1$.)

REFERENCES

- [A] K. Arzdorf, *On local models with special parahoric level structure*, Michigan Math. J. **58** (2009), no. 3, 683–710.
- [BT] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Inst. Hautes Études Sci. Publ. Math. (1984), no. 60, 197–376 (French).
- [G1] U. Görtz, *On the flatness of models of certain Shimura varieties of PEL-type*, Math. Ann. **321** (2001), no. 3, 689–727.
- [G2] ———, *On the flatness of local models for the symplectic group*, Adv. Math. **176** (2003), no. 1, 89–115.
- [G3] ———, *Computing the alternating trace of Frobenius on the sheaves of nearby cycles on local models for GL_4 and GL_5* , J. Algebra **278** (2004), no. 1, 148–172.
- [G4] ———, *Topological flatness of local models in the ramified case*, Math. Z. **250** (2005), no. 4, 775–790.
- [H] T. J. Haines, *Test functions for Shimura varieties: the Drinfeld case*, Duke Math. J. **106** (2001), no. 1, 19–40.
- [HN1] T. J. Haines and Ngô B. C., *Nearby cycles for local models of some Shimura varieties*, Compositio Math. **133** (2002), no. 2, 117–150.
- [HN2] ———, *Alcoves associated to special fibers of local models*, Amer. J. Math. **124** (2002), no. 6, 1125–1152.
- [HR] T. Haines and M. Rapoport, *On parahoric subgroups*, appendix to: G. Pappas and M. Rapoport, *Twisted loop groups and their affine flag varieties*, Adv. Math. **219** (2008), no. 1, 118–198.
- [IM] N. Iwahori and H. Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups*, Inst. Hautes Études Sci. Publ. Math. (1965), no. 25, 5–48.
- [KR] R. Kottwitz and M. Rapoport, *Minuscule alcoves for GL_n and GSp_{2n}* , Manuscripta Math. **102** (2000), no. 4, 403–428.
- [K] N. Krämer, *Local models for ramified unitary groups*, Abh. Math. Sem. Univ. Hamburg **73** (2003), 67–80.
- [P] G. Pappas, *On the arithmetic moduli schemes of PEL Shimura varieties*, J. Algebraic Geom. **9** (2000), no. 3, 577–605.
- [PR1] G. Pappas and M. Rapoport, *Local models in the ramified case. I. The EL-case*, J. Algebraic Geom. **12** (2003), no. 1, 107–145.
- [PR2] ———, *Local models in the ramified case. II. Splitting models*, Duke Math. J. **127** (2005), no. 2, 193–250.
- [PR3] ———, *Twisted loop groups and their affine flag varieties*, with an appendix by T. Haines and Rapoport, Adv. Math. **219** (2008), no. 1, 118–198.
- [PR4] ———, *Local models in the ramified case. III. Unitary groups*, J. Inst. Math. Jussieu **8** (2009), no. 3, 507–564.
- [R] M. Rapoport, *A guide to the reduction modulo p of Shimura varieties*, Astérisque (2005), no. 298, 271–318 (English, with English and French summaries). Automorphic forms. I.
- [RZ] M. Rapoport and Th. Zink, *Period spaces for p -divisible groups*, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996.
- [S1] B. D. Smithling, *Admissibility and permissibility for minuscule cocharacters in orthogonal groups* (March 12, 2010), preprint, [arXiv:1001.0937v2](https://arxiv.org/abs/1001.0937v2) [math.AG].
- [S2] ———, *Topological flatness of local models for ramified unitary groups. I. The odd dimensional case* (February 18, 2010), preprint, [arXiv:1002.3520v1](https://arxiv.org/abs/1002.3520v1) [math.AG].
- [S3] ———, *Topological flatness of local models for ramified unitary groups. II. The even dimensional case*, in preparation.

UNIVERSITY OF TORONTO, DEPARTMENT OF MATHEMATICS, 40 ST. GEORGE ST., TORONTO, ON M5S 2E4, CANADA

E-mail address: bds@math.toronto.edu