

## CONSEQUENCES OF THE CAUCHY'S INTEGRAL FORMULA



UNIVERSITY OF  
**TORONTO**

October 16<sup>th</sup>, 2020 to October 21<sup>st</sup>, 2020

# Holomorphic functions are analytic<sup>1</sup> – 1

## Theorem

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be holomorphic/analytic/ $\mathbb{C}$ -differentiable. Then  $f$  can be **locally** expressed as a power series in a neighborhood of any point of  $U$ .

More precisely, if  $\overline{D_r}(z_0) \subset U$  then

$$\forall z \in D_r(z_0), f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$$

where

$$a_n = \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{(w - z_0)^{n+1}} dw$$

and  $\gamma : [0, 1] \rightarrow \mathbb{C}$  is defined by  $\gamma(t) = z_0 + re^{2i\pi t}$  and the radius of convergence of this power series is greater than or equal to  $r$ .

---

<sup>1</sup>I will stop complaining about the term *analytic*... Now we know that holomorphic and analytic are synonyms.

# Holomorphic functions are analytic – 2

*Proof.* By Cauchy's integral formula

$$\begin{aligned} f(z) &= \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{w-z} dw \\ &= \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{w-z_0} \frac{1}{1 - \frac{z-z_0}{w-z_0}} dw \\ &= \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{w-z_0} \sum_{n=0}^{+\infty} \left( \frac{z-z_0}{w-z_0} \right)^n dw && \text{since } |z-z_0| < |w-z_0| = r \\ &= \sum_{n=0}^{+\infty} \left( \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{(w-z_0)^{n+1}} dw \right) (z-z_0)^n \end{aligned}$$

Actually, we should justify the permutation  $\int - \sum$  more carefully.

It can be done by noting that for  $\varepsilon > 0$  there exists  $N$  such that if  $k \geq N$  then the remainder satisfies  $\left| \sum_{n=k}^{+\infty} \left( \frac{z-z_0}{w-z_0} \right)^n \right| \leq \varepsilon$ , so that

$$\left| f(z) - \sum_{n=0}^k \left( \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{(w-z_0)^{n+1}} dw \right) (z-z_0)^n \right| \leq \left| \frac{1}{2i\pi} \int_{\gamma} \frac{f(w)}{w-z_0} \sum_{n=k}^{+\infty} \left( \frac{z-z_0}{w-z_0} \right)^n dw \right| \leq \frac{\max_{|w-z_0|=r} |f|}{r} \varepsilon \text{Length}(\gamma) \leq \varepsilon \max_{|w-z_0|=r} |f|.$$

# Holomorphic functions are analytic – 3

## Corollary

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be a function.

Then  $f$  is holomorphic/analytic/ $\mathbb{C}$ -differentiable if and only if  $f$  can be **locally** expressed as a power series in a neighborhood of any point of  $U$ .

*Proof.*

$\Rightarrow$ : the main theorem from the previous slide.

$\Leftarrow$ : if  $f(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n$  in  $D_r(z_0)$  then  $f$  is holomorphic on  $D_r(z_0)$  from last week lecture.



## Corollary

A holomorphic/analytic/ $\mathbb{C}$ -differentiable function is infinitely many times  $\mathbb{C}$ -differentiable.

# Holomorphic functions are analytic – 4

The previous results are false for  $\mathbb{R}$ -differentiability.

- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by 
$$f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

Then  $f$  is  $\mathbb{R}$ -differentiable but not  $C^1$ .

- Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by 
$$g(x) = \begin{cases} e^{-\frac{1}{x^2}} & \text{if } x \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

Then  $g$  is  $\mathbb{R}$ -differentiable, even  $C^\infty$ , but not analytic at 0,

i.e. it can't be expressed as a power series around 0:

Indeed  $\forall n \in \mathbb{N}_{\geq 0}$ ,  $g^{(n)}(0) = 0$ , so if  $g$  were equal to its Taylor series around 0 then it would be constant equal to 0 but  $g$  is non-zero in any neighborhood of 0.

- By a theorem of Borel<sup>2</sup>, given a real sequence  $(a_n)_{n \in \mathbb{N}_{\geq 0}}$ , there exists a  $C^\infty$  function defined in a neighborhood of 0 in  $\mathbb{R}$  such that  $\forall n \in \mathbb{N}_{\geq 0}$ ,  $f^{(n)}(0) = a_n$ .  
Otherwise stated, any real power series is the Taylor expansion of a  $C^\infty$  function.  
If we take  $a_n = (n!)^2$  then we obtain a power series whose radius of convergence is  $R = 0$ , hence a function with such a Taylor expansion can't be analytic.

---


<sup>2</sup>It generalizes to multivariable functions.

# Continuation of analytic functions – 1

## Theorem

Let  $U \subset \mathbb{C}$  be a **domain** and  $f : U \rightarrow \mathbb{C}$  be a holomorphic/analytic function. If there exists  $z_0 \in U$  such that  $\forall n \in \mathbb{N}_{\geq 0}$ ,  $f^{(n)}(z_0) = 0$  then  $f \equiv 0$  on  $U$ .

*Proof.*



Let  $z \in U$ . Since  $U$  is path connected, there exists a  $C^0$  curve  $\gamma : [a, b] \rightarrow U$  from  $\gamma(a) = z_0$  to  $\gamma(b) = z$ . Since  $U$  is open, for every  $w \in \gamma([a, b])$  there exists  $r_w > 0$  such that  $D_{r_w}(w) \subset U$ . Since  $\gamma([a, b])$  is compact we may assume that it is covered by finitely many of these disks  $D_{r_1}(w_1), \dots, D_{r_k}(w_k)$ .

By the theorem on Slide 2, if there exists  $v \in D_{r_i}(w_i)$  such that  $\forall n \in \mathbb{N}_{\geq 0}$ ,  $f^{(n)}(v) = 0$  then  $f \equiv 0$  on  $D_{r_i}(w_i)$ . Two consecutive disks intersect (since they cover  $\gamma$ ), so we conclude using the previous remark disk by disk from  $z_0$  to  $z$ . ■

*If you attend MAT327, another proof consists in showing that  $\{z \in U : \forall n \in \mathbb{N}_{\geq 0}, f^{(n)}(z) = 0\}$  is open, closed, and non-empty, hence equals to  $U$  by connectedness.*

### Corollary

Let  $U \subset \mathbb{C}$  be a **domain** and  $f, g : U \rightarrow \mathbb{C}$  be holomorphic/analytic functions.  
If  $f$  and  $g$  coincide in the neighborhood of a point,

$$\text{i.e. } \exists z_0 \in U, \exists r > 0, \forall z \in D_r(z_0) \cap U, f(z) = g(z),$$

then they coincide on  $U$ ,

$$\text{i.e. } \forall z \in U, f(z) = g(z).$$

*Proof.* Then  $\forall n \in \mathbb{N}_{\geq 0}$ ,  $(f - g)^{(n)}(z_0) = 0$  (since  $f - g \equiv 0$  on  $D_r(z_0) \cap U$ ).  
Hence  $f - g \equiv 0$  on  $U$  by the previous theorem. ■

The previous results are false for  $\mathbb{R}$ -differentiability.

## Example

Define  $f : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f(x) = \begin{cases} e^{-\frac{1}{x}} & \text{if } x > 0 \\ 0 & \text{otherwise} \end{cases}$$

then  $f$  is  $\mathbb{R}$ -differentiable (even  $C^\infty$ ). And

- $\forall n \in \mathbb{N}_{\geq 0}, f^{(n)}(0) = 0$  but  $f \not\equiv 0$ .
- $\forall x \in (-\infty, 0), f(x) = 0$  but  $f \not\equiv 0$ .



# Continuation of analytic functions – 4

A common way to construct an analytic function consists in defining it in a "small" domain and then to extend it to an analytic function with a bigger domain.

By the above result, this analytic continuation, if it exists, is unique.

That's a very powerful tool: knowing a function on a "small" domain determines the function everywhere else<sup>3</sup>. *Holomorphic/analytic functions are very rigid!*

⚠ The maximal domain may not be  $\mathbb{C}$ : for instance, if we try to extend  $\text{Log}$ , we won't be able to do a full turn around the origin since we won't recover the same values (it increases by  $2i\pi$ ).

## Example

Let  $f : D_1(0) \rightarrow \mathbb{C}$  be defined by  $f(z) = \sum_{n=0}^{+\infty} z^n$ .

Then  $f$  coincides with  $\frac{1}{1-z}$  on  $D_1(0)$ .

Hence we may extend  $f$  with  $F : \mathbb{C} \setminus \{1\} \rightarrow \mathbb{C}$  defined by  $F(z) = \frac{1}{1-z}$ .

<sup>3</sup>And we will even weaken the assumptions later this term: it is enough to know the function on a set with a limit point.

# Order of a zero

## Definition: order of a zero

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be holomorphic/analytic. Let  $z_0 \in U$  be such that  $f(z_0) = 0$ . We define the **order of vanishing of  $f$  at  $z_0$**  by  $m_f(z_0) := \min \{n \in \mathbb{N} : f^{(n)}(z_0) \neq 0\}$ .

*Note that  $m_f(z_0) > 0$  since  $f(z_0) = 0$ .*

## Proposition

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be holomorphic/analytic. Let  $z_0 \in U$  be such that  $f(z_0) = 0$ .

Denote the power series expansion of  $f$  at  $z_0$  by  $f(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n$ .

Then  $m_f(z_0) = \min \{n \in \mathbb{N} : a_n \neq 0\}$ .

*Proof.*  $a_n = \frac{f^{(n)}(z_0)}{n!}$

## Proposition

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be holomorphic/analytic. Then  $z_0$  is a zero of order  $n$  of  $f$  if and only if there exists  $g : U \rightarrow \mathbb{C}$  holomorphic such that  $f(z) = (z - z_0)^n g(z)$  and  $g(z_0) \neq 0$ .

# Morera's theorem

## Theorem: Morera's theorem

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be continuous.

If for every (full) triangle  $T$  lying in  $U$  we have  $\int_{\partial T} f = 0$  then  $f$  is holomorphic/analytic on  $U$ .

*Proof.* Let  $z_0 \in U$  and  $r > 0$  be such that  $D_r(z_0) \subset U$ . We define  $F : D_r(z_0) \rightarrow \mathbb{C}$  by  $F(z) = \int_{[z_0, z]} f$ .

Let  $z, h \in \mathbb{C}$  be such that  $z, z+h \in D_r(z_0)$  then, considering the triangle whose vertices are  $z_0, z$  and  $z+h$ , we obtain  $\int_{[z_0, z]} f + \int_{[z, z+h]} f + \int_{[z+h, z_0]} f = 0$ , i.e.  $F(z+h) - F(z) = \int_{[z, z+h]} f$ . Then

$$\begin{aligned} \left| \frac{F(z+h) - F(z)}{h} - f(z) \right| &= \left| \frac{\left( \int_{[z, z+h]} f \right) - hf(z)}{h} \right| = \frac{1}{|h|} \left| \int_{[z, z+h]} (f(w) - f(z)) dw \right| \leq \frac{\sup_{w \in [z, z+h]} |f(w) - f(z)|}{|h|} \text{Length}([z, z+h]) \\ &= \sup_{w \in [z, z+h]} |f(w) - f(z)| \xrightarrow{h \rightarrow 0} 0 \end{aligned}$$

Hence  $F$  is holomorphic on  $D_r(z_0)$  and  $F' = f$ . Furthermore,  $f$  is holomorphic on  $D_r(z_0)$  (and hence at  $z_0$ ) as the complex derivative of a holomorphic function. ■

# Theorem: characterizations of holomorphicity/analyticity

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$ . Then the following are equivalent:

- ①  $f$  is holomorphic/analytic/ $\mathbb{C}$ -differentiable, i.e.  $\forall z_0 \in U, \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$  exists.
- ②  $\tilde{f} : \tilde{U} \rightarrow \mathbb{R}^2$  is  $\mathbb{R}$ -differentiable and satisfies the Cauchy–Riemann equations on  $\tilde{U}$ .
- ③  $f$  may be written as a power series  $f(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n$  on a neighborhood of every  $z_0 \in U$ .
- ④  $f$  is continuous and for every (full) triangle  $T$  lying in  $U$  we have  $\int_{\partial T} f = 0$ .
- ⑤  $f$  is continuous and for every simple closed curve  $\gamma$  on  $U$  whose inside is also included in  $U$ , we have  $\int_{\gamma} f = 0$ .
- ⑥  $f$  admits local primitives/antiderivatives: for every  $z_0 \in U$  there exists  $F : D_r(z_0) \cap U \rightarrow \mathbb{C}$  holomorphic for some  $r > 0$  such that  $F' = f$  on  $D_r(z_0) \cap U$ .

When  $U$  is simply-connected, we may drop the assumption that the inside of the triangle/curve is included in  $U$  in ④ ⑤ furthermore we may also replace "local primitives/antiderivatives" by "a primitive/antiderivative" in ⑥ i.e. there exists  $F : U \rightarrow \mathbb{C}$  holomorphic s.t.  $F' = f$ .

# Liouville's theorem – 1

## Definition

We say that a function  $f : \mathbb{C} \rightarrow \mathbb{C}$  is **entire** if it is holomorphic (everywhere) on  $\mathbb{C}$ .

## Theorem: Liouville's theorem

A bounded entire function is constant

# Liouville's theorem – 2

## Lemma: Cauchy's inequalities

Let  $U \subset \mathbb{C}$  be open and  $f : U \rightarrow \mathbb{C}$  be holomorphic. Let  $r > 0$ . If  $D_r(z_0) \subset U$  then

$$|f^{(n)}(z_0)| \leq \frac{n!}{r^n} \max_{|z-z_0|=r} |f(z)|$$

*Proof.* From the theorem on Slide 2

$$\left| \frac{f^{(n)}(z_0)}{n!} \right| = \left| \frac{1}{2i\pi} \int_{|z-z_0|=r} \frac{f(w)}{(w-z_0)^{n+1}} dw \right| \leq \frac{\max_{|z-z_0|=r} |f(z)|}{r^{n+1}} \frac{\text{Length}(|z-z_0|=r)}{2\pi} = \frac{\max_{|z-z_0|=r} |f(z)|}{r^n}$$

*Proof of Liouville's theorem.*

Assume that there exists  $M > 0$  such that  $\forall z \in \mathbb{C}, |f(z)| \leq M$ .

Let  $z \in \mathbb{C}$ . For  $r > 0$ , by Cauchy's inequality with  $n = 1$ , we have  $|f'(z)| \leq \frac{M}{r} \xrightarrow{r \rightarrow +\infty} 0$ .

Hence,  $\forall z \in \mathbb{C}, f'(z) = 0$ . Therefore  $f$  is constant on  $\mathbb{C}$ .

### Theorem: d'Alembert–Gauss theorem or the Fundamental Theorem of Algebra

A non-constant polynomial with coefficients in  $\mathbb{C}$  admits a root/zero in  $\mathbb{C}$ .

*Proof.* Assume that  $P$  is a complex polynomial with no root in  $\mathbb{C}$ .

Then  $Q = \frac{1}{P}$  is an entire function and  $Q$  is bounded since  $\lim_{z \rightarrow +\infty} Q = 0$ .

Therefore, by Liouville's theorem,  $Q$  is constant, and so is  $P$ . ■

## Theorem

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f : U \rightarrow \mathbb{C}$  a holomorphic/analytic function which doesn't vanish, i.e.  $\forall z \in U, f(z) \neq 0$ .


Then there exists  $g : U \rightarrow \mathbb{C}$  holomorphic/analytic such that  $e^g = f$ .

*Proof.* Since  $f$  doesn't vanish,  $\frac{f'}{f}$  is holomorphic on  $U$ .

Then, since  $U$  is simply connected,  $\frac{f'}{f}$  admits a complex primitive/antiderivative, i.e. there exists  $\tilde{g} : U \rightarrow \mathbb{C}$  holomorphic/analytic such that  $\tilde{g}' = \frac{f'}{f}$ .

Then  $(f e^{-\tilde{g}})' = (f' - \tilde{g}' f) e^{-\tilde{g}} = 0$  and therefore  $f e^{-\tilde{g}} = K$  is constant since  $U$  is connected.

Since  $K \neq 0$ , there exists  $w \in \mathbb{C}$  such that  $e^w = K$ .

Then, for  $g = \tilde{g} + w$ , we have  $e^g = f$ . 

## Careful

Such a function  $g$  is not unique! For instance  $g + 2i\pi$  is another suitable function.



# Multiplication of complex power series – 1

We are going to prove the following theorem, but using results related to analytic functions.

## Theorem

Let  $S_A(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n$  and  $S_B(z) = \sum_{n=0}^{+\infty} b_n(z - z_0)^n$  be two power series of radii  $R_A$  and  $R_B$ .

We define  $S_{AB}(z) := \sum_{n=0}^{+\infty} \left( \sum_{k=0}^n a_k b_{n-k} \right) (z - z_0)^n$  and denote its radius of convergence by  $R_{AB}$ .

Then

①  $R_{AB} \geq \min(R_A, R_B)$ .

② If  $|z - z_0| < \min(R_A, R_B)$  then  $S_{AB}(z) = S_A(z)S_B(z)$ ,

i.e. 
$$\sum_{n=0}^{+\infty} \left( \sum_{k=0}^n a_k b_{n-k} \right) (z - z_0)^n = \left( \sum_{n=0}^{+\infty} a_n (z - z_0)^n \right) \left( \sum_{n=0}^{+\infty} b_n (z - z_0)^n \right).$$

# Multiplication of complex power series – 1

*Proof.* We know that  $S_A$  and  $S_B$  are holomorphic on  $z \in \mathbb{C}$  such that  $|z - z_0| < \min(R_A, R_B)$ .

Then  $f(z) = S_A(z)S_B(z)$  is holomorphic on  $|z - z_0| < \min(R_A, R_B)$ , so it can be written as a

power series  $f(z) = \sum_{n=0}^{+\infty} c_n(z - z_0)^n$  on  $|z - z_0| < \min(R_A, R_B)$  (particularly its radius of convergence is at least  $\min(R_A, R_B)$ ).

Then

$$\begin{aligned} c_n &= \frac{f^{(n)}(z_0)}{n!} = \frac{1}{n!} \sum_{k=0}^n \frac{n!}{(n-k)!k!} S_A^{(n-k)}(z_0) S_B^{(k)}(z_0) \text{ by Leibniz rule} \\ &= \sum_{k=0}^n \frac{S_A^{(n-k)}(z_0)}{(n-k)!} \frac{S_B^{(k)}(z_0)}{k!} \\ &= \sum_{k=0}^n a_{n-k} b_k = \sum_{k=0}^n a_k b_{n-k} \end{aligned}$$