## MAT246H1-S – LEC0201/9201 Concepts in Abstract Mathematics

# NATURAL NUMBERS – 1



## January 12<sup>th</sup>, 2021

What is  $\mathbb{N}$ ?





- There is an initial element: 0 "zero".
- Every natural number *n* has a successor s(n), intuitively n + 1.



- There is an initial element: 0 "zero".
- Every natural number *n* has a *successor* s(n), intuitively n + 1.



- There is an initial element: 0 "zero".
- Every natural number *n* has a *successor* s(n), intuitively n + 1.

• • • • • • 0 is not the successor of another natural number



- There is an initial element: 0 "zero".
- Every natural number *n* has a successor s(n), intuitively n + 1.



0 is not the successor of another natural number

A natural number can't be the successor of two distinct natural numbers



- There is an initial element: 0 "zero".
- Every natural number *n* has a successor s(n), intuitively n + 1.



0 is not the successor of another natural number

A natural number can't be the successor of two distinct natural numbers

The successor can't loop back to a previous natural number



- There is an initial element: 0 "zero".
- Every natural number *n* has a successor s(n), intuitively n + 1.



0 is not the successor of another natural number

A natural number can't be the successor of two distinct natural numbers

The successor can't loop back to a previous natural number

The successor of a natural number is uniquely defined



- There is an initial element: 0 "zero".
- Every natural number *n* has a successor s(n), intuitively n + 1.





- There is an initial element: 0 "zero".
- Every natural number *n* has a successor s(n), intuitively n + 1.



So we need extra assumptions on 0 and *s*.

# Peano's axioms - 1

We admit the following theorem (we have to start from somewhere...).

#### Theorem: Peano axioms

There exists a set  $\mathbb{N}$  together with an element  $0 \in \mathbb{N}$  "zero" and a function  $s : \mathbb{N} \to \mathbb{N}$  "successor" such that:

**1** 0 is not the successor of any element of  $\mathbb{N}$ , i.e. 0 is not in the image of *s*:

 $0 \notin s(\mathbb{N})$ 

2 If the successor of *n* equals the successor of *m* then n = m, i.e. *s* is injective:

 $\forall n, m \in \mathbb{N}, \ s(n) = s(m) \implies n = m$ 

**3** The induction principle. If a subset of  $\mathbb{N}$  contains 0 and is closed under s then it is  $\mathbb{N}$ :

$$\forall A \subset \mathbb{N}, \left\{ \begin{array}{c} 0 \in A \\ s(A) \subset A \end{array} \right. \implies A = \mathbb{N}$$

# Peano's axioms - 1

We admit the following theorem (we have to start from somewhere...).

### Theorem: Peano axioms

There exists a set  $\mathbb{N}$  together with an element  $0 \in \mathbb{N}$  "zero" and a function  $s : \mathbb{N} \to \mathbb{N}$  "successor" such that:

**1** 0 is not the successor of any element of  $\mathbb{N}$ , i.e. 0 is not in the image of *s*:

 $0 \not\in s(\mathbb{N})$ 

2 If the successor of *n* equals the successor of *m* then n = m, i.e. *s* is injective:

 $\forall n, m \in \mathbb{N}, \ s(n) = s(m) \implies n = m$ 

**3** The induction principle. If a subset of  $\mathbb{N}$  contains 0 and is closed under s then it is  $\mathbb{N}$ :

$$\forall A \subset \mathbb{N}, \left\{ \begin{array}{c} 0 \in A \\ s(A) \subset A \end{array} \right. \implies A = \mathbb{N}$$

The last point means we obtain  $\mathbb{N}$  taking iteratively the successor starting at 0 (we don't miss any element).

- That's why we can define a sequence  $(u_n)_{n \in \mathbb{N}}$  iteratively by fixing  $u_0$  and setting  $u_{n+1} = f(u_n)$ .
- It is closely related to the notion of proof by induction.

### Remarks

- We can't use + and × yet, because they are still not defined. Currently, we can only use the three Peano's axioms from above.
- Intuitively, s(n) = n + 1 (which will become true after we define the addition).
- All the results about N will derive from these three basic properties (and even more since we will construct the other sets Z, Q, R, C from N).

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

Proof. Set  $A = \{n \in \mathbb{N} : n \neq s(n)\}.$ 

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

Proof. Set  $A = \{n \in \mathbb{N} : n \neq s(n)\}$ . Then •  $A \subset \mathbb{N}$ 

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

#### Proof.

#### Set $A = \{n \in \mathbb{N} : n \neq s(n)\}$ . Then

- $A \subset \mathbb{N}$
- $0 \in A$

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

#### Proof.

Set 
$$A = \{n \in \mathbb{N} : n \neq s(n)\}$$
. Then

- $A \subset \mathbb{N}$
- $0 \in A$  since  $0 \neq s(0)$  (remember that  $0 \notin s(\mathbb{N})$ ).

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

#### Proof.

Set 
$$A = \{n \in \mathbb{N} : n \neq s(n)\}$$
. Then

- $A \subset \mathbb{N}$
- $0 \in A$  since  $0 \neq s(0)$  (remember that  $0 \notin s(\mathbb{N})$ ).

•  $s(A) \subset A$ 

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

#### Proof.

```
Set A = \{n \in \mathbb{N} : n \neq s(n)\}. Then
```

- $A \subset \mathbb{N}$
- $0 \in A$  since  $0 \neq s(0)$  (remember that  $0 \notin s(\mathbb{N})$ ).

```
• s(A) \subset A
Indeed, let m \in s(A). Then m = s(n) for some n \in A. So s(n) \neq n.
Since s is injective, we get that s(s(n)) \neq s(n), i.e. s(m) \neq m.
Hence m \in A.
```

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

#### Proof.

```
Set A = \{n \in \mathbb{N} : n \neq s(n)\}. Then
```

```
• A \subset \mathbb{N}
```

•  $0 \in A$  since  $0 \neq s(0)$  (remember that  $0 \notin s(\mathbb{N})$ ).

```
• s(A) \subset A
Indeed, let m \in s(A). Then m = s(n) for some n \in A. So s(n) \neq n.
Since s is injective, we get that s(s(n)) \neq s(n), i.e. s(m) \neq m.
Hence m \in A.
```

So, by the induction principle,  $A = \mathbb{N}$ .

A natural number is never its own successor, i.e.

 $\forall n \in \mathbb{N}, n \neq s(n)$ 

#### Proof.

```
Set A = \{n \in \mathbb{N} : n \neq s(n)\}. Then
```

```
• A \subset \mathbb{N}
```

•  $0 \in A$  since  $0 \neq s(0)$  (remember that  $0 \notin s(\mathbb{N})$ ).

```
• s(A) \subset A
Indeed, let m \in s(A). Then m = s(n) for some n \in A. So s(n) \neq n.
Since s is injective, we get that s(s(n)) \neq s(n), i.e. s(m) \neq m.
Hence m \in A.
```

So, by the induction principle,  $A = \mathbb{N}$ . Thus, for every  $n \in \mathbb{N}$  we have that  $n \neq s(n)$ .