

# *Concepts in Abstract Mathematics*

## REAL NUMBERS – 2




UNIVERSITY OF  
TORONTO

March 16<sup>th</sup>, 2021

# Characterization of the supremum

## Proposition


Let  $A \subset \mathbb{R}$  and  $M \in \mathbb{R}$ . Then  $M = \sup(A) \Leftrightarrow \begin{cases} \forall x \in A, x \leq M \\ \forall \varepsilon > 0, \exists x \in A, M - \varepsilon < x \end{cases}$



# Characterization of the supremum

## Proposition

Let  $A \subset \mathbb{R}$  and  $M \in \mathbb{R}$ . Then  $M = \sup(A) \Leftrightarrow \begin{cases} \forall x \in A, x \leq M \\ \forall \varepsilon > 0, \exists x \in A, M - \varepsilon < x \end{cases}$



*Proof.*

$\Rightarrow$  Assume that  $M = \sup(A)$ . Then  $M$  is an upper bound of  $A$  so  $\forall x \in A, x \leq M$ .

We know that if  $T$  is an other upper bound of  $A$  then  $M \leq T$  (since  $M$  is the least upper bound).

So, by taking the contrapositive, if  $T < M$  then  $T$  isn't an upper bound of  $A$ .

Let  $\varepsilon > 0$ . Since  $M - \varepsilon < M$ , we know that  $M - \varepsilon$  is not an upper bound of  $A$ , meaning that there exists  $x \in A$  such that  $M - \varepsilon < x$ .

$\Leftarrow$  Assume that  $\begin{cases} \forall x \in A, x \leq M \\ \forall \varepsilon > 0, \exists x \in A, M - \varepsilon < x \end{cases}$

Then, by the first condition,  $M$  is an upper bound of  $A$ . Let's prove it is the least one.

We will show the contrapositive: if  $T < M$  then  $T$  isn't an upper bound of  $A$ .

Let  $T \in \mathbb{R}$ . Assume that  $T < M$ . Set  $\varepsilon = M - T > 0$ . Then there exists  $x \in A$  such that  $M - \varepsilon < x$ , i.e.  $T < x$ .


Hence  $T$  isn't an upper bound of  $A$ .

# Characterization of the infimum

## Proposition

Let  $A \subset \mathbb{R}$  and  $m \in \mathbb{R}$ . Then

$$m = \inf(A) \Leftrightarrow \begin{cases} \forall x \in A, m \leq x \\ \forall \varepsilon > 0, \exists x \in A, x < m + \varepsilon \end{cases}$$



# $\mathbb{R}$ is archimedean

Theorem:  $\mathbb{R}$  is archimedean

$$\forall \varepsilon > 0, \forall A > 0, \exists n \in \mathbb{N}, n\varepsilon > A$$

## Theorem: $\mathbb{R}$ is archimedean

$\forall \varepsilon > 0, \forall A > 0, \exists n \in \mathbb{N}, n\varepsilon > A$

*Proof.*

Let  $\varepsilon > 0$  and  $A > 0$ .

Assume by contradiction that  $\forall n \in \mathbb{N}, n\varepsilon \leq A$ .

Then  $E = \{n\varepsilon : n \in \mathbb{N}\}$  is non-empty and bounded from above so it admits a supremum  $M = \sup E$  by the least upper bound principle.

Since  $M - \varepsilon < M$ ,  $M - \varepsilon$  is not an upper bound of  $E$ , so there exists  $n \in \mathbb{N}$  such that  $n\varepsilon > M - \varepsilon$ .  
Therefore  $(n + 1)\varepsilon > M$ , hence a contradiction. ■

## Theorem: $\mathbb{R}$ is archimedean

$$\forall \varepsilon > 0, \forall A > 0, \exists n \in \mathbb{N}, n\varepsilon > A$$

*Proof.*

Let  $\varepsilon > 0$  and  $A > 0$ .

Assume by contradiction that  $\forall n \in \mathbb{N}, n\varepsilon \leq A$ .

Then  $E = \{n\varepsilon : n \in \mathbb{N}\}$  is non-empty and bounded from above so it admits a supremum  $M = \sup E$  by the least upper bound principle.

Since  $M - \varepsilon < M$ ,  $M - \varepsilon$  is not an upper bound of  $E$ , so there exists  $n \in \mathbb{N}$  such that  $n\varepsilon > M - \varepsilon$ . Therefore  $(n + 1)\varepsilon > M$ , hence a contradiction. ■

The above theorem means that  $\lim_{n \rightarrow +\infty} \frac{1}{n} = 0$ , or equivalently that  $\mathbb{R}$  doesn't contain infinitesimal elements (i.e. there is not infinitely large or infinitely small elements).

# Floor function

## Proposition: floor function

For every  $x \in \mathbb{R}$ , there exists a unique  $n \in \mathbb{Z}$  such that  $n \leq x < n + 1$ .

We say that  $n$  is the integer part (or the floor function value) of  $x$  and we denote it by  $\lfloor x \rfloor$ .



## Proposition: floor function

For every  $x \in \mathbb{R}$ , there exists a unique  $n \in \mathbb{Z}$  such that  $n \leq x < n + 1$ .

We say that  $n$  is the integer part (or the floor function value) of  $x$  and we denote it by  $\lfloor x \rfloor$ .

*Proof.* Let  $x \in \mathbb{R}$ .

### Existence.

- *First case: if  $x \geq 0$ .*

We set  $E = \{n \in \mathbb{N} : x < n\}$ .

By the archimedean property (with  $\varepsilon = 1$ ), there exists  $m \in \mathbb{N}$  such that  $m > x$ . Hence  $E \neq \emptyset$ .

By the well-ordering principle,  $E$  admits a least element  $p$ .

We have that  $x < p$  since  $p \in E$  and that  $p - 1 \leq x$  since  $p - 1 \notin E$ .

Therefore  $n = p - 1$  satisfies  $n \leq x < n + 1$ .

- *Second case: if  $x < 0$ .* Then we apply the first case to  $-x$ .

**Uniqueness.** Assume that  $n, n' \in \mathbb{Z}$  are two suitable integers. Then (1)  $n \leq x < n + 1$  and  $n' \leq x < n' + 1$ .

We deduce from the last inequality that (2)  $-n' - 1 < -x \leq -n'$ .

Summing (1) and (2), we get that  $n - n' - 1 < 0 < n - n' + 1$ .

Hence  $n - n' < 1$ , i.e.  $n - n' \leq 0$ , and  $-1 < n - n'$  i.e.  $0 \leq n - n'$ .

Therefore  $n = n'$ .