## BLOWUP PROPERTIES FOR CRITICAL AND SUPERCRITICAL NLS AT LOW REGULARITY

#### J. Colliander

Toronto

56th Midwest Partial Differential Equations Seminar

Cauchy Problem Scaling

#### CRITICAL REGIMES & LOW REGULARITY GWP?

Mass Critical Case  $H^{1/2}$  Critical Case Energy Critical Case Energy Supercritical Case

#### CRITICAL NORM EXPLOSION FOR $H^{1/2}$ CRITICAL CASE

Heuristics Contradiction Strategy Mass Freezing

글 🖌 🖌 글

-

A B + 
 A B +
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A

Consider the initial value problem  $NLS_{p}^{\pm}(\mathbb{R}^{d})$ :

$$\begin{cases} i\partial_t u + \Delta u = \pm |u|^{p-1}u\\ u(0,x) = u_0(x) \end{cases}$$

∃ ► < ∃</p>

Consider the initial value problem  $NLS_p^{\pm}(\mathbb{R}^d)$ :

$$\begin{cases} i\partial_t u + \Delta u = \pm |u|^{p-1} u \\ u(0, x) = u_0(x) \end{cases}$$

We seek  $u: (-T_*, T^*) \times \mathbb{R}^d \mapsto \mathbb{C}$ . (+ focusing, - defocusing)

Consider the initial value problem  $NLS_p^{\pm}(\mathbb{R}^d)$ :

$$\begin{cases} i\partial_t u + \Delta u = \pm |u|^{p-1} u \\ u(0, x) = u_0(x) \end{cases}$$

We seek 
$$u: (-T_*, T^*) \times \mathbb{R}^d \mapsto \mathbb{C}$$
.  
(+ focusing, - defocusing)

Time Invariant Quantities

$$\begin{aligned} \mathsf{Mass} &= \|u(t)\|_{L^2_x}\\ \mathsf{Hamiltonian} &= \int_{R^d} |\nabla u(t)|^2 dx \mp \frac{2}{p+1} |u(t)|^{p+1} dx \end{aligned}$$

• 3 • 4 3

イロト イポト イヨト イヨト

• If *u* solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-T_*, T^*) \times \mathbb{R}^2$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-\lambda^2 T_*, \lambda^2 T^*) \times \mathbb{R}^2$ .

▶ If *u* solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-T_*, T^*) \times \mathbb{R}^2$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-\lambda^2 T_*, \lambda^2 T^*) \times \mathbb{R}^2$ .

 Dilation invariant norms play decisive role in the theory of NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>) :

- A I I A I I A I I A

▶ If *u* solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-T_*, T^*) \times \mathbb{R}^2$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-\lambda^2 T_*, \lambda^2 T^*) \times \mathbb{R}^2$ .

 Dilation invariant norms play decisive role in the theory of NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>) :

- A I I A I I A I I A

▶ If *u* solves  $NLS^{\pm}_{p}(\mathbb{R}^{d})$  on  $(-T_{*}, T^{*}) \times \mathbb{R}^{2}$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-\lambda^2 T_*, \lambda^2 T^*) \times \mathbb{R}^2$ .

 Dilation invariant norms play decisive role in the theory of NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>) :

$$\|D_y^{\sigma}u_{\lambda}\|_{L^q(\mathbb{R}^d_y)} = \left(\frac{1}{\lambda}\right)^{\frac{2}{p-1}+\sigma-\frac{d}{q}} \|D_x^{\sigma}u\|_{L^q(\mathbb{R}^d_x)}.$$

- A I I A I I A I I A

▶ If *u* solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-T_*, T^*) \times \mathbb{R}^2$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-\lambda^2 T_*, \lambda^2 T^*) \times \mathbb{R}^2$ .

 Dilation invariant norms play decisive role in the theory of NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>) :

$$\|D_y^{\sigma}u_{\lambda}\|_{L^q(\mathbb{R}^d_y)} = \left(\frac{1}{\lambda}\right)^{\frac{2}{p-1}+\sigma-\frac{d}{q}} \|D_x^{\sigma}u\|_{L^q(\mathbb{R}^d_x)}.$$

 $\dot{W}^{\sigma,q}$  is critical if  $\frac{2}{p-1} + \sigma - \frac{d}{q} = 0$ .

▶ If *u* solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-T_*, T^*) \times \mathbb{R}^2$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS_p^{\pm}(\mathbb{R}^d)$  on  $(-\lambda^2 T_*, \lambda^2 T^*) \times \mathbb{R}^2$ .

 Dilation invariant norms play decisive role in the theory of *NLS*<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>) :

$$\|D_y^{\sigma}u_{\lambda}\|_{L^q(\mathbb{R}^d_y)} = \left(\frac{1}{\lambda}\right)^{\frac{2}{p-1}+\sigma-\frac{d}{q}} \|D_x^{\sigma}u\|_{L^q(\mathbb{R}^d_x)}.$$

$$\dot{W}^{\sigma,q}$$
 is critical if  $\frac{2}{p-1} + \sigma - \frac{d}{q} = 0$ .  
 $\sim NLS_p^{\pm}(\mathbb{R}^d)$  is  $\dot{H}^{s_c}$ -critical for  $s_c := \frac{d}{2} - \frac{2}{p-1}$ .

▶ If *u* solves  $NLS^{\pm}_{p}(\mathbb{R}^{d})$  on  $(-T_{*}, T^{*}) \times \mathbb{R}^{2}$  then

$$u_{\lambda}(\tau, y) := \lambda^{\frac{2}{1-p}} u(\tau \lambda^{-2}, y \lambda^{-1})$$

solves  $NLS^{\pm}_{p}(\mathbb{R}^{d})$  on  $(-\lambda^{2}T_{*},\lambda^{2}T^{*})\times\mathbb{R}^{2}$ .

 Dilation invariant norms play decisive role in the theory of NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>) :

$$\|D_y^{\sigma}u_{\lambda}\|_{L^q(\mathbb{R}^d_y)} = \left(\frac{1}{\lambda}\right)^{\frac{2}{p-1}+\sigma-\frac{d}{q}} \|D_x^{\sigma}u\|_{L^q(\mathbb{R}^d_x)}.$$

 $\dot{W}^{\sigma,q}$  is critical if  $\frac{2}{p-1} + \sigma - \frac{d}{q} = 0$ .

- $NLS_p^{\pm}(\mathbb{R}^d)$  is  $\dot{H}^{s_c}$ -critical for  $s_c := \frac{d}{2} \frac{2}{p-1}$ .
- ▶  $L^2$  and  $\dot{H}^1$  critical cases distinguished by conservation laws.

イロト イポト イヨト イヨト

• Theory for  $NLS_p^{\pm}(\mathbb{R}^d)$  is qualitatively similar in regimes:

- Theory for  $NLS^{\pm}_{\rho}(\mathbb{R}^d)$  is qualitatively similar in regimes:
  - ▶ Mass subcritical (*s<sub>c</sub>* < 0)

向下 イヨト イヨト

- Theory for  $NLS^{\pm}_{\rho}(\mathbb{R}^d)$  is qualitatively similar in regimes:
  - ▶ Mass subcritical (*s<sub>c</sub>* < 0)
  - ▶ Mass critical (s<sub>c</sub> = 0)

向下 イヨト イヨト

- Theory for  $NLS^{\pm}_{p}(\mathbb{R}^{d})$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - Mass critical  $(s_c = 0)$
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )

- - E > - E

- Theory for  $NLS^{\pm}_{\rho}(\mathbb{R}^d)$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - Mass critical (s<sub>c</sub> = 0)
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )
  - Energy critical (s<sub>c</sub> = 1)

- Theory for  $NLS^{\pm}_{\rho}(\mathbb{R}^d)$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - Mass critical (s<sub>c</sub> = 0)
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )
  - Energy critical (s<sub>c</sub> = 1)
  - Energy supercritical (s<sub>c</sub> > 1).

- Theory for  $NLS^{\pm}_{\rho}(\mathbb{R}^d)$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - ▶ Mass critical (s<sub>c</sub> = 0)
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )
  - Energy critical (s<sub>c</sub> = 1)
  - Energy supercritical (s<sub>c</sub> > 1).

• Optimal local-in-time well-posedness (LWP) for  $NLS_p^{\pm}(\mathbb{R}^d)$ :

- Theory for  $NLS^{\pm}_{\rho}(\mathbb{R}^d)$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - ▶ Mass critical (s<sub>c</sub> = 0)
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )
  - Energy critical (s<sub>c</sub> = 1)
  - Energy supercritical (s<sub>c</sub> > 1).

• Optimal local-in-time well-posedness (LWP) for  $NLS_p^{\pm}(\mathbb{R}^d)$ :

- Theory for  $NLS^{\pm}_{p}(\mathbb{R}^{d})$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - Mass critical (s<sub>c</sub> = 0)
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )
  - Energy critical (s<sub>c</sub> = 1)
  - Energy supercritical (s<sub>c</sub> > 1).
- Optimal local-in-time well-posedness (LWP) for NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>):
   ∀ s ≥ max(0, s<sub>c</sub>) ∃ unique continuous data-to-solution map

$$H^{s} \ni u_{0} \longmapsto u \in C([0, T_{lwp}]; H^{s}) \cap L^{q}_{t}L^{p}_{x}$$

with  $T_{lwp} = T_{lwp}(||u_0||_{H^s})$  if  $s > s_c$  and  $T_{lwp} = T(u_0)$  if  $s = s_c$ .

(日) (同) (E) (E) (E) (E)

- Theory for  $NLS^{\pm}_{p}(\mathbb{R}^{d})$  is qualitatively similar in regimes:
  - Mass subcritical (s<sub>c</sub> < 0)</li>
  - ▶ Mass critical (s<sub>c</sub> = 0)
  - Mass supercritical/Energy subcritical ( $0 < s_c < 1$ )
  - Energy critical (s<sub>c</sub> = 1)
  - Energy supercritical (s<sub>c</sub> > 1).
- Optimal local-in-time well-posedness (LWP) for NLS<sup>±</sup><sub>p</sub>(ℝ<sup>d</sup>):
   ∀ s ≥ max(0, s<sub>c</sub>) ∃ unique continuous data-to-solution map

$$H^{s} \ni u_{0} \longmapsto u \in C([0, T_{lwp}]; H^{s}) \cap L^{q}_{t}L^{p}_{x}$$

with  $T_{lwp} = T_{lwp}(||u_0||_{H^s})$  if  $s > s_c$  and  $T_{lwp} = T(u_0)$  if  $s = s_c$ .

Optimal maximal-in-time well-posedness (GWP) is known only in the defocusing energy critical case. What is the fate of local-in-time solutions with critical initial regularity?

CRITICAL REGIMES & LOW REGULARITY GWP?

## LOW REGULARITY POLEMICS

イロン イヨン イヨン

Why should we care about low regularity data?

글 🖌 🖌 글

-

Why should we care about low regularity data?

Envelope equation derivation of NLS is "band limited".

Why should we care about low regularity data?

- Envelope equation derivation of NLS is "band limited".
- Physically relevant solutions are smooth.

Why should we care about low regularity data?

- Envelope equation derivation of NLS is "band limited".
- Physically relevant solutions are smooth.

Some answers:

Why should we care about low regularity data?

- Envelope equation derivation of NLS is "band limited".
- Physically relevant solutions are smooth.

Some answers:

► ∃ global-in-time classical solutions of energy critical NLS was unknown until finite energy solutions were shown to be global.

Why should we care about low regularity data?

- Envelope equation derivation of NLS is "band limited".
- Physically relevant solutions are smooth.

Some answers:

- ► ∃ global-in-time classical solutions of energy critical NLS was unknown until finite energy solutions were shown to be global.
- Typical, slightly bigger than necessary, finite time blowup solutions of mass critical NLS with nice initial data are "L<sup>2</sup>-typical" after removing the blowup.

Why should we care about low regularity data?

- Envelope equation derivation of NLS is "band limited".
- Physically relevant solutions are smooth.

Some answers:

- ► ∃ global-in-time classical solutions of energy critical NLS was unknown until finite energy solutions were shown to be global.
- Typical, slightly bigger than necessary, finite time blowup solutions of mass critical NLS with nice initial data are "L<sup>2</sup>-typical" after removing the blowup.
- Invariant measures live on low regularity phase space.

Why should we care about low regularity data?

- Envelope equation derivation of NLS is "band limited".
- Physically relevant solutions are smooth.

Some answers:

- ► ∃ global-in-time classical solutions of energy critical NLS was unknown until finite energy solutions were shown to be global.
- Typical, slightly bigger than necessary, finite time blowup solutions of mass critical NLS with nice initial data are "L<sup>2</sup>-typical" after removing the blowup.
- Invariant measures live on low regularity phase space.

The talk will first discuss ideas and questions in the  $L^2$  mass critical case, the  $H^{1/2}$  critical case, and the  $H^1$  energy critical case. Then, I'll discuss a qualitative property of blowup in an  $H^{1/2}$  critical case.

・ロト ・同ト ・ヨト ・ヨト

# Typical blowups leave an $L^2$ stain at time $T^*$

メロト メタト メヨト メヨト

# Typical blowups leave an $L^2$ stain at time $T^*$

[Merle-Raphaël]

-

(4) (3) (4) (4) (4)

A B + 
 A B +
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
### Typical blowups leave an $L^2$ stain at time $T^*$

[Merle-Raphaël]

▶ Consider  $H^1 \cap \{ \|Q\|_{L^2} < \|u_0\|_{L^2} < \|Q\|_{L^2} + \alpha^* \} \ni u_0 \longmapsto u$  solving  $NLS_3^-(\mathbb{R}^2)$  on  $[0, T^*)$  (maximal) with  $T^* < \infty$ .

### Typical blowups leave an $L^2$ stain at time $T^*$

[Merle-Raphaël]

- ▶ Consider  $H^1 \cap \{ \|Q\|_{L^2} < \|u_0\|_{L^2} < \|Q\|_{L^2} + \alpha^* \} \ni u_0 \longmapsto u$  solving  $NLS_3^-(\mathbb{R}^2)$  on  $[0, T^*)$  (maximal) with  $T^* < \infty$ .
- ▶  $\exists \ \lambda(t), x(t), \theta(t) \in \mathbb{R}^+, \mathbb{R}^2, \mathbb{R}/(2\pi\mathbb{Z})$  and  $u^*$  such that

$$u(t) - \lambda(t)^{-1} Q\left(rac{x-x(t)}{\lambda(t)}
ight) e^{i heta(t)} 
ightarrow u^*$$

strongly in  $L^2(\mathbb{R}^2)$ .

### Typical blowups leave an $L^2$ stain at time $T^*$

[Merle-Raphaël]

- ▶ Consider  $H^1 \cap \{ \|Q\|_{L^2} < \|u_0\|_{L^2} < \|Q\|_{L^2} + \alpha^* \} \ni u_0 \longmapsto u$  solving  $NLS_3^-(\mathbb{R}^2)$  on  $[0, T^*)$  (maximal) with  $T^* < \infty$ .
- ▶  $\exists \ \lambda(t), x(t), \theta(t) \in \mathbb{R}^+, \mathbb{R}^2, \mathbb{R}/(2\pi\mathbb{Z})$  and  $u^*$  such that

$$u(t) - \lambda(t)^{-1} Q\left(rac{x-x(t)}{\lambda(t)}
ight) e^{i heta(t)} 
ightarrow u^*$$

strongly in  $L^2(\mathbb{R}^2)$ .

▶ Typically,  $u^* \notin H^s \cup L^p$  for s > 0, p > 2!

#### MASS CRITICAL CASE

### $L^2$ CRITICAL CASE: LWP THEORY

イロト イポト イヨト イヨト

3 × 4 3 ×

[Cazenave-Weissler]

(E) (E)

[Cazenave-Weissler]

▶  $\forall u_0 \in L^2$  there exists  $T_{lwp}(u_0)$  determined by

$$\|e^{it\Delta}u_0\|_{L^4_{tx}}([0, T_{lwp}] imes \mathbb{R}^2) < rac{1}{100}.$$

 $\exists \text{ unique solution } u \in C([0, T_{lwp}]; L^2) \cap L^4_{tx}([0, T_{lwp}] \times \mathbb{R}^2).$ 

[Cazenave-Weissler]

▶  $\forall u_0 \in L^2$  there exists  $T_{lwp}(u_0)$  determined by

$$\|e^{it\Delta}u_0\|_{L^4_{tx}}([0, T_{lwp}] imes \mathbb{R}^2) < rac{1}{100}.$$

 $\exists \text{ unique solution } u \in C([0, T_{lwp}]; L^2) \cap L^4_{tx}([0, T_{lwp}] \times \mathbb{R}^2).$ 

• Define the maximal forward existence time  $T^*(u_0)$  by

$$\|u\|_{L^4_{tx}([0,T^*-\delta]\times\mathbb{R}^2)}<\infty$$

for all  $\delta > 0$  but diverges to  $\infty$  as  $\delta \downarrow 0$ .

[Cazenave-Weissler]

▶  $\forall u_0 \in L^2$  there exists  $T_{lwp}(u_0)$  determined by

$$\|e^{it\Delta}u_0\|_{L^4_{tx}}([0, T_{lwp}] imes \mathbb{R}^2) < rac{1}{100}.$$

 $\exists \text{ unique solution } u \in C([0, T_{lwp}]; L^2) \cap L^4_{tx}([0, T_{lwp}] \times \mathbb{R}^2).$ 

• Define the maximal forward existence time  $T^*(u_0)$  by

$$\|u\|_{L^4_{tx}([0,T^*-\delta]\times\mathbb{R}^2)}<\infty$$

for all  $\delta > 0$  but diverges to  $\infty$  as  $\delta \downarrow 0$ .

▶ ∃ small data scattering threshold  $\mu_0 > 0$ 

$$||u_0||_{L^2} < \mu_0 \implies ||u||_{L^4_{tx}(\mathbb{R} \times \mathbb{R}^2)} < 2\mu_0.$$

#### MASS CRITICAL CASE

### $L^2$ CRITICAL CASE: GWP THEORY

イロト イポト イヨト イヨト

▶  $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .

イロト イヨト イヨト

#### MASS CRITICAL CASE

### $L^2$ Critical Case: GWP Theory

- $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .
- ►  $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

A B N A B N

**A** ►

#### MASS CRITICAL CASE

### $L^2$ Critical Case: GWP Theory

- $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .
- ►  $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

A B N A B N

**A** ►

- ▶  $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .
- $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies H^1 \ni u_0 \longmapsto u, \, T^* = \infty.$$

[Weinstein]

- E - - E -

▶  $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .

•  $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies H^1 \ni u_0 \longmapsto u, T^* = \infty.$$

[Weinstein]

Here Q is the ground state solution to  $-Q + \Delta Q = Q^3$ .  $e^{it}Q(x)$  is the ground state solution solution to  $NLS_3^-(\mathbb{R}^2)$ .

▶  $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .

•  $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies H^1 \ni u_0 \longmapsto u, T^* = \infty.$$

[Weinstein]

Here Q is the ground state solution to  $-Q + \Delta Q = Q^3$ .  $e^{it}Q(x)$  is the ground state solution solution to  $NLS_3^-(\mathbb{R}^2)$ .

► 'I Method' yields  $H^s$ -GWP for  $s > \frac{4}{7}$   $(s > \frac{1}{2}$ ?). [CKSTT]

▶  $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .

•  $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies H^1 \ni u_0 \longmapsto u, T^* = \infty.$$

[Weinstein]

Here Q is the ground state solution to  $-Q + \Delta Q = Q^3$ .  $e^{it}Q(x)$  is the ground state solution solution to  $NLS_3^-(\mathbb{R}^2)$ .

► 'I Method' yields  $H^s$ -GWP for  $s > \frac{4}{7}$   $(s > \frac{1}{2}$ ?). [CKSTT]

# 1<sup>2</sup> Critical Case: GWP Theory

- ▶  $H^1$ -GWP for  $NLS_3^+(\mathbb{R}^2)$ .
- $H^1$ -GWP mass threshold  $||Q||_{L^2}$  for  $NLS_3^-(\mathbb{R}^2)$ :

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies H^1 \ni u_0 \longmapsto u, T^* = \infty.$$

### [Weinstein]

Here Q is the ground state solution to  $-Q + \Delta Q = Q^3$ .  $e^{it}Q(x)$  is the ground state solution solution to  $NLS_3^-(\mathbb{R}^2)$ .

*'I Method'* yields H<sup>s</sup>-GWP for s > <sup>4</sup>/<sub>7</sub> (s > <sup>1</sup>/<sub>2</sub>?). [CKSTT] NLS<sup>+</sup><sub>5</sub>(ℝ<sup>1</sup>) is similarly H<sup>s</sup>-GWP for s > <sup>4</sup>/<sub>9</sub>. [Tzirakis] NLS<sup>+</sup><sub><sup>4</sup>/<sub>4</sub>+1</sub>(ℝ<sup>d</sup>) is H<sup>s</sup>-GWP for s > <sup>d+8</sup>/<sub>d+10</sub>. [Visan-Zhang]

### $L^2$ CRITICAL CASE: BLOWUP SOLUTION PROPERTIES

-

(3)

A B + 
 A B +
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A

### $L^2$ CRITICAL CASE: BLOWUP SOLUTION PROPERTIES

**Explicit Blowup Solutions** 

• 3 • 4 3

#### **Explicit Blowup Solutions**

• Arise as *pseudoconformal* image of  $e^{it}Q(x)$ :

$$S(t,x) = \frac{1}{t}Q\left(\frac{x}{t}\right)e^{-i\frac{|x|^2}{4t}+\frac{i}{t}}.$$

#### **Explicit Blowup Solutions**

• Arise as *pseudoconformal* image of  $e^{it}Q(x)$ :

$$S(t,x) = \frac{1}{t}Q\left(\frac{x}{t}\right)e^{-i\frac{|x|^2}{4t}+\frac{i}{t}}.$$

► S has minimal mass:

$$\|S(-1)\|_{L^2_x} = \|Q\|_{L^2}.$$

All mass in S is conically concentrated into a point.

#### **Explicit Blowup Solutions**

• Arise as *pseudoconformal* image of  $e^{it}Q(x)$ :

$$S(t,x) = \frac{1}{t}Q\left(\frac{x}{t}\right)e^{-i\frac{|x|^2}{4t}+\frac{i}{t}}.$$

S has minimal mass:

$$\|S(-1)\|_{L^2_x} = \|Q\|_{L^2}.$$

All mass in S is conically concentrated into a point.

▶ Minimal mass H<sup>1</sup> blowup solution characterization: u<sub>0</sub> ∈ H<sup>1</sup>, ||u<sub>0</sub>||<sub>L<sup>2</sup></sub> = ||Q||<sub>L<sup>2</sup></sub>, T<sup>\*</sup>(u<sub>0</sub>) < ∞ implies that u = S up to an explicit solution symmetry. [Merle]</p>

### $L^2$ CRITICAL CASE: BLOWUP SOLUTION PROPERTIES

Virial Identity  $\implies \exists$  Many Blowup Solutions

( ) ) ( ) )

### $L^2$ CRITICAL CASE: BLOWUP SOLUTION PROPERTIES

### Virial Identity $\implies \exists$ Many Blowup Solutions

Integration by parts and the equation yields

$$\partial_t^2 \int_{\mathbb{R}^2_x} |x|^2 |u(t,x)|^2 dx = 8H[u_0].$$

[Vlasov-Petrischev-Talanov], [Glassey]

#### Virial Identity $\implies \exists$ Many Blowup Solutions

Integration by parts and the equation yields

$$\partial_t^2 \int_{\mathbb{R}^2_x} |x|^2 |u(t,x)|^2 dx = 8H[u_0].$$

[Vlasov-Petrischev-Talanov], [Glassey]

•  $H[u_0] < 0, \int |x|^2 |u_0(x)|^2 dx < \infty$  blows up.

### Virial Identity $\implies \exists$ Many Blowup Solutions

Integration by parts and the equation yields

$$\partial_t^2 \int_{\mathbb{R}^2_x} |x|^2 |u(t,x)|^2 dx = 8H[u_0].$$

[Vlasov-Petrischev-Talanov], [Glassey]

- $H[u_0] < 0, \int |x|^2 |u_0(x)|^2 dx < \infty$  blows up.
- How do these solutions blow up?

#### $H^1$ Theory of Mass Concentration

(4) (2) (4) (2)

A B + A B +
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A
 A

#### $H^1$ Theory of Mass Concentration

► 
$$H^1 \cap \{ radial \} \ni u_0 \longmapsto u, T^* < \infty \text{ implies}$$
  
$$\liminf_{t \uparrow T^*} \int_{|x| < (T^* - t)^{1/2-}} |u(t, x)|^2 dx \ge ||Q||_{L^2}^2.$$

[Merle-Tsutsumi]

(3)

## $I^2$ Critical Case: Mass Concentration

#### $H^1$ Theory of Mass Concentration

► 
$$H^1 \cap \{ radial \} \ni u_0 \longmapsto u, T^* < \infty$$
 implies  
$$\liminf_{t \uparrow T^*} \int_{|x| < (T^* - t)^{1/2-}} |u(t, x)|^2 dx \ge ||Q||_{L^2}^2.$$

### [Merle-Tsutsumi]

 $\blacktriangleright$   $H^1$  blowups parabolically concentrate at least the ground state mass. Explicit blowups S concentrate mass much faster.

#### $H^1$ Theory of Mass Concentration

► 
$$H^1 \cap \{ radial \} \ni u_0 \longmapsto u, T^* < \infty$$
 implies  
$$\liminf_{t \uparrow T^*} \int_{|x| < (T^* - t)^{1/2-}} |u(t, x)|^2 dx \ge ||Q||_{L^2}^2.$$

### [Merle-Tsutsumi]

- ► H<sup>1</sup> blowups parabolically concentrate at least the ground state mass. Explicit blowups S concentrate mass much faster.
- ▶ Fantastic recent progress on the H<sup>1</sup> blowup theory. [Merle-Raphaël]

#### L<sup>2</sup> Theory of Mass Concentration

**A** ►

CRITICAL REGIMES & LOW REGULARITY GWP? MASS CRITICAL CASE

## $L^2$ CRITICAL CASE: MASS CONCENTRATION

#### L<sup>2</sup> Theory of Mass Concentration

▶ 
$$L^2 \ni u_0 \longmapsto u, T^* < \infty$$
 implies

$$\limsup_{t \uparrow T^*} \sup_{cubes} \sup_{I,side(I) \le (T^*-t)^{1/2}} \int_I |u(t,x)|^2 dx \ge ||u_0||_{L^2}^{-M}.$$

^

[Bourgain]

(3)

CRITICAL REGIMES & LOW REGULARITY GWP? MASS CRITICAL CASE

## $L^2$ CRITICAL CASE: MASS CONCENTRATION

#### L<sup>2</sup> Theory of Mass Concentration

▶ 
$$L^2 \ni u_0 \longmapsto u, T^* < \infty$$
 implies

$$\limsup_{t \uparrow T^*} \sup_{cubes} \sup_{I,side(I) \le (T^*-t)^{1/2}} \int_I |u(t,x)|^2 dx \ge ||u_0||_{L^2}^{-M}.$$

^

[Bourgain]

(3)
## $L^2$ CRITICAL CASE: MASS CONCENTRATION

#### L<sup>2</sup> Theory of Mass Concentration

▶ 
$$L^2 \ni u_0 \longmapsto u, T^* < \infty$$
 implies

$$\limsup_{t \uparrow T^*} \sup_{cubes} \sup_{I,side(I) \le (T^*-t)^{1/2}} \int_I |u(t,x)|^2 dx \ge ||u_0||_{L^2}^{-M}.$$

0

[Bourgain]

 $L^2$  blowups parabolically concentrate some mass.

∃ ► 4.

## $L^2$ CRITICAL CASE: MASS CONCENTRATION

#### L<sup>2</sup> Theory of Mass Concentration

▶ 
$$L^2 \ni u_0 \longmapsto u, T^* < \infty$$
 implies

$$\limsup_{t \uparrow T^*} \sup_{cubes} \sup_{I,side(I) \le (T^*-t)^{1/2}} \int_{I} |u(t,x)|^2 dx \ge ||u_0||_{L^2}^{-M}.$$

[Bourgain]

L<sup>2</sup> blowups parabolically concentrate some mass.
For large L<sup>2</sup> data, do there exist tiny concentrations?

# $L^2$ CRITICAL CASE: MASS CONCENTRATION

#### L<sup>2</sup> Theory of Mass Concentration

▶ 
$$L^2 \ni u_0 \longmapsto u, T^* < \infty$$
 implies

$$\limsup_{t \uparrow T^*} \sup_{cubes} \sup_{I,side(I) \le (T^*-t)^{1/2}} \int_I |u(t,x)|^2 dx \ge ||u_0||_{L^2}^{-M}.$$

•

[Bourgain]

 $L^2$  blowups parabolically concentrate some mass.

- For large  $L^2$  data, do there exist tiny concentrations?
- Extensions in [Merle-Vega], [Carles-Keraani], [Bégout-Vargas].

イロト イヨト イヨト イヨト

# $L^2$ CRITICAL CASE: CONJECTURES/QUESTIONS Consider focusing $NLS_3^-(\mathbb{R}^2)$ :

向下 イヨト イヨト

Consider focusing  $NLS_3^-(\mathbb{R}^2)$ :

#### Scattering Below the Ground State Mass

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies \stackrel{???}{\Longrightarrow} u_0 \longmapsto u \text{ with } \|u\|_{L^4_{tx}} < \infty.$$

(Also,  $L^2$  solutions of  $NLS_3^+(\mathbb{R}^2)$  satisfy???  $||u||_{L^4_{tx}} < \infty$ .)

Consider focusing  $NLS_3^-(\mathbb{R}^2)$ :

### Scattering Below the Ground State Mass

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies \stackrel{???}{\longrightarrow} u_0 \longmapsto u \text{ with } \|u\|_{L^4_{tx}} < \infty.$$

(Also,  $L^2$  solutions of  $NLS_3^+(\mathbb{R}^2)$  satisfy???  $||u||_{L^4_{tv}} < \infty$ .)

Minimal Mass Blowup Characterization

$$\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \longmapsto u, T^* < \infty \implies \stackrel{???}{\Longrightarrow} u = S,$$

modulo a solution symmetry. An intermediate step would extend characterization of the minimal mass blowup solutions in  $H^s$  for s < 1.

Consider focusing  $NLS_3^-(\mathbb{R}^2)$ :

### Scattering Below the Ground State Mass

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies \stackrel{???}{\longrightarrow} u_0 \longmapsto u \text{ with } \|u\|_{L^4_{tx}} < \infty.$$

(Also,  $L^2$  solutions of  $NLS_3^+(\mathbb{R}^2)$  satisfy???  $||u||_{L^4_{tx}} < \infty$ .)

Minimal Mass Blowup Characterization

$$\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \longmapsto u, T^* < \infty \implies \stackrel{???}{\Longrightarrow} u = S,$$

modulo a solution symmetry. An intermediate step would extend characterization of the minimal mass blowup solutions in  $H^s$  for s < 1.

#### Concentrated mass amounts are quantized

The explicit blowups constructed by pseudoconformally transforming time periodic solutions with ground and excited state profiles are the only asymptotic profiles.

イロト 不得 トイヨト イヨト

Consider focusing  $NLS_3^-(\mathbb{R}^2)$ :

### Scattering Below the Ground State Mass

$$\|u_0\|_{L^2} < \|Q\|_{L^2} \implies \stackrel{???}{\longrightarrow} u_0 \longmapsto u \text{ with } \|u\|_{L^4_{tx}} < \infty.$$

(Also,  $L^2$  solutions of  $NLS_3^+(\mathbb{R}^2)$  satisfy???  $||u||_{L^4_{tx}} < \infty$ .)

Minimal Mass Blowup Characterization

$$\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \longmapsto u, T^* < \infty \implies \stackrel{???}{\Longrightarrow} u = S,$$

modulo a solution symmetry. An intermediate step would extend characterization of the minimal mass blowup solutions in  $H^s$  for s < 1.

#### Concentrated mass amounts are quantized

The explicit blowups constructed by pseudoconformally transforming time periodic solutions with ground and excited state profiles are the only asymptotic profiles.

#### Are there any general upper bounds?

For  

$$0.86 \sim \frac{1}{5}(1 + \sqrt{11}) < s < 1, H^s \cap \{radial\} \ni u_0 \longmapsto u, T^* < \infty \implies$$

$$\lim \sup_{t \uparrow T^*} \int_{|x| < (T^* - t)^{s/2-}} |u(t, x)|^2 dx \ge ||Q||_{L^2}^2.$$

*H<sup>s</sup>*-blowup solutions concentrate ground state mass. [With Raynor, Sulem and Wright]

3 ×

For  

$$0.86 \sim \frac{1}{5}(1 + \sqrt{11}) < s < 1, H^s \cap \{radial\} \ni u_0 \longmapsto u, T^* < \infty \implies$$

$$\lim_{t \to \infty} \sup_{t \to 0} \int_{0}^{t} |u(t, x)|^2 dx > ||Q||^2$$

$$\limsup_{t \uparrow T^*} \int_{|x| < (T^* - t)^{s/2-}} |u(t, x)|^2 dx \ge \|Q\|_{L^2}^2.$$

*H<sup>s</sup>*-blowup solutions concentrate ground state mass. [With Raynor, Sulem and Wright]

►  $\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \in H^s, \sim 0.86 < s < 1, T^* < \infty \implies \exists t_n \uparrow T^*$ s.t.  $u(t_n) \to Q$  in  $H^{\tilde{s}(s)}$  (mod symmetry sequence).

For  

$$0.86 \sim \frac{1}{5}(1 + \sqrt{11}) < s < 1, H^s \cap \{radial\} \ni u_0 \longmapsto u, T^* < \infty \implies$$

$$\lim_{t \to \infty} \sup_{t \to 0} \int_{0}^{t} |u(t, x)|^2 dx > ||Q||^2$$

$$\limsup_{t \uparrow T^*} \int_{|x| < (T^* - t)^{s/2-}} |u(t, x)|^2 dx \ge \|Q\|_{L^2}^2.$$

*H<sup>s</sup>*-blowup solutions concentrate ground state mass. [With Raynor, Sulem and Wright]

►  $\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \in H^s, \sim 0.86 < s < 1, T^* < \infty \implies \exists t_n \uparrow T^*$ s.t.  $u(t_n) \to Q$  in  $H^{\tilde{s}(s)}$  (mod symmetry sequence).

For  

$$0.86 \sim \frac{1}{5}(1 + \sqrt{11}) < s < 1, H^s \cap \{radial\} \ni u_0 \longmapsto u, T^* < \infty \implies$$

$$\limsup_{t \uparrow T^*} \int_{|x| < (T^* - t)^{s/2-}} |u(t, x)|^2 dx \ge \|Q\|_{L^2}^2.$$

*H<sup>s</sup>*-blowup solutions concentrate ground state mass. [With Raynor, Sulem and Wright]

▶  $\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \in H^s, \sim 0.86 < s < 1, T^* < \infty \implies \exists t_n \uparrow T^*$ s.t.  $u(t_n) \to Q$  in  $H^{\tilde{s}(s)}$  (mod symmetry sequence). For  $H^s$  blowups with  $\|u_0\|_{L^2} > \|Q\|_{L^2}, u(t_n) \rightarrow V \in H^1$  (mod symmetry sequence). [Hmidi-Keraani]

For  

$$0.86 \sim \frac{1}{5}(1 + \sqrt{11}) < s < 1, H^s \cap \{radial\} \ni u_0 \longmapsto u, T^* < \infty \implies$$

$$\limsup_{t\uparrow T^*} \int_{|x|<(T^*-t)^{s/2-}} |u(t,x)|^2 dx \ge \|Q\|_{L^2}^2.$$

*H<sup>s</sup>*-blowup solutions concentrate ground state mass. [With Raynor, Sulem and Wright]

▶  $\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \in H^s, \sim 0.86 < s < 1, T^* < \infty \implies \exists t_n \uparrow T^*$ s.t.  $u(t_n) \to Q$  in  $H^{\tilde{s}(s)}$  (mod symmetry sequence). For  $H^s$  blowups with  $\|u_0\|_{L^2} > \|Q\|_{L^2}, u(t_n) \to V \in H^1$  (mod symmetry sequence). [Hmidi-Keraani] This is an  $H^s$  analog of an  $H^1$  result of [Weinstein] which preceded the minimal  $H^1$  blowup solution characterization.

For  

$$0.86 \sim \frac{1}{5}(1 + \sqrt{11}) < s < 1, H^s \cap \{radial\} \ni u_0 \longmapsto u, T^* < \infty \implies$$

$$\limsup_{t \uparrow T^*} \int_{|x| < (T^* - t)^{s/2-}} |u(t, x)|^2 dx \ge \|Q\|_{L^2}^2.$$

*H<sup>s</sup>*-blowup solutions concentrate ground state mass. [With Raynor, Sulem and Wright]

- ▶  $\|u_0\|_{L^2} = \|Q\|_{L^2}, u_0 \in H^s, \sim 0.86 < s < 1, T^* < \infty \implies \exists t_n \uparrow T^*$ s.t.  $u(t_n) \to Q$  in  $H^{\tilde{s}(s)}$  (mod symmetry sequence). For  $H^s$  blowups with  $\|u_0\|_{L^2} > \|Q\|_{L^2}, u(t_n) \to V \in H^1$  (mod symmetry sequence). [Hmidi-Keraani] This is an  $H^s$  analog of an  $H^1$  result of [Weinstein] which preceded the minimal  $H^1$  blowup solution characterization.
- ▶ Same results for  $NLS^{-}_{\frac{4}{d}+1}(\mathbb{R}^d)$  in  $H^s$ ,  $s > \frac{d+8}{d+10}$ . [Visan-Zhang]

Spacetime norm divergence rate

$$\|u\|_{L^4_{tx}([0,t] imes \mathbb{R}^2)}\gtrsim (T^*-t)^{-eta}$$

is linked with mass concentration rate

$$\limsup_{t \uparrow T^*} \sup_{cubes \ I, side(I) \le (T^* - t)^{\frac{1}{2} + \frac{\beta}{2}}} \int_{I} |u(t, x)|^2 dx \ge \|u_0\|_{L^2}^{-M}$$

[Work in progress with Roudenko]

◆□ > ◆□ > ◆臣 > ◆臣 > ○臣 - のへで

### Consider $NLS_3^-(\mathbb{R}^3)$ . Also $L_x^3$ -Critical. Typical Case?

イロト イヨト イヨト

Consider NLS<sub>3</sub><sup>-</sup>(ℝ<sup>3</sup>). Also L<sub>x</sub><sup>3</sup>-Critical. Typical Case?
LWP theory similar to NLS<sub>3</sub><sup>±</sup>(ℝ<sup>2</sup>):

$$\mathcal{L}^{2}(\mathbb{R}^{2})\longmapsto\mathcal{H}^{1/2}(\mathbb{R}^{3})$$
  
 $\mathcal{L}^{4}_{tx}\longmapsto\mathcal{L}^{5}_{tx}.$ 

向下 イヨト イヨト

Consider NLS<sub>3</sub><sup>-</sup>(ℝ<sup>3</sup>). Also L<sub>x</sub><sup>3</sup>-Critical. Typical Case?
LWP theory similar to NLS<sub>3</sub><sup>±</sup>(ℝ<sup>2</sup>):

$$L^2(\mathbb{R}^2) \longmapsto H^{1/2}(\mathbb{R}^3)$$
  
 $L^4_{tx} \longmapsto L^5_{tx}.$ 

• There cannot be an  $H^1$ -GWP mass threshold.

Consider  $NLS_3^-(\mathbb{R}^3)$ . Also  $L_x^3$ -Critical. Typical Case? • LWP theory similar to  $NLS_3^{\pm}(\mathbb{R}^2)$ :

$$\mathcal{L}^{2}(\mathbb{R}^{2})\longmapsto\mathcal{H}^{1/2}(\mathbb{R}^{3})$$
  
 $\mathcal{L}^{4}_{tx}\longmapsto\mathcal{L}^{5}_{tx}.$ 

- There cannot be an  $H^1$ -GWP mass threshold.
- No explicit blowup solutions are known.

### $H^{1/2}$ CRITICAL CASE

Consider NLS<sub>3</sub><sup>-</sup>(ℝ<sup>3</sup>). Also L<sub>x</sub><sup>3</sup>-Critical. Typical Case?
LWP theory similar to NLS<sub>3</sub><sup>±</sup>(ℝ<sup>2</sup>):

$$\mathcal{L}^{2}(\mathbb{R}^{2})\longmapsto\mathcal{H}^{1/2}(\mathbb{R}^{3})$$
  
 $\mathcal{L}^{4}_{tx}\longmapsto\mathcal{L}^{5}_{tx}.$ 

- There cannot be an  $H^1$ -GWP mass threshold.
- ▶ No explicit blowup solutions are known.
- Virial identity  $\implies \exists$  many blowup solutions.

Consider  $NLS_3^-(\mathbb{R}^3)$ . Also  $L_x^3$ -Critical. Typical Case? • LWP theory similar to  $NLS_3^{\pm}(\mathbb{R}^2)$ :

$$\mathcal{L}^{2}(\mathbb{R}^{2})\longmapsto\mathcal{H}^{1/2}(\mathbb{R}^{3})$$
  
 $\mathcal{L}^{4}_{tx}\longmapsto\mathcal{L}^{5}_{tx}.$ 

- There cannot be an  $H^1$ -GWP mass threshold.
- No explicit blowup solutions are known.
- Virial identity  $\implies \exists$  many blowup solutions.
- ▶  $H^1 \cap \{ radial \} \ni u_0 \longmapsto u, T^* < \infty$  then for any a > 0

$$\|
abla u(t)\|_{L^2_{|x| as  $t\uparrow T^*.$$$

Thus, radial solutions must explode at the origin.

Proof.

- 22

イロン イヨン イヨン

#### Proof.

By Hamiltonian conservation,

$$\|\nabla u(t)\|_{L^2}^2 = H[u_0] + \frac{1}{2} \|u(t)\|_{L^4_{|x|a}}^4$$

イロン イヨン イヨン

#### PROOF.

By Hamiltonian conservation,

$$\|\nabla u(t)\|_{L^2}^2 = H[u_0] + \frac{1}{2} \|u(t)\|_{L^4_{|x|a}}^4$$

Inner contribution estimated using Gagliardo-Nirenberg by  $C(Mass, a) \|\nabla u(t)\|_{L^2_{|x| < a}}^3$ .

- 4 回 ト 4 ヨ ト 4 ヨ ト

#### PROOF.

By Hamiltonian conservation,

$$\|\nabla u(t)\|_{L^2}^2 = H[u_0] + \frac{1}{2} \|u(t)\|_{L^4_{|x|a}}^4.$$

Inner contribution estimated using Gagliardo-Nirenberg by  $C(Mass, a) \|\nabla u(t)\|_{L^2_{|x|<a}}^3$ . Exterior region estimated by pulling out two factors in  $L^{\infty}_x$  then using radial Sobolev to get control by  $\|u(t)\|_{L^2}^3 \|\nabla u(t)\|_{L^2}$ .

#### PROOF.

By Hamiltonian conservation,

$$\|\nabla u(t)\|_{L^2}^2 = H[u_0] + \frac{1}{2} \|u(t)\|_{L^4_{|x|a}}^4$$

Inner contribution estimated using Gagliardo-Nirenberg by  $C(Mass, a) \|\nabla u(t)\|_{L^{2}_{|x| < a}}^{3}$ . Exterior region estimated by pulling out two factors in  $L^{\infty}_{x}$  then using radial Sobolev to get control by  $\|u(t)\|_{L^{2}}^{3} \|\nabla u(t)\|_{L^{2}}$ . Absorb the exterior kinetic energy to left side

$$|
abla u(t)||^2_{L^2} \lesssim C(a, Mass[u_0], H[u_0]) + C(a, Mass[u_0]) ||
abla u(t)||^3_{L^2_{|x| < a}}$$

- 25

イロン イヨン イヨン

► Radial blowup solutions of energy subcritical NLS<sub>p</sub>(ℝ<sup>d</sup>) with p < 5 must explode at the origin.</p>

(3)

- ► Radial blowup solutions of energy subcritical NLS<sub>p</sub>(ℝ<sup>d</sup>) with p < 5 must explode at the origin.</p>
- For H<sup>1/2</sup>-critical NLS<sub>5</sub><sup>-</sup>(ℝ<sup>2</sup>), there exists H<sup>1</sup> ∩ {radial} ∋ v<sub>0</sub> → v, T<sup>\*</sup>(v<sub>0</sub>) < ∞ which blows up precisely on a circle! [Raphaël]

- ► Radial blowup solutions of energy subcritical NLS<sub>p</sub>(ℝ<sup>d</sup>) with p < 5 must explode at the origin.</p>
- For H<sup>1/2</sup>-critical NLS<sub>5</sub><sup>-</sup>(ℝ<sup>2</sup>), there exists H<sup>1</sup> ∩ {radial} ∋ v<sub>0</sub> → v, T<sup>\*</sup>(v<sub>0</sub>) < ∞ which blows up precisely on a circle! [Raphaël]
- Numerics/heuristics suggest: Finite time blowup solutions of NLS<sub>3</sub>(ℝ<sup>3</sup>) satisfy ||u(t)||<sub>L<sup>3</sup>/x</sub> ↑ ∞ as t ↑ T\*.
   [Work in progress with Raynor, Sulem, Wright]
   (Analogous to [Escauriaza-Seregin-Šverák] on Navier-Stokes)

・ 同 ト ・ ヨ ト ・ ヨ ト

- ► Radial blowup solutions of energy subcritical NLS<sub>p</sub>(ℝ<sup>d</sup>) with p < 5 must explode at the origin.</p>
- For H<sup>1/2</sup>-critical NLS<sub>5</sub><sup>-</sup>(ℝ<sup>2</sup>), there exists H<sup>1</sup> ∩ {radial} ∋ v<sub>0</sub> → v, T<sup>\*</sup>(v<sub>0</sub>) < ∞ which blows up precisely on a circle! [Raphaël]
- Numerics/heuristics suggest: Finite time blowup solutions of NLS<sub>3</sub>(ℝ<sup>3</sup>) satisfy ||u(t)||<sub>L<sup>3</sup>/2</sub> ↑ ∞ as t ↑ T\*.
   [Work in progress with Raynor, Sulem, Wright]
   (Analogous to [Escauriaza-Seregin-Šverák] on Navier-Stokes)
- H<sup>1/2</sup>-blowups parabolically concentrate in L<sup>3</sup> and H<sup>1/2</sup>. [Work in progress with Roudenko]

・ 同下 ・ ヨト ・ ヨト

# $H^1(\mathbb{R}^d), d \geq 3$ Critical Case

2

・ロト ・回ト ・ヨト ・ヨト
## $H^1(\mathbb{R}^d), d \geq 3$ Critical Case

▶ Defocusing energy critical NLS<sup>+</sup><sub>1+4/(d-2)</sub>(ℝ<sup>d</sup>), d ≥ 3 is globally well-posed and scatters in H<sup>1</sup>:

向下 イヨト イヨト

# $H^1(\mathbb{R}^d), d > 3$ Critical Case

 Defocusing energy critical NLS<sup>+</sup><sub>1+4/(d-2)</sub>(ℝ<sup>d</sup>), d ≥ 3 is globally well-posed and scatters in H<sup>1</sup>: [Bourgain], [Grillakis]: Radial Case for d = 3 [CKSTT]: d = 3 [Tao]: Radial Case for d = 4 [Ryckman-Visan], [Visan], [Tao-Visan]: d ≥ 4

### $H^1(\mathbb{R}^d), d \geq 3$ CRITICAL CASE

▶ Defocusing energy critical  $NLS^+_{1+4/(d-2)}(\mathbb{R}^d), d \ge 3$  is globally well-posed and scatters in  $H^1$ : [Bourgain], [Grillakis]: Radial Case for d = 3[CKSTT]: d = 3[Tao]: Radial Case for d = 4[Ryckman-Visan], [Visan], [Tao-Visan]:  $d \ge 4$ Induction on Energy; Interaction Morawetz; Mass Freezing

・ 同下 ・ ヨト ・ ヨト

### $H^1(\mathbb{R}^d), d \geq 3$ Critical Case

- ▶ Defocusing energy critical NLS<sup>+</sup><sub>1+4/(d-2)</sub>(ℝ<sup>d</sup>), d ≥ 3 is globally well-posed and scatters in H<sup>1</sup>: [Bourgain], [Grillakis]: Radial Case for d = 3 [CKSTT]: d = 3 [Tao]: Radial Case for d = 4 [Ryckman-Visan], [Visan], [Tao-Visan]: d ≥ 4 Induction on Energy; Interaction Morawetz; Mass Freezing
- Focusing energy critical case?

マロト イヨト イヨト

CRITICAL REGIMES & LOW REGULARITY GWP?

ENERGY CRITICAL CASE

## $H^1(\mathbb{R}^2)$ "Critical" Case

∃ 900

イロン イヨン イヨン

## $H^1(\mathbb{R}^2)$ "CRITICAL" CASE

▶  $NLS_p(\mathbb{R}^2)$  is energy subcritical for all p. Is there an "energy critical" *NLS* equation on  $\mathbb{R}^2$ ?

• 3 • 4 3

## $H^1(\mathbb{R}^2)$ "CRITICAL" CASE

- ► NLS<sub>p</sub>(ℝ<sup>2</sup>) is energy subcritical for all p. Is there an "energy critical" NLS equation on ℝ<sup>2</sup>?
- Consider the defocusing initial value problem  $NLS_{exp}(\mathbb{R}^2)$

$$\begin{cases} i\partial_t u + \Delta u = u(e^{4\pi|u|^2} - 1) \\ u(0, \cdot) = u_0(\cdot) \in H^1(\mathbb{R}^2) \end{cases}$$

向下 イヨト イヨ

# $H^1(\mathbb{R}^2)$ "Critical" Case

- ► NLS<sub>p</sub>(ℝ<sup>2</sup>) is energy subcritical for all p. Is there an "energy critical" NLS equation on ℝ<sup>2</sup>?
- Consider the defocusing initial value problem  $NLS_{exp}(\mathbb{R}^2)$

$$\begin{cases} i\partial_t u + \Delta u = u(e^{4\pi|u|^2} - 1) \\ u(0, \cdot) = u_0(\cdot) \in H^1(\mathbb{R}^2) \end{cases}$$

with Hamiltonian

$$H[u(t)] := \int_{\mathbb{R}^2} |
abla u(t,x)|^2 + \int_{\mathbb{R}^2} rac{e^{4\pi |u(t,x)|^2} - 1}{4\pi} dx.$$

CRITICAL REGIMES & LOW REGULARITY GWP?

ENERGY CRITICAL CASE

## $H^1(\mathbb{R}^2)$ "Critical" Case

∃ 900

イロン イヨン イヨン

## $H^1(\mathbb{R}^2)$ "Critical" Case

If H[u<sub>0</sub>] - M[u<sub>0</sub>] ≤ 1 then NLS<sub>exp</sub>(ℝ<sup>2</sup>) is globally well-posed. Uniform continuity of data-to-solution map fails to hold for data satisfying H[u<sub>0</sub>] - M[u<sub>0</sub>] > 1.

[Work in progress with Ibrahim, Majdoub, Masmoudi]

・ 同 ト ・ ヨ ト ・ ヨ ト

## $H^1(\mathbb{R}^2)$ "Critical" Case

- If H[u<sub>0</sub>] − M[u<sub>0</sub>] ≤ 1 then NLS<sub>exp</sub>(ℝ<sup>2</sup>) is globally well-posed. Uniform continuity of data-to-solution map fails to hold for data satisfying H[u<sub>0</sub>] − M[u<sub>0</sub>] > 1.
   [Work in progress with Ibrahim, Majdoub, Masmoudi]
- Well-posedness result relies upon Strichartz estimates, Moser-Trudinger inequality, and a log-Sobolev inequality. (Largely based on similar result for NLKG by [Ibrahim-Majdoub-Masmoudi])

イロト イポト イヨト イヨト 二日

## $H^1(\mathbb{R}^2)$ "CRITICAL" CASE

- If H[u<sub>0</sub>] − M[u<sub>0</sub>] ≤ 1 then NLS<sub>exp</sub>(ℝ<sup>2</sup>) is globally well-posed. Uniform continuity of data-to-solution map fails to hold for data satisfying H[u<sub>0</sub>] − M[u<sub>0</sub>] > 1.
   [Work in progress with Ibrahim, Majdoub, Masmoudi]
- Well-posedness result relies upon Strichartz estimates, Moser-Trudinger inequality, and a log-Sobolev inequality. (Largely based on similar result for NLKG by [Ibrahim-Majdoub-Masmoudi])
- Ill-posedness result relies upon optimizing sequence for Moser-Trudinger and small dispersion approximation following [Christ-C-Tao].

イロト イボト イヨト イヨト 二日

## $H^1(\mathbb{R}^2)$ "Critical" Case

- If H[u<sub>0</sub>] − M[u<sub>0</sub>] ≤ 1 then NLS<sub>exp</sub>(ℝ<sup>2</sup>) is globally well-posed. Uniform continuity of data-to-solution map fails to hold for data satisfying H[u<sub>0</sub>] − M[u<sub>0</sub>] > 1.
   [Work in progress with Ibrahim, Majdoub, Masmoudi]
- Well-posedness result relies upon Strichartz estimates, Moser-Trudinger inequality, and a log-Sobolev inequality. (Largely based on similar result for NLKG by [Ibrahim-Majdoub-Masmoudi])
- Ill-posedness result relies upon optimizing sequence for Moser-Trudinger and small dispersion approximation following [Christ-C-Tao].
- Scattering?

イロト イボト イヨト イヨト 二日

イロト イヨト イヨト イヨト

3

Consider  $NLS_7^+(\mathbb{R}^3)$ . Typical case?

-

(3)

A ■

Consider  $NLS_7^+(\mathbb{R}^3)$ . Typical case?

 Numerical experiments by [Blue-Sulem] and also for corresponding NLKG [Strauss-Vazquez] suggest GWP and scattering.

Consider  $NLS_7^+(\mathbb{R}^3)$ . Typical case?

- Numerical experiments by [Blue-Sulem] and also for corresponding NLKG [Strauss-Vazquez] suggest GWP and scattering.
- ► Conjecture: NLS<sub>7</sub><sup>+</sup>(ℝ<sup>3</sup>) is GWP and scatters in H<sup>7/6</sup>(ℝ<sup>3</sup>). [See discussion by Bourgain, GAFA Special Volume, 2000]

[Work in progress with Raynor, Sulem, Wright....details remain.]

• 3 • 4 3

[Work in progress with Raynor, Sulem, Wright....details remain.]

Question: Qualitative properties mass supercritical NLS blowup?

[Work in progress with Raynor, Sulem, Wright....details remain.]

Question: Qualitative properties mass supercritical NLS blowup? Restrict attention to  $H^{1/2}$ -critical  $NLS_3^-(\mathbb{R}^3)$ .

[Work in progress with Raynor, Sulem, Wright....details remain.]

Question: Qualitative properties mass supercritical NLS blowup? Restrict attention to  $H^{1/2}$ -critical  $NLS_3^-(\mathbb{R}^3)$ .

►  $T^*$  defined via divergence of  $||u||_{L^5_{tx}}$  or  $||D^{1/2}u||_{L^{10/3}_{tx}}$ .

[Work in progress with Raynor, Sulem, Wright....details remain.]

Question: Qualitative properties mass supercritical NLS blowup? Restrict attention to  $H^{1/2}$ -critical  $NLS_3^-(\mathbb{R}^3)$ .

- $T^*$  defined via divergence of  $||u||_{L^5_{tx}}$  or  $||D^{1/2}u||_{L^{10/3}}$ .
- ► Finite energy radial blowups explode at spatial origin.

[Work in progress with Raynor, Sulem, Wright....details remain.]

Question: Qualitative properties mass supercritical NLS blowup? Restrict attention to  $H^{1/2}$ -critical  $NLS_3^-(\mathbb{R}^3)$ .

- $T^*$  defined via divergence of  $||u||_{L^5_{tx}}$  or  $||D^{1/2}u||_{L^{10/3}}$ .
- ► Finite energy radial blowups explode at spatial origin.
- ▶ Heuristics and numerics suggest asymptotic profile Q which decays near spatial infinity like  $|y|^{-1} \implies Q \notin L^3(\mathbb{R}^3)$ . Sobolev embedding  $H^{1/2} \hookrightarrow L^3$  suggests as  $t \uparrow T^*$

$$\|u(t)\|_{H^{1/2}}\sim |\log(T^*-t)|\to\infty.$$

- 4 回 ト 4 ヨ ト 4 ヨ ト

[Work in progress with Raynor, Sulem, Wright....details remain.]

Question: Qualitative properties mass supercritical NLS blowup? Restrict attention to  $H^{1/2}$ -critical  $NLS_3^-(\mathbb{R}^3)$ .

- $T^*$  defined via divergence of  $||u||_{L^5_{tx}}$  or  $||D^{1/2}u||_{L^{10/3}}$ .
- ► Finite energy radial blowups explode at spatial origin.
- ▶ Heuristics and numerics suggest asymptotic profile Q which decays near spatial infinity like  $|y|^{-1} \implies Q \notin L^3(\mathbb{R}^3)$ . Sobolev embedding  $H^{1/2} \hookrightarrow L^3$  suggests as  $t \uparrow T^*$

$$\|u(t)\|_{H^{1/2}} \sim |\log(T^*-t)| \rightarrow \infty.$$

 Frequency heuristic: Bounded H<sup>1/2</sup> blowup inconsistent with mass conservation.

イロン イヨン イヨン

▶ Contradiction Hypothesis (CH): Assume  $\exists \Lambda < \infty$  such that

$$\|u\|_{L^{\infty}_t H^{1/2}_x([0,T^*)\times\mathbb{R}^3)} < \Lambda.$$

• 3 • 4 3

▶ Contradiction Hypothesis (CH): Assume  $\exists \Lambda < \infty$  such that

$$\left\|u\right\|_{L^{\infty}_{t}H^{1/2}_{x}\left([0,T^{*})\times\mathbb{R}^{3}\right)}<\Lambda.$$

▶ Concentration Property: If  $H^1 \cap \{radial\} \ni u_0 \mapsto u$  solves  $NLS_3^-(\mathbb{R}^3), T^* < \infty$  and we assume (CH) then

$$\liminf_{t\uparrow T^*} \|u(t)\|_{L^3_{|x|<(T^*-t)^{1/2-}}} \geq \frac{\sqrt{2}}{\pi^{2/3}} = c^*.$$

向下 イヨト イヨト

▶ Contradiction Hypothesis (CH): Assume  $\exists \Lambda < \infty$  such that

$$\left\|u\right\|_{L^{\infty}_{t}H^{1/2}_{x}\left([0,T^{*})\times\mathbb{R}^{3}\right)}<\Lambda.$$

▶ Concentration Property: If  $H^1 \cap \{radial\} \ni u_0 \mapsto u$  solves  $NLS_3^-(\mathbb{R}^3), T^* < \infty$  and we assume (CH) then

$$\liminf_{t\uparrow T^*} \|u(t)\|_{L^3_{|x|<(T^*-t)^{1/2-}}} \geq \frac{\sqrt{2}}{\pi^{2/3}} = c^*.$$

向下 イヨト イヨト

▶ Contradiction Hypothesis (CH): Assume  $\exists \Lambda < \infty$  such that

$$\|u\|_{L^{\infty}_t H^{1/2}_x([0,T^*)\times\mathbb{R}^3)} < \Lambda.$$

▶ Concentration Property: If  $H^1 \cap \{ radial \} \ni u_0 \longmapsto u$  solves  $NLS_3^-(\mathbb{R}^3), T^* < \infty$  and we assume (CH) then

$$\liminf_{t\uparrow \mathcal{T}^*} \|u(t)\|_{L^3_{|x|<(\mathcal{T}^*-t)^{1/2-}}} \geq \frac{\sqrt{2}}{\pi^{2/3}} = c^*.$$

The proof follows [Merle-Tsutsumi] with the (CH) upper bound as a proxy for  $L^2$  conservation. Explicit constant from sharp Gagliardo-Nirenberg estimate. [Delpino-Dolbeaut]

イロト イボト イヨト イヨト

イロン イヨン イヨン

► Frequency level Sets:

$$R_{\mu}(t) := \sup\{R : \|P_{|\xi| > R}u(t)\|_{\dot{H}^{1/2}_{x}} > \mu\}$$

イロト イヨト イヨト

► Frequency level Sets:

$$R_{\mu}(t) := \sup\{R : \|P_{|\xi| > R}u(t)\|_{\dot{H}^{1/2}_{x}} > \mu\}$$

イロト イヨト イヨト

#### Frequency level Sets:

$$R_{\mu}(t) := \sup\{R : \|P_{|\xi|>R}u(t)\|_{\dot{H}^{1/2}_{x}} > \mu\}$$
  
Concentration  $\implies R_{c^{*}}(t) \ge (T^{*} - t)^{-1/2}.$ 

イロト イヨト イヨト

Frequency level Sets:

$$R_{\mu}(t) := \sup\{R : \|P_{|\xi|>R}u(t)\|_{\dot{H}^{1/2}_{x}} > \mu\}$$
Concentration  $\implies R_{c^{*}}(t) \ge (T^{*} - t)^{-1/2}.$ 

$$M := \sup\{\mu : R_\mu(t) = O(R_{c^*}(t)) \text{ as } t \uparrow T^*\}$$

イロト イヨト イヨト イヨト

#### Frequency level Sets:

$$R_{\mu}(t) := \sup\{R : \|P_{|\xi|>R}u(t)\|_{\dot{H}_{x}^{1/2}} > \mu\}$$

Concentration  $\implies R_{c^*}(t) \ge (T^* - t)^{-1/2}.$ 

$$M := \sup\{\mu : R_\mu(t) = O(R_{c^*}(t)) \text{ as } t \uparrow T^*\}$$

By design  $R_{M+\gamma_0}(t) = o(R_M(t))$  for all  $\gamma_0 > 0$  as  $t \uparrow T^*$ . There exists  $\mu_0 > 0$  such that  $R_M(t) \sim R_{M-\mu_0}(t)$  as  $t \uparrow T^*$ .

・ロト ・同ト ・ヨト ・ヨト

Frequency level Sets:

$$R_{\mu}(t) := \sup\{R : \|P_{|\xi| > R}u(t)\|_{\dot{H}_{x}^{1/2}} > \mu\}$$

Concentration  $\implies R_{c^*}(t) \ge (T^* - t)^{-1/2}.$ 

$$M := \sup\{\mu : R_\mu(t) = O(R_{c^*}(t)) \text{ as } t \uparrow T^*\}$$

By design  $R_{M+\gamma_0}(t) = o(R_M(t))$  for all  $\gamma_0 > 0$  as  $t \uparrow T^*$ . There exists  $\mu_0 > 0$  such that  $R_M(t) \sim R_{M-\mu_0}(t)$  as  $t \uparrow T^*$ .

▶ Fix a number *K* by the condition

$$K^{1/2}\mu_0=3\Lambda.$$

- 4 同 6 4 三 6 4 三 6
イロト イヨト イヨト

▶ Solution Decomposition: At a time  $t_0 < T^*$ , decompose

$$u(t_0) = u^{low}(t_0) + u^{gap}(t_0) + u^{hi}(t_0)$$

3 ×

▶ Solution Decomposition: At a time  $t_0 < T^*$ , decompose

$$u(t_0) = u^{low}(t_0) + u^{gap}(t_0) + u^{hi}(t_0)$$

3 ×

▶ Solution Decomposition: At a time  $t_0 < T^*$ , decompose

$$u(t_0) = u^{low}(t_0) + u^{gap}(t_0) + u^{hi}(t_0)$$

with respect to frequency regions

$$egin{aligned} &\xi| < R_{\mathcal{M}+\gamma_0}(t_0) \ &R_{\mathcal{M}+\gamma_0}(t_0) < |\xi| < R_\mathcal{M}(t_0) \ &R_\mathcal{M}(t_0) < |\xi|. \end{aligned}$$

▶ Solution Decomposition: At a time  $t_0 < T^*$ , decompose

$$u(t_0) = u^{low}(t_0) + u^{gap}(t_0) + u^{hi}(t_0)$$

with respect to frequency regions

$$egin{aligned} &\xi| < R_{M+\gamma_0}(t_0) \ &R_{M+\gamma_0}(t_0) < |\xi| < R_{M}(t_0) \ &R_{M}(t_0) < |\xi|. \end{aligned}$$

Evolve  $u^{l}$  and  $u^{g}$  forward on  $[t_{0}, T^{*})$  using  $NLS_{3}^{-}(\mathbb{R}^{3})$ . Evolve  $u^{h}$  according to  $\widetilde{NLS}$  so that

$$u(t) = u'(t) + u^{g}(t) + u^{h}(t).$$

▶ Solution Decomposition: At a time  $t_0 < T^*$ , decompose

$$u(t_0) = u^{low}(t_0) + u^{gap}(t_0) + u^{hi}(t_0)$$

with respect to frequency regions

$$egin{aligned} &\xi| < R_{\mathcal{M}+\gamma_0}(t_0) \ &R_{\mathcal{M}+\gamma_0}(t_0) < |\xi| < R_{\mathcal{M}}(t_0) \ &R_{\mathcal{M}}(t_0) < |\xi|. \end{aligned}$$

Evolve  $u^l$  and  $u^g$  forward on  $[t_0, T^*)$  using  $NLS_3^-(\mathbb{R}^3)$ . Evolve  $u^h$  according to  $\widetilde{NLS}$  so that

$$u(t) = u'(t) + u^{g}(t) + u^{h}(t).$$

► *K*th Doubling Time after *t*<sub>0</sub>:

$$t_1 := \inf\{t \in (t_0, T^*) : R_M(t_1) > KR_{M-\mu_0}(t_0)\}$$

イロト イヨト イヨト

 High Frequency Mass Freezing Contradiction: Suppose we show the high frequency mass freezing property

$$egin{aligned} &\|P_{|\xi|>R_M(t_0)}u(t_1)\|_{L^2} \geq rac{1}{2}\|P_{|\xi|>R_M}(t_0)u(t_0)\|_{L^2}\ &\gtrsim \mu_0 R_{M-\mu_0}^{-1/2}(t_0). \end{aligned}$$

High Frequency Mass Freezing Contradiction: Suppose we show the high frequency mass freezing property

$$egin{aligned} &\|P_{|\xi|>R_{M}(t_{0})}u(t_{1})\|_{L^{2}}\geqrac{1}{2}\|P_{|\xi|>R_{M}}(t_{0})u(t_{0})\|_{L^{2}}\ &\gtrsim\mu_{0}R_{M-\mu_{0}}^{-1/2}(t_{0}). \end{aligned}$$

For small γ<sub>0</sub>, we can not park this mass inside the gap R<sub>M</sub>(t<sub>0</sub>) < |ξ| < R<sub>M</sub>(t<sub>1</sub>) so we have to put it in the high frequency boondox |ξ| > R<sub>M</sub>(t<sub>1</sub>).

High Frequency Mass Freezing Contradiction: Suppose we show the high frequency mass freezing property

$$egin{aligned} \|P_{|\xi|>R_{M}(t_{0})}u(t_{1})\|_{L^{2}}&\geqrac{1}{2}\|P_{|\xi|>R_{M}}(t_{0})u(t_{0})\|_{L^{2}}\ &\gtrsim\mu_{0}R_{M-\mu_{0}}^{-1/2}(t_{0}). \end{aligned}$$

- For small *γ*<sub>0</sub>, we can not park this mass inside the gap *R<sub>M</sub>(t*<sub>0</sub>) < |ξ| < *R<sub>M</sub>(t*<sub>1</sub>) so we have to put it in the high frequency boondox |ξ| > *R<sub>M</sub>(t*<sub>1</sub>).
- Since, at time  $t_1, R_M(t_1) \ge KR_{M-\mu_0}(t_0)$ , we conclude

$$||u(t_1)||_{H^{1/2}} \ge 3\Lambda,$$

a contradiction.

イロト イヨト イヨト

• Main issue is to control the  $L^2$  mass increment of  $P_{|\xi|>R_M(t_0)}u^h(\cdot)$ under the  $\widetilde{NLS}$  evolution from  $t_0$  to  $t_1$ .

- ► Main issue is to control the  $L^2$  mass increment of  $P_{|\xi|>R_M(t_0)}u^h(\cdot)$ under the  $\widetilde{NLS}$  evolution from  $t_0$  to  $t_1$ .
- ▶ We must control 4-linear spacetime integrals like

$$\int_{t_0}^{t_1} \int_{\mathbb{R}^3} P_{>R_M(t_0)}(u^{l}\overline{u^g}u^h) P_{>R_M(t_0)}\overline{u^h} dx dt.$$

- ▶ Main issue is to control the  $L^2$  mass increment of  $P_{|\xi|>R_M(t_0)}u^h(\cdot)$ under the  $\widetilde{NLS}$  evolution from  $t_0$  to  $t_1$ .
- ▶ We must control 4-linear spacetime integrals like

$$\int_{t_0}^{t_1} \int_{\mathbb{R}^3} P_{>R_M(t_0)}(u^{I}\overline{u^g}u^h) P_{>R_M(t_0)}\overline{u^h} dx dt.$$

Since L<sup>4</sup><sub>tx</sub> is H<sup>1/4</sup>-critical and we have H<sup>1/2</sup> control on u we can control such integrals with some gain:

$$\lesssim \left\{ (t_1 - t_0)^{1/5} \|u\|_{L^5_{t,x}([t_0,t_1] imes \mathbb{R}^3)} 
ight\}^{5/2} \Lambda^{3/2}.$$

- ► Main issue is to control the  $L^2$  mass increment of  $P_{|\xi|>R_M(t_0)}u^h(\cdot)$ under the  $\widetilde{NLS}$  evolution from  $t_0$  to  $t_1$ .
- ▶ We must control 4-linear spacetime integrals like

$$\int_{t_0}^{t_1} \int_{\mathbb{R}^3} P_{>R_M(t_0)}(u'\overline{u^g}u^h) P_{>R_M(t_0)}\overline{u^h} dx dt.$$

Since L<sup>4</sup><sub>tx</sub> is H<sup>1/4</sup>-critical and we have H<sup>1/2</sup> control on u we can control such integrals with some gain:

$$\lesssim \left\{ (t_1 - t_0)^{1/5} \| u \|_{L^5_{t,x}([t_0,t_1] \times \mathbb{R}^3)} \right\}^{5/2} \Lambda^{3/2}.$$

► Assuming that  $||u||_{L^{5}_{tx}([0,t]\times\mathbb{R}^{3})} \lesssim (T^{*}-t)^{-1/5+}$  and the Concentration Property we contradict (CH) proving critical norm explosion.

#### Remarks

-2

▲口 → ▲圖 → ▲ 臣 → ▲ 臣 → □

#### Remarks

## • Spacetime $L_{tx}^5$ upper bound is consistent with heuristics.

-

(3)

< 🗇 🕨

#### REMARKS

- Spacetime  $L_{tx}^5$  upper bound is consistent with heuristics.
- Concentration Property following [Merle-Tsutsumi] proof assumed H<sup>1</sup> ∩ {radial} data. The rest of the argument is at the critical level.

#### REMARKS

- Spacetime  $L_{tx}^5$  upper bound is consistent with heuristics.
- Concentration Property following [Merle-Tsutsumi] proof assumed H<sup>1</sup> ∩ {radial} data. The rest of the argument is at the critical level.
- ► Under (CH) bound, Bourgain's L<sup>2</sup> critical concentration result extends to the NLS<sub>3</sub><sup>-</sup>(ℝ<sup>3</sup>) case to prove L<sup>3</sup> and H<sup>1/2</sup> concentration. [with Roudenko]

This relaxes the  $H^1 \cap \{radial\}$  assumptions to  $H^{1/2}$ .

#### REMARKS

- Spacetime  $L_{tx}^5$  upper bound is consistent with heuristics.
- Concentration Property following [Merle-Tsutsumi] proof assumed H<sup>1</sup> ∩ {radial} data. The rest of the argument is at the critical level.
- Under (CH) bound, Bourgain's L<sup>2</sup> critical concentration result extends to the NLS<sub>3</sub><sup>-</sup>(ℝ<sup>3</sup>) case to prove L<sup>3</sup> and H<sup>1/2</sup> concentration. [with Roudenko]
   This relaxes the H<sup>1</sup> ∩ {radial} assumptions to H<sup>1/2</sup>.
- Extends to the general mass supercritical case?