The *I*-method with a Morawetz BOOTSTRAP

J. Colliander

University of Toronto

Séminaire, Université de Paris-Nord 13, Villetaneuse

1 GENERALIZED VIRIAL IDENTITIES

2 A Priori Spacetime Estimates

3 *I*-Method with Morawetz Bootstrap

1. Generalized Virial Identities

Consider the initial value problem:

$$\begin{cases} (i\partial_t + \Delta)u = \pm F'(|u|^2)u\\ u(0, x) = u_0(x) \end{cases} \qquad (NLS_F^{\pm}(\mathbb{R}^d))$$

Remarks:

• Assume $F' \ge 0$. The + case is defocusing; - is focusing.

Generalized NLS with Lagrangian derivation.

• U(1) solution symmetry: $u \rightarrow e^{i\theta}u$.

TIME INVARIANT QUANTITIES

The following quantities do not change with time:

$$\begin{split} \mathsf{Mass} &= \int_{\mathbb{R}^d} |u(t,x)|^2 dx.\\ \mathsf{Momentum} &= 2\Im \int_{\mathbb{R}^2} \overline{u}(t) \nabla u(t) dx.\\ \mathsf{Energy} &= \mathcal{H}[u(t)] = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla u(t)|^2 dx \pm \mathcal{F}(|u(t)|^2) dx. \end{split}$$

 \implies a priori conservation controls (defocusing case):

$$\|u\|_{L^{\infty}_{t}L^{2}_{x}} \leq \|u_{0}\|_{L^{2}} \|\nabla u\|_{L^{\infty}_{t}L^{2}_{x}} \leq E[u_{0}].$$

These are very useful bounds but do not give any decay in time.

We consider an even more general NLS equation.

• Suppose $\phi : [0, T] \times \mathbb{R}^d \to \mathbb{C}$ solves generalized NLS

$$(i\partial_t + \Delta)\phi = \mathcal{N}$$
 $(GNLS(\mathbb{R}^d))$

for $\mathcal{N} = \mathcal{N}(t, x, \phi) : [0, T] \times \mathbb{R}^d \times \mathbb{C} \to \mathbb{C}$. Assume ϕ is nice.

- Not necessarily Lagrangian; No U(1) symmetry.
- Express mass & momentum (non)conservation for GNLS.

Write $\partial_{x_j}\phi = \partial_j\phi = \phi_j$.

LOCAL MASS/MOMENTUM (NON)CONSERVATION

- mass density: $T_{00} = |\phi|^2$
- momentum density/mass current:

$$T_{0j}=T_{j0}=2\Im(\overline{\phi}\phi_j)$$

- (linear part of the) momentum current: $L_{jk} = L_{kj} = -\partial_j \partial_k |\phi|^2 + 4\Re(\overline{\phi_j}\phi_k)$
- mass bracket: $\{f,g\}_m = \Im(f\overline{g})$
- momentum bracket: $\{f, g\}_{p}^{j} = \Re(f\partial_{j}\overline{g} g\partial_{j}\overline{f})$

Local mass (non)conservation identity:

$$\partial_t T_{00} + \partial_j T_{0j} = 2\{\mathcal{N}, \phi\}_m$$

Local momentum (non)conservation identity:

$$\partial_t T_{0j} + \partial_k L_{kj} = 2\{\mathcal{N}, \phi\}_p^j$$

Consider $\mathcal{N} = F'(|\phi|^2)\phi$ for polynomial $F : \mathbb{R}^+ \to \mathbb{R}$.

We calculate the mass bracket

$$\{F'(|\phi|^2)\phi,\phi\}_m = \Im(F'(|\phi|^2)\phi\overline{\phi}) = 0.$$

Thus mass is conserved for these nonlinearities.

We calculate the momentum bracket

$$\{F'(|\phi|^2)\phi,\phi\}_p^j = -\partial_j G(|\phi|^2)$$

where $G(z) = zF'(z) - F(z) \sim F(z)$.

Thus the momentum bracket contributes a divergence and momentum is conserved for these nonlinearities.

Let $a : \mathbb{R}^d \to \mathbb{R}$ (virial weight). Form the virial potential

$$V_{a}(t) = \int_{\mathbb{R}^d} a(x) |\phi(t,x)|^2 dx.$$

Form the Morawetz action

$$M_{a}(t) = \int_{\mathbb{R}^{d}} \nabla a \cdot 2\Im(\overline{\phi} \nabla \phi) dx.$$

Conservation identities lead to the generalized virial identities

$$\partial_t V_a = M_a + \int_{\mathbb{R}^d} a(x) \{\mathcal{N}, \phi\}_m(t, x) dx,$$

$$\partial_t M_{\boldsymbol{a}} = \int_{\mathbb{R}^d} (-\Delta \Delta \boldsymbol{a}) |\phi|^2 + 4a_{jk} \Re(\overline{\phi_j}\phi_k) + 2a_j \{\mathcal{N}, \phi\}_p^j d\boldsymbol{x}.$$

Remarks on Virial Identities

- The virial potential is a weighted average of the mass density against the virial weight a.
- The Morawetz action is a contraction of the momentum density against ∇a . Vector fields not arising as gradients could also be considered.
- Useful estimates emerge from monotonicity and boundedness of terms in the virial identities.
- Monotone quantities provide dynamical insights.
- Idea of Morawetz Estimates: Cleverly choose the weight function *a* so that $\partial_t M_a \ge 0$ but $M_a \le C(\phi_0)$ to obtain spacetime control on ϕ . This strategy imposes various constraints on *a* which suggest choosing a(x) = |x|.

[Glassey], [Vlasov-Petrischev-Talanov]

- Consider GNLS with $\mathcal{N} = \pm |u|^{4/d} u$. This is the L^2 critical focusing equation $NLS_{1+\frac{d}{2}}^{\pm}(\mathbb{R}^d)$.
- Choose $a(x) = |x|^2$. Calculations reveal that

$$\partial_t^2 \int_{\mathbb{R}^d} |x|^2 |u(t,x)|^2 dx = 16H[u(t)].$$

■ In the focusing case, we can consider initial data *u*₀ with *H*[*u*₀] < 0 and finite variance. Such data must blow up in finite time.

2. A Priori Spacetime Estimates

[LIN-STRAUSS] MORAWETZ IDENTITY

Consider $(i\partial_t + \Delta)\phi = F'(|\phi|^2)\phi$ with $F' \ge 0$ and $x \in \mathbb{R}^3$. Choose a(x) = |x|. Observe that *a* is weakly convex, $\nabla a = \frac{x}{|x|}$ is bounded, and $-\Delta\Delta a = 4\pi\delta_0$. From monotonicity $\partial_t M_a \ge 0$ and the bound $|M_a| \le \sqrt{H[u_0]}$ emerges the Lin-Strauss Morawetz identity

$$M_{a}(T) - M_{a}(0) = \int_{0}^{T} \int_{\mathbb{R}^{3}} 4\pi \delta_{0}(x) |\phi(t,x)|^{2} + (\geq 0) + 4 \frac{G(|\phi|^{2})}{|x|} dx dt.$$

This implies the spacetime control estimate (centered at x = 0)

$$(H[u_0])^{1/2} \|u_0\|_{L^2} \gtrsim \int_0^T \int_{\mathbb{R}^3} \frac{G(|\phi|^2)}{|x|} dx dt.$$

[Morawetz] Reward & Anchor. [Ginibre-Velo] H¹-Scattering.

[BOURGAIN] & [GRILLAKIS] TRUNCATION

Let χ_{B_R} denote a smooth cutoff adapted to B_R = {|x| < R}.
Choose cutoff virial weight a(x) = χ_{B_R}(x)|x|. and calculate

$$M_{a}\Big|_{0}^{T} \geq \int_{0}^{T} \int_{\mathbb{R}^{3}} 4\pi \delta_{0}(x) |\phi(t,x)|^{2} + 4 \int_{0}^{T} \int_{|x| < R/2} \frac{G(|\phi|^{2})}{|x|} dx dt$$

$$|M_{a}|_{0}^{T}| \leq R^{-1}TH[u_{0}] + RH[u_{0}] \implies \text{choose } R \sim T^{1/2} \implies T$$

$$\int_{0}^{\infty} \int_{|x| < T^{1/2}} \frac{G(|\phi|^2)}{|x|} dx \lesssim T^{1/2} \|\nabla u\|_{L^{\infty}_{[0,T]}L^2_x}^2.$$

[Bourgain][Grillakis]: Energy critical bubbles sparse along time axis.

AVERAGING OVER [LIN-STRAUSS] CENTER?

- Translation invariance? Weight $|x|^{-1}$ difficult in proofs.
- Recenter [L-S] at fixed $y \in \mathbb{R}^d$. Set a(x) = |x y|.
- Recentered Morawetz action can be expressed

$$M_{y}[u](t) = \int_{\mathbb{R}^{d}} \frac{(x-y)}{|x-y|} 2\Im(u\nabla\overline{u})(t,x)dx.$$

- Monotonicity $\partial_t M_y[u] \ge 0$: mass is repelled from any $y \in \mathbb{R}^d$.
- Can we average with respect to center y and obtain new translation invariant spacetime control?
- Yes, if we average against the natural density $|u(t, y)|^2$.

INTERACTION MORAWETZ VIA AVERAGING

Define the Morawetz interaction potential

$$M[u](t) = \int_{\mathbb{R}^d_y} |u(t,y)|^2 M_y[u](t) dy.$$

It is bounded: $\left| M[u](t) \right| \lesssim \|u(t)\|_{L^2_x}^3 \| \nabla u(t) \|_{L^2_x}$. We calculate

$$\partial_t M[u] = \int_{\mathbb{R}^d_y} |u(t,y)|^2 \{\partial_t M_y[u]\} + \{\partial_t |u(y)|^2\} M_y[u] dy.$$

• Local conservation & [L-S] \implies monotonicity: $\exists I, II, III, IV$ such that $I, III \ge 0$ and $II + IV \ge 0$ and $\partial_t M[u] = I + II + III + IV$. Integrating in time gives

$$\int_0^T \int_{\mathbb{R}^3} |u(t,x)|^4 dx dt \lesssim \|u(t)\|_{L^{\infty}_T L^2_x}^3 \|\nabla u(t)\|_{L^{\infty}_T L^2_x} \, .$$

2-particle interaction Morawetz

(Hassell 04)

Suppose ϕ_1, ϕ_2 are two solutions of $(i\partial_t + \Delta)\phi = F'(|\phi|^2)\phi$ with $F' \ge 0$ and $x \in \mathbb{R}^3$. The "2-particle" wave function

$$\Psi(t, x_1, x_2) = \phi_1(t, x_1)\phi_2(t, x_2)$$

satisfies an NLS-type equation on \mathbb{R}^{1+6}

$$(i\partial_t + \Delta_1 + \Delta_2)\Psi = [F'(|\phi_1|^2) + F'(|\phi_2|^2)]\Psi.$$

- Note that $[F'(|\phi_1|^2) + F'(|\phi_2|^2)] \ge 0$ so defocusing.
- Reparametrize ℝ⁶ using center-of-mass coordinates (x̄, y) with x̄ = ½(x₁ + x₂) ∈ ℝ³. Note that y = 0 corresponds to the diagonal x₁ = x₂ = x̄. Apply the generalized virial identity with the **choice** a(x₁, x₂) = |y|. Dismissing terms with favorable signs, one obtains...

2-particle interaction Morawetz

$$\begin{split} \|\nabla u\|_{L^{\infty}_{[0,T]}L^{2}_{x}}\|u_{0}\|^{3}_{L^{2}} &\geq \int_{0}^{T}\int_{\mathbb{R}^{6}}(-\Delta_{6}\Delta_{6}|y|)|\Psi(x_{1},x_{2})|^{2}dx_{1}dx_{2}dt\\ &\geq c\int_{0}^{T}\int_{\mathbb{R}^{6}}\delta_{\{y=0\}}(x_{1},x_{2})|\phi_{1}(x_{1})\phi_{2}(x_{2})|^{2}dx_{1}dx_{2}dt\\ &\geq c\int_{0}^{T}\int_{\mathbb{R}^{3}}|\phi_{1}(t,\overline{x})\phi_{2}(t,\overline{x})|^{2}d\overline{x}dt. \end{split}$$

Specializing to $\phi_1 = \phi_2$ gives the **2-particle Morawetz estimate**

$$\int_0^T \int_{\mathbb{R}^3} |\phi(t,x)|^4 dx dt \le C \|\nabla u\|_{L^{\infty}_{[0,T]}L^2_x} \|u_0\|^3_{L^2_x}$$

valid uniformly for all defocusing NLS equations on \mathbb{R}^3 .

"The" 2-particle Morawetz Estimate

Efforts to extend the $L^4(\mathbb{R}_t \times \mathbb{R}^3_x)$ interaction Morawetz to the \mathbb{R}^2_x setting led to...

THEOREM (C-GRILLAKIS-TZIRAKIS & PLANCHON-VEGA)

Finite energy solutions of any defocusing $NLS^+(\mathbb{R}^d)$ satisfy

$$\|D^{\frac{3-d}{2}}|u|^2\|_{L^2_{t,x}}^2 \lesssim \|u_0\|_{L^2_x}^3 \|\nabla u\|_{L^{\infty}_t L^2_x}.$$

- Independently & simultaneously by [Planchon-Vega].
- Gives simple proof of H¹-scattering in mass supercritical case. [Nakanishi]
- Simplified proof extends to H^s for certain s < 1.
- May play a role in resolving the L² scattering conjecture?

4-particle Morawetz Estimate

(Hassel-Tao) [C-Holmer-Visan-Zhang]

$$\mathbb{R}^{4} = \{ \mathbf{x} = (x_{1}, x_{2}, x_{3}, x_{4}) : x_{i} \in \mathbb{R}; i = 1, 2, 3, 4 \}$$

$$\overline{x} = \text{center of mass} = \frac{1}{4}(x_{1} + x_{2} + x_{3} + x_{4}).$$
Define $y = (x_{1} - \overline{x}, x_{2} - \overline{x}, x_{3} - \overline{x}, x_{4} - \overline{x}).$

$$\mathbb{R}^{4} \ni \mathbf{x} = (x_{1}, x_{2}, x_{3}, x_{4}) \iff (\overline{x}, y) \in \mathbb{R} \times \mathbb{R}^{3}$$

The 4-particle wave function

$$\Psi(t,\mathbf{x}) = \prod_{i=1}^{4} \phi_1(t,x_i)$$

satisfies a defocusing NLS equation on \mathbb{R}^{1+4} .

• Choice of virial weight $a(\mathbf{x}) = |y|$ spawns

$$\int_0^T \int_{\mathbb{R}} |u|^8 d\overline{x} dt \lesssim \|u\|_{L^\infty_T L^2_x}^7 \|\nabla u\|_{L^\infty_T L^2_x}$$

How does this estimate generalize to other dimensions?

Application: **Subcritical scattering** for certain $NLS_p^+(\mathbb{R}^d)$. [CKSTT], [CHVZ], [Fang-Grillakis], [C-Grillakis-Tzirakis]

- 2-particle Morawetz is an $\dot{H}^{1/4}$ -critical input. 4-particle Morawetz is an $\dot{H}^{1/8}$ -critical input.
- Scaling invariant H^s for $NLS_p^+(\mathbb{R}^d): s_c = \frac{d}{2} \frac{2}{p-1}$
- When $1/4 < s_c < 1 \exists s_* \in (s_c, 1)$ and $\forall s \ge s_*$ the H^s -solutions of $NLS_p^+(\mathbb{R}^d)$ scatter.
- We obtain scattering for certain energy subcritcal $(s_c < 1)$ NLS for infinite energy data of subcritical regularity $(s_c < s_*)$.
- The critical scattering conjecture corresponds to $s_c = s_*$. This is known (for general data) only for $s_c = 1$.

I-METHOD WITH MORAWETZ BOOTSTRAP

Consider
$$NLS^+_{2k+1}(\mathbb{R})$$
 (nonlinearity $+|u|^{2k}u$) for $k = 3, 4, ...$
Note that $s_c = \frac{1}{2} - \frac{1}{k}$.
Define
 $8k - 16$

$$s_k=\frac{8k-10}{9k-14}<1.$$

THEOREM (C-HOLMER-VISAN-ZHANG)

 $\forall s > s_k, H^s(\mathbb{R}) \ni u_0 \longmapsto u \text{ solving } NLS^+_{2k+1}(\mathbb{R}) \text{ is global in time and scatters: } \exists u_{\pm} \in H^s(\mathbb{R}) \text{ such that}$

$$\lim_{t\to\pm\infty}\|u(t)-e^{it\Delta}u_{\pm}\|_{H^s(\mathbb{R})}=0.$$

Proof treats a family of equations; Wish that $s_k \rightarrow 1/2$ as $k \rightarrow \infty$.

The 4-particle $L^8(\mathbb{R}_t \times \mathbb{R}_x)$ estimate may be reexpressed:

$$\int_0^T \int_{\mathbb{R}} |u|^8 d\overline{x} dt \lesssim \|u\|_{L^\infty_T L^2_x}^6 \|u\|_{L^\infty_T \dot{H}^{1/2}_x},$$

 $\implies \|u\|_{L^3_{t,x}} \lesssim \|u\|_{L^\infty_t H^1_x} \implies H^1\text{-scattering using some interpolation.}$

 Recall that, based on a Hölder-in-time step, subcritical local-in-time theory gives

$$T_{Iwp} \sim \|u_0\|_{H^s}^{-\frac{2}{s-s_c}}.$$

Bootstrap toward scattering: Hölder-in-time is forbidden.

Suppose we have almost conservation of modified energy.

- \blacksquare RHS of 4-particle \lesssim almost conserved modified energy
- **2** \implies $L^8_{t,x}$ controlled on long time interval $t \in [0, T]$
- **3** \implies spacetime slab decomposition: $[0, T] \times \mathbb{R} = \cup_{j=1}^{J} [t_j, t_{j+1}) \times \mathbb{R}$ such that

$$\|u\|_{L^{\otimes}([t_{j},t_{j+1})\times\mathbb{R})} = \eta \sim \frac{1}{100}$$

4 \implies almost conserved modified energy on $[t_j, t_{j+1}]$ 5 \implies RHS of 4-particle.... bootstrap loop! LEMMA (ALMOST CONSERVATION ON SLAB)

Let $H^s \ni u_0 \to u$ solve $NLS^+_{2k+1}(\mathbb{R}^d)$ with $s > s_k$. Suppose we have a spacetime slab $[t_-, t_+]$ on which

 $\|u\|_{L^8([t_-,t_+]\times\mathbb{R})} \lesssim \eta$

and $\exists t_0 \in [t_-, t_+]$ such that $H[I_N u(t_0)] \leq 1$. Then for large N we have almost conservation:

$$\sup_{t\in[t_-,t_+]}H[I_Nu(t)]=H[I_Nu(t_0)]+O(N^{-1+}).$$

Rescaling and continuity arguments glue it all together.