# Nonlinear Schrödinger Evolutions from Low Regularity Initial Data

J. Colliander

University of Toronto

Warwick, June 2009

### **1** CUBIC NLS ON $\mathbb{R}^2$

- **2** High-Low Fourier Truncation
- **3** BILINEAR STRICHARTZ ESTIMATE
- 4 The *I*-Method of Almost Conservation

# 1. Cubic NLS Initial Value Problem on $\mathbb{R}^2$

# 1. CUBIC NLS INITIAL VALUE PROBLEM ON $\mathbb{R}^2$

We consider the initial value problems:  $\begin{cases}
(i\partial_t + \Delta)u = \pm |u|^2 u \\
u(0, x) = u_0(x).
\end{cases}$ (NLS<sup>±</sup><sub>3</sub>(R<sup>2</sup>))

The + case is called defocusing; – is focusing.  $NLS_3^{\pm}$  is ubiquitous in physics. The solution has a dilation symmetry

$$u^{\lambda}(\tau, y) = \lambda^{-1} u(\lambda^{-2}\tau, \lambda^{-1}y).$$

which is invariant in  $L^2(\mathbb{R}^2)$ . This problem is  $L^2$ -critical.

(This talk mostly addresses the defocusing case.)

# TIME INVARIANT QUANTITIES

$$\begin{split} \mathsf{Mass} &= \int_{\mathbb{R}^d} |u(t,x)|^2 dx.\\ \mathsf{Momentum} &= 2\Im \int_{\mathbb{R}^2} \overline{u}(t) \nabla u(t) dx.\\ \mathsf{Energy} &= H[u(t)] = \frac{1}{2} \int_{R^2} |\nabla u(t)|^2 dx \pm \frac{1}{2} |u(t)|^4 dx.\\ \mathsf{Hamiltonian} & \mathsf{kinetic} & \mathsf{potential} \end{split}$$

- Mass is  $L^2$ ; Momentum is close to  $H^{1/2}$ ; Energy involves  $H^1$ .
- Dynamics on a sphere in L<sup>2</sup>; focusing/defocusing energy.
- Local conservation laws express **how** quantity is conserved: e.g.,  $\partial_t |u|^2 = \nabla \cdot 2\Im(\overline{u}\nabla u)$ . Frequency Localizations?

The solution of the linear Schrödinger initial value problem

$$\begin{cases} (i\partial_t + \Delta)u = 0\\ u(0, x) = u_0(x). \end{cases}$$
 (LS( $\mathbb{R}^d$ ))

is denoted  $u(t,x) = e^{it\Delta}u_0$ . The solution can be given explicitly Fourier Multiplier Representation:

$$e^{it\Delta}u_0(x)=c_{\pi}\int_{\mathbb{R}^d}e^{ix\cdot\xi}e^{-it|\xi|^2}\widehat{u_0}(\xi)d\xi.$$

Convolution Representation:

$$e^{it\Delta}u_0(x)=c_{\pi}^1rac{1}{(it)^{d/2}}\int_{\mathbb{R}^d}e^{irac{|x-y|^2}{4t}}u_0(y)dy.$$

The solution of the linear Schrödinger initial value problem

$$\begin{cases} (i\partial_t + \Delta)u = 0\\ u(0, x) = u_0(x). \end{cases}$$
 (LS( $\mathbb{R}^d$ ))

is denoted  $u(t,x) = e^{it\Delta}u_0$ . The solution can be given explicitly **Fourier Multiplier Representation**:

$$e^{it\Delta}u_0(x)=c_{\pi}\int_{\mathbb{R}^d}e^{ix\cdot\xi}\underline{e^{-it|\xi|^2}}\widehat{u_0}(\xi)d\xi.$$

Convolution Representation:

Modulus 1 Multiplier

$$e^{it\Delta}u_0(x) = c_{\pi}^1 \frac{1}{(it)^{d/2}} \int_{\mathbb{R}^d} e^{i \frac{|x-y|^2}{4t}} u_0(y) dy.$$

# LINEAR SCHRÖDINGER PROPAGATOR

The solution of the linear Schrödinger initial value problem

$$\begin{cases} (i\partial_t + \Delta)u = 0\\ u(0, x) = u_0(x). \end{cases}$$
 (LS( $\mathbb{R}^d$ ))

is denoted  $u(t,x) = e^{it\Delta}u_0$ . The solution can be given explicitly Fourier Multiplier Representation:

$$e^{it\Delta}u_0(x)=c_{\pi}\int_{\mathbb{R}^d}e^{ix\cdot\xi}e^{-it|\xi|^2}\widehat{u_0}(\xi)d\xi.$$

Convolution Representation:

$$e^{it\Delta}u_0(x) = c_{\pi}^1 \frac{1}{(it)^{d/2}} \int_{\mathbb{R}^d} e^{i\frac{|x-y|^2}{4t}}u_0(y)dy$$
  
Time Decay

# Estimates for Linear Schrödinger Propagator

• Fourier Multiplier Representation  $\implies$  Unitary in  $H^s$ :

$$\|D_x^s e^{it\Delta} u_0\|_{L^2_x} = \|D_x^s u_0\|_{L^2_x}.$$

• Convolution Representation  $\implies$  Dispersive estimate:

$$\|e^{it\Delta}u_0\|_{L^{\infty}_x} \leq \frac{C}{t^{d/2}}\|u_0\|_{L^1_x}.$$

Spacetime estimates? Strichartz estimates hold, for example,

$$\|e^{it\Delta}u_0\|_{L^4(\mathbb{R}_t\times\mathbb{R}^2_x)}\leq C\|u_0\|_{L^2(\mathbb{R}^2_x)}$$

(Strichartz estimates linked to Fourier restriction phenomena.)

# Local-in-time theory for $NLS_3^{\pm}(\mathbb{R}^2)$

■ 
$$\forall u_0 \in L^2(\mathbb{R}^2) \exists T_{lwp}(u_0)$$
 determined by  
 $\|e^{it\Delta}u_0\|_{L^4_{tx}([0,T_{lwp}]\times\mathbb{R}^2)} < \frac{1}{100}$  such that  
 $\exists \text{ unique } u \in C([0, T_{lwp}]; L^2) \cap L^4_{tx}([0, T_{lwp}] \times \mathbb{R}^2)$  solving  
 $NLS^+_3(\mathbb{R}^2).$   
■  $\forall u_0 \in H^s(\mathbb{R}^2), s > 0, T_{lwp} \sim \|u_0\|_{H^s}^{-\frac{2}{s}}$  and regularity persists:  
 $u \in C([0, T_{lwp}]; H^s(\mathbb{R}^2)).$ 

• Define the maximal forward existence time  $T^*(u_0)$  by

$$\|u\|_{L^4_{tx}([0,T^*-\delta]\times\mathbb{R}^2)}<\infty$$

for all  $\delta > 0$  but diverges to  $\infty$  as  $\delta \searrow 0$ .

**a**  $\exists$  small data scattering threshold  $\mu_0 > 0$ 

$$||u_0||_{L^2} < \mu_0 \implies ||u||_{L^4_{tx}(\mathbb{R}\times\mathbb{R}^2)} < 2\mu_0.$$

What is the ultimate fate of the local-in-time solutions?

 $\frac{L^2\text{-critical Scattering Conjecture:}}{L^2 \ni u_0 \longmapsto u \text{ solving } NLS_3^+(\mathbb{R}^2) \text{ is global-in-time and}} \|u\|_{L^4_{t,x}} < A(u_0) < \infty.$ 

Moreover,  $\exists \ u_{\pm} \in L^2(\mathbb{R}^2)$  such that

$$\lim_{t\to\pm\infty}\|e^{\pm it\Delta}u_{\pm}-u(t)\|_{L^2(\mathbb{R}^2)}=0.$$

Same statement for focusing  $NLS_3^-(\mathbb{R}^2)$  if  $||u_0||_{L^2} < ||Q||_{L^2}$ . Remarks:

- Known for small data  $||u_0||_{L^2(\mathbb{R}^2)} < \mu_0$ .
- Known for large radial data [Killip-Tao-Visan 07].

# $NLS_3^{\pm}(\mathbb{R}^2)$ : Present Status for General Data

 regularity	idea	reference	]
$s > \frac{2}{3}$	high/low frequency decomposition	[Bourgain98]	
$s > \frac{4}{7}$	H(lu)	[CKSTT02]	
$s>rac{1}{2}$	resonant cut of 2nd energy	[CKSTT07]	
$s \geq \frac{1}{2}$	H(Iu) & Interaction Morawetz	[Fang-Grillakis05]	
$s > \frac{2}{5}$	<i>H</i> ( <i>Iu</i> ) & Interaction <i>I</i> -Morawetz	[CGTz07]	
$s>rac{1}{3}$	resonant cut & <i>I</i> -Morawetz	[C-Roy08]	
<i>s</i> > 0?			

- Morawetz-based arguments are only for defocusing case.
- Focusing results assume  $||u_0||_{L^2} < ||Q||_{L^2}$ .
- Unify theory of focusing-under-ground-state and defocusing?

# 2. BOURGAIN'S HIGH-LOW FOURIER TRUNCATION

# 2. BOURGAIN'S HIGH-LOW FOURIER TRUNCATION

IMRN International Mathematics Research Notices 1998, No. 5

#### Refinements of Strichartz' Inequality and Applications to 2D-NLS with Critical Nonlinearity

J. Bourgain

#### Summary

Consider the 2D IVP

$$\begin{split} & \mathrm{i} u_t + \Delta u + \lambda |u|^2 u = 0 \\ & u(0) = \phi \in L^2(\mathbb{R}^2). \end{split}$$

The theory on the Cauchy problem asserts a unique maximal solution

 $\mathfrak{u} \in \mathcal{C}(\mathbb{I} - \mathbb{T} \mathbb{T}^*[\mathbb{I}^2(\mathbb{R}^2)) \cap \mathbb{I}^4(\mathbb{I} - \mathbb{T} \mathbb{T}^*[\mathbb{I}^4(\mathbb{R}^2))]$ 

Consider the Cauchy problem for defocusing cubic NLS on  $\mathbb{R}^2$ :

$$\begin{cases} (i\partial_t + \Delta)u = +|u|^2 u\\ u(0, x) = \phi_0(x). \end{cases}$$
 (NLS<sub>3</sub><sup>+</sup>( $\mathbb{R}^2$ ))

We describe the first result to give global well-posedness below  $H^1$ .

- $NLS_3^+(\mathbb{R}^2)$  is GWP in  $H^s$  for  $s > \frac{2}{3}$  [Bourgain 98].
- First use of Bilinear Strichartz estimate was in this proof.
- Proof cuts solution into low and high frequency parts.
- For  $u_0 \in H^s$ ,  $s > \frac{2}{3}$ , Proof gives (and crucially exploits),

$$u(t) - e^{it\Delta}\phi_0 \in H^1(\mathbb{R}^2_x).$$

## Setting up; Decomposing Data

- Fix a large target time *T*.
- Let N = N(T) be large to be determined.
- Decompose the initial data:

$$\phi_0 = \phi_{low} + \phi_{high}$$

where

$$\phi_{low}(x) = \int_{|\xi| < N} e^{ix \cdot \xi} \widehat{\phi_0}(\xi) d\xi.$$

Our plan is to evolve:

$$\phi_0 = \phi_{low} + \phi_{high}$$

$$u(t) = u_{low}(t) + u_{high}(t).$$

# Setting up; Decomposing Data

Low Frequency Data Size:

Kinetic Energy:

$$\begin{split} \|\nabla\phi_{low}\|_{L^{2}}^{2} &= \int_{|\xi| < N} |\xi|^{2} |\widehat{\phi_{0}}(\xi)|^{2} dx \\ &= \int_{|\xi| < N} |\xi|^{2(1-s)} |\xi|^{2s} |\widehat{\phi_{0}}(\xi)|^{2} dx \\ &\stackrel{\checkmark}{\leq} N^{2(1-s)} \|\phi_{0}\|_{H^{s}}^{2} \leq C_{0} N^{2(1-s)}. \end{split}$$

■ Potential Energy:  $\|\phi_{low}\|_{L^4_x} \leq \|\phi_{low}\|_{L^2}^{1/2} \|\nabla\phi_{low}\|_{L^2}^{1/2}$  $\implies H[\phi_{low}] \leq CN^{2(1-s)}.$ 

High Frequency Data Size:

$$\|\phi_{high}\|_{L^2} \le C_0 N^{-s}, \ \|\phi_{high}\|_{H^s} \le C_0.$$

The NLS Cauchy Problem for the low frequency data

$$\begin{cases} (i\partial_t + \Delta)u_{low} = +|u_{low}|^2 u_{low} \\ u_{low}(0, x) = \phi_{low}(x) \end{cases}$$

is well-posed on  $[0, T_{lwp}]$  with  $T_{lwp} \sim \|\phi_{low}\|_{H^1}^{-2} \sim N^{-2(1-s)}$ .

We obtain, as a consequence of the local theory, that

$$\|u_{low}\|_{L^4_{[0,T_{lwp}],x}} \leq \frac{1}{100}.$$

The NLS Cauchy Problem for the low frequency data

$$\begin{cases} (i\partial_t + \Delta)u_{high} = +2|u_{low}|^2 u_{high} + \text{similar} + |u_{high}|^2 u_{high} \\ u_{high}(0, x) = \phi_{high}(x) \end{cases}$$

is also well-posed on  $[0, T_{lwp}]$ .

**Remark:** The LWP lifetime of *NLS* evolution of  $u_{low}$  AND the LWP lifetime of the *DE* evolution of  $u_{high}$  are controlled by  $||u_{low}(0)||_{H^1}$ .

The high frequency evolution may be written

$$u_{high}(t) = e^{it\Delta}u_{high} + w.$$

The local theory gives  $||w(t)||_{L^2} \leq N^{-s}$ . Moreover, due to smoothing (obtained via bilinear Strichartz), we have that

$$w \in H^1, \ \|w(t)\|_{H^1} \lesssim N^{1-2s+}.$$
 (SMOOTH!)

Let's postpone the proof of (SMOOTH!).

•  $\forall t \in [0, T_{lwp}]$ , we have

$$u(t) = u_{low}(t) + e^{it\Delta}\phi_{high} + w(t).$$

• At time  $T_{lwp}$ , we define data for the progressive sheme:

$$u(T_{lwp}) = u_{low}(T_{lwp}) + w(T_{lwp}) + e^{iT_{lwp}\Delta}\phi_{high}.$$

$$u(t) = u_{low}^{(2)}(t) + u_{high}^{(2)}(t)$$

for  $t > T_{lwp}$ .

# HAMILTONIAN INCREMENT: $\phi_{low}(0) \longmapsto u_{low}^{(2)}(T_{lwp})$

The Hamiltonian increment due to  $w(T_{lwp})$  being added to low frequency evolution can be calcluated. Indeed, by Taylor expansion, using the bound (SMOOTH!) and energy conservation of  $u_{low}$  evolution, we have using

$$H[u_{low}^{(2)}(T_{lwp})] = H[u_{low}(0)] + (H[u_{low}(T_{lwp}) + w(T_{lwp})] - H[u_{low}(T_{lwp})])$$
  
  $\sim N^{2(1-s)} + N^{2-3s+} \sim N^{2(1-s)}.$ 

Moreover, we can accumulate  $N^s$  increments of size  $N^{2-3s+}$  before we double the size  $N^{2(1-s)}$  of the Hamiltonian. During the iteration, Hamiltonian of "low frequency" pieces remains of size  $\lesssim N^{2(1-s)}$  so the LWP steps are of uniform size  $N^{-2(1-s)}$ . We advance the solution on a time interval of size:

$$N^{s}N^{-2(1-s)} = N^{-2+3s}$$

For  $s > \frac{2}{3}$ , we can choose N to go past target time T.

# How do we prove (SMOOTH!)?

Bourgain's Bilinear Strichartz Estimate: For (dyadic)  $N \leq L$ 

$$\|e^{it\Delta}f_Le^{it\Delta}g_N\|_{L^2_{t,x}} \leq \frac{N^{\frac{2-1}{2}}}{L^{\frac{1}{2}}}\|f_L\|_{L^2_x}\|g_N\|_{L^2_x}.$$

# COROLLARY For $s \ge \frac{1}{2}$ $\|D_x^s(u_1u_2)\|_{L^2_{[0,\delta],x}} \le C(\|u_1\|_{X^{s,1/2+}_{[0,\delta]}}\|u_2\|_{X^{0,1/2+}_{[0,\delta]}} + \|u_1\|_{X^{1/2,1/2+}_{[0,\delta]}}\|u_2\|_{X^{s-1/2,1/2+}_{[0,\delta]}}).$

Thus, the Bilinear Estimate allows us move half a derivative off the high frequency part and instead onto of the low frequency part.

#### Treatment of a typical term in w

 Using the controls we have on u<sub>low</sub>, u<sub>high</sub> from the local theory on [0, T<sub>lwp</sub>], we want to prove for

$$w = \int_0^t e^{i(t-t')\Delta} |u_{low}|^2 u_{high}(t') dt'$$

that 
$$\sup_{t \in [0, T_{lwp}]} \|\nabla w\|_{L^2} < N^{1-2S+}$$
.

By Sobolev embedding, we have

$$\|w\|_{L^{\infty}_{[0,T_{lwp}]}H^{1}} \leq \|w\|_{X^{1,1/2+}_{[0,T_{lwp}]}}$$

• The mapping  $f \mapsto \int_0^t e^{i(t-t')\Delta}$  is formally  $f \mapsto (i\partial_t + \Delta)^{-1}f$  which, due to time localization, is essentially  $\widehat{f} \mapsto \langle \tau + |\xi|^2 \rangle \widehat{f}$ . It suffices to control  $\|D_x|u_{low}|^2 u_{high}\|_{X^{0,-1/2+}}$ . Proceed by duality....

$$\begin{split} \|w\|_{L^{\infty}_{[0,T_{lwp}]}H^{1}} &\leq \sup_{\|g\|_{X^{0,1/2-}} \leq 1} \langle g, D_{x}(|u_{low}|^{2}u_{high}) \rangle. \\ &\lesssim \sup_{g} \langle gD_{x}u_{low}, u_{low}u_{high} \rangle + \sup_{g} \langle gu_{low}, D_{x}(u_{low}u_{high}) \rangle \\ &= \text{easier} + \sup_{g} \langle D_{x}^{1/2}(gu_{low}), D_{x}^{1/2}(u_{low}u_{high}) \rangle. \end{split}$$

The corollary and the available bounds then give (SMOOTH!).

# 3. Bourgain's Bilinear Strichartz Estimate

# 3. BOURGAIN'S BILINEAR STRICHARTZ ESTIMATE

• Recall the Strichartz estimate for  $(i\partial_t + \Delta)$  on  $\mathbb{R}^2$ :

$$\|e^{it\Delta}u_0\|_{L^4(\mathbb{R}_t\times\mathbb{R}^2_x)}\leq C\|u_0\|_{L^2(\mathbb{R}^2_x)}.$$

• We can view this trivially as a bilinear estimate by writing $\|e^{it\Delta}u_0\ e^{it\Delta}v_0\|_{L^2(\mathbb{R}_t\times\mathbb{R}^2_x)} \leq C\|u_0\|_{L^2(\mathbb{R}^2_x)}\|v_0\|_{L^2(\mathbb{R}^2_x)}.$ 

 Bourgain refined this trivial bilinear estimate for functions having certain Fourier support properties.

# BOURGAIN'S BILINEAR STRICHARTZ ESTIMATE



# 3. BOURGAIN'S PROOF

# B98:IMRN

Proof. Since the standard Strichartz inequality yields (112) without the

$$\left(\frac{M_1}{M_2}\right)^{\frac{1}{2}}$$
-factor,

we may assume  $M_2 \gg M_1$ .

Writing

$$(e^{it\Delta}\psi_1)(e^{it\Delta}\psi_2) = \int \widehat{\psi}_1(\xi_1)\widehat{\psi}_2(\xi_2)e^{i[(\xi_1+\xi_2),x+(|\xi_1|^2+|\xi_2|^2)t]} \,d\xi_1 d\xi_2,$$

it follows from Parseval's identity and Cauchy-Schwarz that

$$\begin{split} \|(e^{it\Delta}\psi_1)(e^{it\Delta}\psi_2)\|_2^2 &= \int d\xi d\lambda \left| \int \widehat{\psi}_1(\xi_1) \widehat{\psi}_2(\xi - \xi_1) \delta_0(|\xi_1|^2 + |\xi - \xi_1|^2 - \lambda) \, d\xi_1 \right|^2 \\ &\leq \|\psi_1\|_2^2 \|\psi_2\|_2^2 \left[ \sup_{\lambda, |\xi| \sim M_2} mes_{(1)}[\xi_1| \, |\xi_1| \sim M_1 \right] \\ & \text{ and } |\xi_1|^2 + |\xi - \xi_1|^2 = \lambda ] \\ & < C \frac{M_1}{M_2}. \end{split}$$

# PROOF BASED ON CHANGE OF VARIABLES

Ideas from (Kenig-Ponce-Vega); see [C-Delort-Kenig-Staffilani].

Recall the Fourier multiplier representation of the propagator:

$$e^{it\Delta}f(x) = c_{\pi} \int_{\mathbb{R}^{2}} e^{ix \cdot \xi} e^{-it|\xi|^{2}} \widehat{f}(\xi) d\xi$$
$$= c_{\pi} \int_{\mathbb{R}^{1+2}} e^{i(x \cdot \xi + t\tau)} \delta_{0}(\tau + |\xi|^{2}) \widehat{f}(\xi) d\tau d\xi.$$
Spacetime inverse  
Fourier transform  
With  $f = f_{L}$  and  $g = g_{N}$ , we wish to estimate

$$\|e^{it\Delta}f e^{it\Delta}g\|_{L^2_{t,x}} = \|\mathcal{F}[e^{it\Delta}f e^{it\Delta}g]\|_{L^2_{\tau,\xi}}.$$

Using Fourier tranform property,  $\mathcal{F}(ab) = \widehat{a} * \widehat{b}$ , we find....

# FOURIER MANIPULATIONS; DIRAC EVALUATIONS

We wish to estimate (in  $L^2_{\tau,\xi}$ ) the expression

$$\int_{\substack{\tau = \tau_1 + \tau_2 \\ \xi = \xi_1 + \xi_2}} \delta_0(\tau_1 + |\xi_1|^2) \widehat{f}(\xi_1) \delta_0(\tau_2 + |\xi_2|^2) \widehat{g}(\xi_2).$$

Evaluating the  $\delta$  functions, we find  $\tau_j = -|\xi_j|^2$ , so

$$\int_{\substack{\tau = -|\xi_1|^2 - |\xi_2|^2\\\xi = \xi_1 + \xi_2}} \widehat{f}(\xi_1) \widehat{g}(\xi_2)$$

We proceed by duality. Let's test this against  $d(\tau, \xi)$ ....

$$\|e^{it\Delta}f \ e^{it\Delta}g\|_{L^{2}_{t,x}} = \sup_{\|d\|_{L^{2}_{\tau,\xi} \le 1}} \left\langle d(\tau,\xi) \ , \int_{\substack{\xi = -|\xi_{1}|^{2} - |\xi_{2}|^{2} \\ \xi = \xi_{1} + \xi_{2}}} \widehat{f}(\xi_{1})\widehat{g}(\xi_{2})\right\rangle.$$

$$= \sup_{d} \int d(-|\xi_1|^2 - |\xi_2|^2, \xi_1 + \xi_2) \ \widehat{f}(\xi_1) \widehat{g}(\xi_2) d\xi_1 d\xi_2.$$

The preceding Fourier manipulations have reduced matters to bounding a certain integral. Our task is to show the integral above is bounded by

$$\lesssim \sqrt{rac{N}{L}} \|f\|_{L^2} \|g\|_{L^2} \|d\|_{L^2}.$$

Let's define a change of variables motivated by the arguments of d:

$$u = -|\xi_1|^2 - |\xi_2|^2, \ v = \xi_1 + \xi_2.$$

- Note that  $u \in \mathbb{R}$  and  $v \in \mathbb{R}^2$ . Thus, *dudv* is a measure in 3d while  $d\xi_1 d\xi_2$  is a measure in 4d.
- Note also that ξ<sub>2</sub> is the argument of g = g<sub>N</sub> so it is localized to the smaller dyadic shell |ξ<sub>2</sub>| ~ N ≪ L.
- Let's denote the components of  $\xi_i \in \mathbb{R}^2$  with superscripts:

$$\xi_j = (\xi_j^1, \xi_j^2).$$

The full change of variables is the defined via

$$dudv \ d\xi_2^1 = |J| \ d\xi_1^1 d\xi_1^2 d\xi_2^2 \ d\xi_2^1.$$

We have an *extra* variable *outside* the changed integral.

The Jacobian matrix J is calculated as

$$J = \begin{bmatrix} \frac{\partial u}{\partial \xi_1^1} & \frac{\partial v^1}{\partial \xi_1^1} & \frac{\partial v^2}{\partial \xi_1^1} \\ \frac{\partial u}{\partial \xi_2^1} & \frac{\partial v^1}{\partial \xi_2^1} & \frac{\partial v^2}{\partial \xi_2^1} \\ \frac{\partial u}{\partial \xi_2^2} & \frac{\partial v^1}{\partial \xi_2^2} & \frac{\partial v^2}{\partial \xi_2^2} \end{bmatrix}.$$

The explicit forms for u, v permit calculating

$$|J| = 2|\xi_1^1 - \xi_1^2|.$$

Since  $|\xi_1| \sim L$ , we may assume by rotation that  $|J| \sim L$ .

# CHANGING VARIABLES

Our task: Estimate, for 
$$|\xi_1| \sim L$$
,  $|\xi_2| \sim N$ , the integral  

$$\int_{|\xi_2^2| \lesssim N} \int_{\xi_1, \xi_2^1} d(-|\xi_1|^2 - |\xi_2|^2, \xi_1 + \xi_2) \ \widehat{f}(\xi_1) \widehat{g}(\xi_2) d\xi_1^1 d\xi_1^2 d\xi_2^1 \ d\xi_2^2.$$

We insert the Jacobian and reexpress inner integration as

$$\int_{\xi_1,\xi_2^1} d(-|\xi_1|^2 - |\xi_2|^2,\xi_1 + \xi_2) \frac{\widehat{f}(\xi_1)\widehat{g}(\xi_2)}{|J|} |J| d\xi_1^1 d\xi_1^2 d\xi_2^1.$$

Changing variables, we observe this equals

$$\int_{u,v} d(u,v)H(u,v;\xi_2^2)|J|dudv$$

where

$$H(u,v;\xi_2^2) = \frac{\widehat{f}(\xi_1)\widehat{g}(\xi_2)}{|J|}.$$

We apply Cauchy-Schwarz in u, v to bound by

$$\|d\|_{L^2} \left(\int_{u,v} |H(u,v;\xi_2^2)|^2 du dv\right)^{1/2}$$

We drop  $\|d\|_{L^2} \leq 1$  by duality and change variables back. We get

$$\left(\int\limits_{\xi_1,\xi_2^2} \left|\frac{\widehat{f}(\xi_1)\widehat{g}(\xi_2)}{|J|}\right|^2 |J|d\xi_1^1d\xi_1^2d\xi_2^1\right)^{1/2}$$

One factor of the Jacobian denominator remains! We gain  $L^{-1/2}$ . We still have the *extra outside* integration....

Recalling what we must control, using what we have obtained....

$$\int\limits_{|\xi_2^2| \lesssim N} \left( \int\limits_{\xi_1,\xi_2^2} \left| \frac{\widehat{f}(\xi_1)\widehat{g}(\xi_2)}{|J|} \right|^2 |J| d\xi_1^1 d\xi_1^2 d\xi_2^1 \right)^{1/2} d\xi_2^2.$$

Apply Cauchy-Schwarz in  $\xi_2^2$  and pay the penalty in the numerator of  $N^{1/2}$ .

We gain over the trivial bilinear estimate by the factor

$$\sqrt{\frac{(\text{measure of extra support})}{|J|}} = \sqrt{\frac{N}{L}}$$

# 4. The *I*-Method of Almost Conservation

Let  $H^s \ni u_0 \longmapsto u$  solve *NLS* for  $t \in [0, T_{lwp}], T_{lwp} \sim ||u_0||_{H^s}^{-2/s}$ . Consider two ingredients (to be defined):

- A smoothing operator  $I = I_N : H^s \mapsto H^1$ . The *NLS* evolution  $u_0 \mapsto u$  induces a smooth reference evolution  $H^1 \ni Iu_0 \mapsto Iu$  solving I(NLS) equation on  $[0, T_{Iwp}]$ .
- A modified energy  $\tilde{E}[lu]$  built using the reference evolution.

We postpone how we actually choose these objects.

For  $s < 1, N \gg 1$  define smooth monotone  $m : \mathbb{R}^2_{\mathcal{E}} \to \mathbb{R}^+$  s.t.

$$m(\xi) = egin{cases} 1 & ext{for } |\xi| < N \ \left( rac{|\xi|}{N} 
ight)^{s-1} & ext{for } |\xi| > 2N. \end{cases}$$

The associated Fourier multiplier operator,  $(Iu)(\xi) = m(\xi)\hat{u}(\xi)$ , satisfies  $I : H^s \to H^1$ . Note that, pointwise in time, we have

$$||u||_{H^s} \lesssim ||Iu||_{H^1} \lesssim N^{1-s} ||u||_{H^s}.$$

Set  $\widetilde{E}[Iu(t)] = H[Iu(t)]$ . Other choices of  $\widetilde{E}$  are mentioned later.

# AC LAW DECAY AND SOBOLEV GWP INDEX

- **1** Modified LWP. Initial  $v_0$  s.t.  $\|\nabla I v_0\|_{L^2} \sim 1$  has  $T_{Iwp} \sim 1$ .
- **2** Goal.  $\forall u_0 \in H^s, \forall T > 0$ , construct  $u : [0, T] \times \mathbb{R}^2 \to \mathbb{C}$ .
- **B**  $\iff$  **Dilated Goal.** Construct  $u^{\lambda} : [0, \lambda^2 T] \times \mathbb{R}^2 \to \mathbb{C}$ .
- **4** Rescale Data.  $\| I \nabla u_0^{\lambda} \|_{L^2} \lesssim N^{1-s} \lambda^{-s} \| u_0 \|_{H^s} \sim 1$  provided we choose  $\lambda = \lambda(N) \sim N^{\frac{1-s}{s}} \iff N^{1-s} \lambda^{-s} \sim 1$ .
- **5** Almost Conservation Law.  $||I \nabla u(t)||_{L^2} \leq H[Iu(t)]$  and

$$\sup_{t\in[0,T_{lwp}]}H[lu(t)]\leq H[lu(0)]+N^{-\alpha}$$

**6** Delay of Data Doubling. Iterate modified LWP  $N^{\alpha}$  steps with  $T_{lwp} \sim 1$ . We obtain rescaled solution for  $t \in [0, N^{\alpha}]$ .

$$\lambda^2(N)T < N^{lpha} \iff T < N^{lpha + rac{2(s-1)}{s}} ext{ so } s > rac{2}{2+lpha} ext{ suffices}.$$

A Fourier analysis established the almost conservation property of  $\tilde{E} = H[Iu]$  with  $\alpha = \frac{3}{2}$  which led to...

#### THEOREM (CKSTT 02)

 $NLS_{3}^{+}(\mathbb{R}^{2})$  is globally well-posed for data in  $H^{s}(\mathbb{R}^{2})$  for  $\frac{4}{7} < s < 1$ . Moreover,  $||u(t)||_{H^{s}} \lesssim \langle t \rangle^{\beta(s)}$  for appropriate  $\beta(s)$ .

- The smoothing property  $u(t) e^{it\Delta}u_0 \in H^1$  is not obtained.
- Same result for NLS<sub>3</sub><sup>-</sup>(ℝ<sup>2</sup>) if ||u<sub>0</sub>||<sub>L<sup>2</sup></sub> < ||Q||<sub>L<sup>2</sup></sub>. Here Q is the ground state (unique positive solution of −Q + ΔQ = −Q<sup>3</sup>).
- Fourier analysis leading to  $\alpha = \frac{3}{2}$  in fact gives  $\alpha = 2$  for most frequency interactions.

# Almost Conservation Law for H[lu]

#### PROPOSITION

Given  $s > \frac{4}{7}$ ,  $N \gg 1$ , and initial data  $\phi_0 \in C_0^{\infty}(\mathbb{R}^2)$  with  $E(I_N u_0) \leq 1$ , then there exists a  $T_{Iwp} \sim 1$  so that the solution

 $u(t,x) \in C([0, T_{lwp}], H^{s}(\mathbb{R}^{2}))$ 

of  $NLS_3^+(\mathbb{R}^2)$  satisfies

$$E(I_N u)(t) = E(I_N u)(0) + O(N^{-\frac{3}{2}+}),$$

for all  $t \in [0, T_{lwp}]$ .

# IDEAS IN THE PROOF OF ALMOST CONSERVATION

Standard Energy Conservation Calculation:

$$\partial_t H(u) = \Re \int_{\mathbb{R}^2} \overline{u_t}(|u|^2 u - \Delta u) dx \qquad \text{cancellation}$$
$$= \Re \int_{\mathbb{R}^2} \overline{u_t}(|u|^2 u - \Delta u - iu_t) dx = 0.$$

For the smoothed reference evolution, we imitate....

$$\partial_t H(lu) = \Re \int_{\mathbb{R}^2} \overline{lu_t}(|lu|^2 lu - \Delta lu - |lu_t|) dx$$
  
=  $\Re \int_{\mathbb{R}^2} \overline{lu_t}(|lu|^2 lu - l(|u|^2 u)) dx \neq 0.$ 

The increment in modified energy involves a commutator,

$$H(Iu)(t) - H(Iu)(0) = \Re \int_0^t \int_{\mathbb{R}^2} \overline{Iu_t}(|Iu|^2 Iu - I(|u|^2 u)) dx dt.$$

Littlewood-Paley, Case-by-Case, (Bi)linear Strichartz, X<sub>s,b</sub>....

## Remarks

The almost conservation property

$$\sup_{t\in[0,\mathcal{T}_{lwp}]}\widetilde{E}[lu(t)]\leq\widetilde{E}[lu_0]+N^{-\alpha}$$

leads to GWP for

$$s > s_{\alpha} = \frac{2}{2+\alpha}.$$

- The *I*-method is a subcritical method. To prove the Scattering Conjecture at s = 0 via the *I*-method would require α = +∞.
- The *I*-method *localizes the conserved density in frequency*. Similar ideas appear in recent critical scattering results.
- There is a *multilinear corrections algorithm* for defining other choices of  $\widetilde{E}$  which yield a better AC property.