

- Topic: Absolute values, intuitive definition of the limit
- **Homework:** Watch videos 2.5 and 2.6 for Wednesday.

All 0s?

Theorem

$\forall N \in \mathbb{N}$, in every set of N MAT137 students, those students will get the same grade.


Proof.

- **Base case.** It is clearly true for $N = 1$.
- **Induction step.**

Assume it is true for N . I'll show it is true for $N + 1$.

Take a set of $N + 1$ students. By induction hypothesis:

- The first N students have the same grade.
- The last N students have the same grade.



Hence the $N + 1$ students all have the same grade as well.



Properties of inequalities

Given $a, b, c \in \mathbb{R}$.

Assume $a < b$. What can we (always) conclude?

① $a + c < b + c$

② $a - c < b - c$

③ $ac < bc$

④ $a^2 < b^2$

⑤ $\frac{1}{a} < \frac{1}{b}$

Properties of absolute value

Given $a, b \in \mathbb{R}$. What can we (always) conclude?

① $|ab| = |a||b|$

② $|a + b| = |a| + |b|$

Sets described by distance

Given $a \in \mathbb{R}$. Given $\delta > 0$.


What are the following sets? Describe them in terms of intervals.

1. $A = \{x \in \mathbb{R} : |x| < \delta\}$
2. $B = \{x \in \mathbb{R} : |x| > \delta\}$
3. $C = \{x \in \mathbb{R} : |x - a| < \delta\}$
4. $D = \{x \in \mathbb{R} : 0 < |x - a| < \delta\}$

Find **all** positive values of A , B , and C which makes the following implications true.

1. $(\forall x \in \mathbb{R},) |x - 3| < 1 \implies |2x - 6| < A.$
2. $(\forall x \in \mathbb{R},) |x - 3| < B \implies |2x - 6| < 1.$
3. $(\forall x \in \mathbb{R},) |x - 3| < 1 \implies |x - 3.5| < C.$


Limits from a graph



Do the following limits exist?

- 1 $\lim_{x \rightarrow 2} f(x)$
- 2 $\lim_{x \rightarrow 0} f(f(x))$

Limits from a graph



Find the value of

- 1 $\lim_{x \rightarrow 2} f(x)$
- 2 $\lim_{x \rightarrow 0} f(f(x))$
- 3 $\lim_{x \rightarrow 2} [f(x)]^2$
- 4 $\lim_{x \rightarrow -3} f(f(x))$

Floor

Given a real number x , we defined the *floor of x* , denoted by $\lfloor x \rfloor$, as the largest integer smaller than or equal to x .

For example:

$$\lfloor \pi \rfloor = 3, \quad \lfloor 7 \rfloor = 7, \quad \lfloor -0.5 \rfloor = -1.$$

Sketch the graph of $y = \lfloor x \rfloor$. Then compute:


① $\lim_{x \rightarrow 0^+} \lfloor x \rfloor$

② $\lim_{x \rightarrow 0^-} \lfloor x \rfloor$

③ $\lim_{x \rightarrow 0} \lfloor x \rfloor$

④ $\lim_{x \rightarrow 0} \lfloor x^2 \rfloor$

More limits from a graph



Find the value of

- 1 $\lim_{x \rightarrow 0^+} g(x)$
- 2 $\lim_{x \rightarrow 0^+} \lfloor g(x) \rfloor$
- 3 $\lim_{x \rightarrow 0^+} g(\lfloor x \rfloor)$
- 4 $\lim_{x \rightarrow 0^-} g(x)$
- 5 $\lim_{x \rightarrow 0^-} \lfloor g(x) \rfloor$
- 6 $\lim_{x \rightarrow 0^-} \lfloor \frac{g(x)}{2} \rfloor$
- 7 $\lim_{x \rightarrow 0^-} g(\lfloor x \rfloor)$