





It is well known that when the Sun rises on midsummer's morning over the "Heel Stone" at Stonehenge, its first rays shine right through the open arms of the horseshoe arrangement. Thus astrological lineups, one of the pillars of modern thought, are much older than the famed Gaussian linking number of two knots.

Recall that the latter is itself an astrological construct: one of the standard ways to compute the Gaussian linking number is to place the two knots in space and then count (with signs) the number of shade points cast on one of the knots by the other knot, with the only lighting coming from some fixed distant star.



Thus we consider the generating function of all stellar coincidences:

$Z(K) := \lim_{N \rightarrow \infty} \sum_{\text{3-valent } D} \frac{1}{2^c c! \binom{N}{e}} \langle D, K \rangle_{\text{signed}} D \cdot \left( \text{framing-dependent counter-term} \right) \in \mathcal{A}(\odot)$



$N := \# \text{ of stars}$       $\mathcal{A}(\odot) := \text{Span} \left\langle \text{oriented vertices} \right\rangle$

$c := \# \text{ of chopsticks}$      AS: + = 0

$e := \# \text{ of edges of } D$      & more relations

**Theorem.** Modulo Relations,  $Z(K)$  is a knot invariant!

When deforming, catastrophes occur when:



$V$ : vector space  
 $dV$ : Lebesgue's measure on  $V$ .  
 $Q$ : A quadratic form on  $V$   
 $Q(v) = \langle Lv, v \rangle$  where  $L: V \rightarrow V^*$  is linear  
**Comaste**  $I = \int_V dv e^{\pm Q + P}$   
 $\approx \sum_{m=0}^{\infty} \frac{1}{m!} \int_V dv P^m e^{Q/2}$   
 $\approx \sum_{m=0}^{\infty} \frac{1}{m!} P^m(\partial_v) e^{\pm Q(v)/2} \Big|_{v=0}$   
 $= \sum_{m=0}^{\infty} \frac{(-1)^m}{2^m m!} P^m(a) (Q^{-1})^m \Big|_{v=0}$

In our case,  
 $\ast Q$  is  $d$ , so  $Q^{-1}$  is an integral operator.  
 $\ast P$  is  $\frac{2}{3} A^3 A^2 A$   
 $\ast H$  is the holonomy, itself a sum of integrals along the knot  $K$ .

& when the dust settles, we get  $Z(K)$ !

The Fourier Transform:  
 $(F: V \rightarrow \mathbb{C}) \Rightarrow (F: V^* \rightarrow \mathbb{C})$   
 via  $F(v) = \int_V F(v) e^{-i\langle v, v \rangle} dv$ .

Simple Facts:  
 1.  $F(0) = \int_V F(v) dv$ .  
 2.  $\frac{\partial}{\partial v} F \sim \sqrt{v} F$ .  
 3.  $(e^{Q/2}) \sim e^{-Q/2}$  where  $Q^{-1}(v) = \langle v, L^{-1}v \rangle$   
 (That's the heart of the Fourier Inversion Formula).



So  $\int_V H(v) e^{\pm Q + P} dv \sim H(a) e^{P(a)} e^{-Q^{-1}(a)/2} \Big|_{v=0}$   
 is  $\sum \text{pairings}$   
 $= \sum c(D) \left( \text{products of } Q^{-1}\text{'s, } P\text{'s (and one H)} \right)$   

Richard Feynman

It all is perturbative Chern-Simons-Witten theory:

$\int_{\text{g-connections}} \mathcal{D}A \text{ hol}_K(A) \exp \left[ \frac{ik}{4\pi} \int_{\mathbb{R}^3} \text{tr} \left( A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right]$

$\rightarrow \sum_{D: \text{Feynman diagram}} W_g(D) \int \mathcal{E}(D) \rightarrow \sum_{D: \text{Feynman diagram}} D \int \mathcal{E}(D)$



"God created the knots, all else in topology is the work of man."

Leopold Kronecker (modified)