

**Axioms.** One axiom is primary and interesting,

- ▶ Contractions commute! Namely,  $c_{x,\xi} \parallel c_{y,\eta} = c_{y,\eta} \parallel c_{x,\xi}$  (or in old-speak,  $c_{y,\eta} \circ c_{x,\xi} = c_{x,\xi} \circ c_{y,\eta}$ ).

And the rest are just what you'd expect:

- ▶  $\sqcup$  is commutative and associative, and it commutes with  $c_{\cdot,\cdot}$  and with  $\sigma_{\cdot}$  whenever that makes sense.
- ▶  $c_{\cdot,\cdot}$  is "natural" relative to renaming:  $c_{x,\xi} = \sigma_y^x \parallel \sigma_\eta^\xi \parallel c_{y,\eta}$ .
- ▶  $\sigma_\xi^\xi = \sigma_x^x = Id$ ,  $\sigma_\eta^\xi \parallel \sigma_\zeta^\eta = \sigma_\zeta^\xi$ ,  $\sigma_y^x \parallel \sigma_z^y = \sigma_z^x$ , and renaming operations commute where it makes sense.

**Comments.**

- ▶ We can relax  $|\mathcal{X}| = |X|$  at no cost.
- ▶ We can lose the distinction between  $\mathcal{X}$  and  $X$  and get "circuit algebras".
- ▶ There is a "coloured version", where  $\mathcal{T}(\mathcal{X}, X)$  is replaced with  $\mathcal{T}(\mathcal{X}, X, \lambda, l)$  where  $\lambda: \mathcal{X} \rightarrow C$  and  $l: X \rightarrow C$  are "colour functions" into some set  $C$  of "colours", and contractions  $c_{x,\xi}$  are allowed only if  $x$  and  $\xi$  are of the same colour,  $l(x) = \lambda(\xi)$ . In the world of tangles, this is "coloured tangles".

## 2. Heaven is a Place on Earth

(A version of the main results of Archibald's thesis, [Ar]).

Let us work over the base ring  $\mathcal{R} = \mathbb{Q}[\{T^{\pm 1/2}: T \in C\}]$ . Set

$$\mathcal{A}(\mathcal{X}, X) := \{w \in \Lambda(\mathcal{X} \sqcup X) : \deg_{\mathcal{X}} w = \deg_X w\}$$

(so in particular the elements of  $\mathcal{A}(\mathcal{X}, X)$  are all of even degree). The union operation is the wedge product, the renaming operations are changes of variables, and  $c_{x,\xi}$  is defined as follows. Write  $w \in \mathcal{A}(\mathcal{X}, X)$  as a sum of terms of the form  $uw'$  where  $u \in \Lambda(\xi, x)$  and  $w' \in \mathcal{A}(\mathcal{X} \setminus \xi, X \setminus x)$ , and map  $u$  to 1 if it is 1 or  $x\xi$  and to 0 if it is  $\xi$  or  $x$ :


$$1w' \mapsto w', \quad \xi w' \mapsto 0, \quad xw' \mapsto 0, \quad x\xi w' \mapsto w'.$$

**Proposition.**  $\mathcal{A}$  is a contraction algebra.

We construct a morphism of coloured contraction algebras  $\mathcal{A}: \mathcal{T} \rightarrow \mathcal{A}$  by declaring

$$\begin{aligned} X_{ijk}[S, T] &\mapsto T^{-1/2} \exp\left(\left(\begin{matrix} \xi_i & \xi_j \end{matrix} \begin{pmatrix} 1 & 1-T \\ 0 & T \end{pmatrix} \begin{pmatrix} x_j \\ x_k \end{pmatrix}\right)\right) \\ \bar{X}_{ijk}[S, T] &\mapsto T^{1/2} \exp\left(\left(\begin{matrix} \xi_i & \xi_j \end{matrix} \begin{pmatrix} T^{-1} & 0 \\ 1-T^{-1} & 1 \end{pmatrix} \begin{pmatrix} x_k \\ x_j \end{pmatrix}\right)\right) \\ P_{ij}[T] &\mapsto \exp(\xi_i x_j) \end{aligned}$$

with



(Note that the matrices appearing in these formulas are the Burau matrices).

**Alternative Formulations.**

- ▶  $c_{x,\xi} w = \iota_{\xi} \iota_x e^{x\xi} w$ , where  $\iota_{\cdot}$  denotes interior multiplication.
- ▶ Using Fermionic integration,  $c_{x,\xi} w = \int e^{x\xi} w \, d\xi \, dx$ .
- ▶  $c_{x,\xi}$  represents composition in exterior algebras! With  $X^* := \{x^* : x \in X\}$ , we have that  $\text{Hom}(\Lambda X, \Lambda Y) \cong \Lambda(X^* \sqcup Y)$  and the following square commutes:

$$\begin{array}{ccc} \text{Hom}(\Lambda X, \Lambda Y) \otimes \text{Hom}(\Lambda Y, \Lambda Z) & \xrightarrow{\parallel} & \text{Hom}(\Lambda X, \Lambda Z) \\ \updownarrow & & \updownarrow \\ \Lambda(X^* \sqcup Y \sqcup Y^* \sqcup Z) & \xrightarrow{\prod_{y \in Y} c_{y,y^*}} & \Lambda(X^*, Z) \end{array}$$

- ▶ Similarly,  $\Lambda(\mathcal{X} \sqcup X) \cong (H^*)^{\otimes \mathcal{X}} \otimes H^{\otimes X}$  where  $H$  is a 2-dimensional "state space" and  $H^*$  is its dual. Under this identification,  $c_{x,\xi}$  becomes the contraction of an  $H$  factor with an  $H^*$  factor.

**Theorem.**

If  $D$  is a classical link diagram with  $k$  components coloured  $T_1, \dots, T_k$  whose first component is open and the rest are closed, if  $MVA$  is the multivariable Alexander polynomial of the closure of  $D$  (with these colours), and if  $p_j$  is the counterclockwise rotation number of the  $j$ th component of  $D$ , then

$$\mathcal{A}(D) = T_1^{-1/2} (T_1 - 1) \left( \prod_j T_j^{p_j/2} \right) \cdot MVA \cdot (1 + \xi_{\text{in}} \wedge x_{\text{out}}).$$

( $\mathcal{A}$  vanishes on closed links).

## 3. An Implementation of $\mathcal{A}$

If I didn't implement I wouldn't believe myself.

Written in Mathematica [Wo], available as the notebook Alpha.nb at <http://drorbn.net/mo21/ap>. Code lines are highlighted in grey, demo lines are plain. We start with an implementation of elements ("Wedge") of exterior algebras, and of the wedge product ("WP"):

```
WP[Wedge[u___], Wedge[v___]] := Signature[{u, v}] * Wedge @@ Sort[{u, v}];
WP[0, _] = WP[_ , 0] = 0;
WP[A_, B_] :=
  Expand[Distribute[A ** B] /.
    (a_. * u_Wedge) ** (b_. * v_Wedge) -> a b WP[u, v]];
WP[Wedge[a_] + Wedge[b] - 2 b ^ a, Wedge[a] - 3 Wedge[b] + 7 c ^ d]
Wedge[] + Wedge[a] - 3 Wedge[b] - a ^ b + 7 c ^ d + 7 a ^ c ^ d + 14 a ^ b ^ c ^ d
```

We then define the exponentiation map in exterior algebras ("WExp") by summing the series and stopping the sum once the current term ("t") vanishes:

```
WExp[A_] := Module[{s = Wedge[a], t = Wedge[a], k = 0},
  While[t != 0, s += (t = Expand[WP[t, A] / (++k)]]; s]
WExp[a ^ b + c ^ d + e ^ f]
Wedge[] + a ^ b + c ^ d + e ^ f + a ^ b ^ c ^ d + a ^ b ^ e ^ f + c ^ d ^ e ^ f + a ^ b ^ c ^ d ^ e ^ f
```