



### Strand Doubling and Reversal.

$$\omega \begin{vmatrix} a & S \\ S & \alpha \theta \\ \phi & \Xi \end{vmatrix} \xrightarrow[\mu=T_a^{-1}]{\Delta_{bc}^a} \begin{pmatrix} \omega & b & c & S \\ b & (\sigma a - \alpha T a - \nu T c)/\mu & (T_b - 1)T c \nu/\mu & (T_b - 1)T c \theta/\mu \\ c & (T c - 1)\nu/\mu & (\alpha - \sigma a T a - \nu T c)/\mu & (T c - 1)\theta/\mu \\ S & \phi & \phi & \Xi \end{pmatrix}$$

$dS^a \downarrow T_a \rightarrow T_a^{-1}$

Where  $\sigma$  assigns to every  $a \in S$  a Laurent monomial  $\sigma_a$  in  $\{t_b\}_{b \in S}$  subject to  $\sigma(a \nearrow b, b \nearrow a) = (a \rightarrow 1, b \rightarrow t_a^{\pm 1})$ ,  $\sigma(T_1 \sqcup T_2) = \sigma(T_1) \sqcup \sigma(T_2)$ , and  $\sigma//m_c^{ab} = (\sigma \setminus \{a, b\}) \cup (c \rightarrow \sigma_a \sigma_b)_{t_a, t_b \rightarrow t_c}$ .

**Vo's Thesis [Vo].** A proof of the Fox-Milnor theorem for ribbon knots using this technology (and more).

**Implementation** key idea:  $\omega\epsilon\beta$ /AlexDemo

```

( $\omega, A = (\alpha_{ab})$ )  $\leftrightarrow$ 
( $\omega, \lambda = \sum \alpha_{ab} t_a t_b$ )

 $\Gamma := \Gamma[\omega, \lambda] := \Gamma[\omega, \lambda]$ 
 $\Gamma[\omega, \lambda] := \Gamma[\omega, \lambda]$ 
 $\Gamma[\omega, \lambda] := \Gamma[\omega, \lambda]$ 

```


**Meta-Associativity**  $\xi = \Gamma[\omega, \{t_1, t_2, t_3, t_5\}]$  **Runs.**  $\{h_1, h_2, h_3, h_5\}$

$$\xi // m_{12 \rightarrow 1} // m_{13 \rightarrow 1} = (\xi // m_{23 \rightarrow 2} // m_{12 \rightarrow 1})$$

**True** **R3** ... divide and conquer!

$\{Rm_{51} Rm_{62} Rp_{34} // m_{14 \rightarrow 1} // m_{25 \rightarrow 2} // m_{36 \rightarrow 3},$   
 $Rp_{61} Rm_{24} Rm_{35} // m_{14 \rightarrow 1} // m_{25 \rightarrow 2} // m_{36 \rightarrow 3}\}$

$\begin{pmatrix} 1 & h_1 & h_2 & h_3 \\ t_1 & T_3 & 0 & 0 \\ t_2 & \frac{-1+T_2}{T_2} & \frac{1}{T_3} & 0 \\ t_3 & \frac{-1+T_3}{T_2} & \frac{-1+T_3}{T_3} & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & h_1 & h_2 & h_3 \\ t_1 & T_3 & 0 & 0 \\ t_2 & \frac{-1+T_2}{T_2} & \frac{1}{T_3} & 0 \\ t_3 & \frac{-1+T_3}{T_2} & \frac{-1+T_3}{T_3} & 1 \end{pmatrix}$
---	---



**Fact.**  $\Gamma$  is better viewed as an invariant of a certain class of 2D knotted objects in  $\mathbb{R}^4$  [BND, BN].

**Fact.**  $\Gamma$  is the “0-loop” part of an invariant that generalizes to “ $n$ -loops” (1D tangles only, see further talks and future publications with van der Veen).

**Speculation.** Stepping stones to categorification?

**Ask me about geography vs. identity!**

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal References.*


*Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*,  $\omega\epsilon\beta$ /KBH, arXiv:1308.1721.

[BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I: w-Knots and the Alexander Polynomial*, Alg. and Geom. Top. **16-2** (2016) 1063–1133, arXiv:1405.1956,  $\omega\epsilon\beta$ /WK01.

[BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.


[GST] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjectures*, Geom. and Top. **14** (2010) 2305–2347, arXiv:1103.1601.

[Vo] H. Vo, *Alexander Invariants of Tangles via Expansions*, University of Toronto Ph.D. thesis,  $\omega\epsilon\beta$ /Vo.




### Algebraic Knot Theory


**Abstract.** This will be a very “light” talk: I will explain why about 13 years ago, in order to have a say on some problems in knot theory, I’ve set out to find tangle invariants with some nice compositional properties. In other talks in Sydney ( $\omega\epsilon\beta$ /talks) I have explained / will explain how such invariants were found - though they are yet to be explored and utilized.



**Genus.** Every knot is the boundary of an orientable “Seifert Surface” ( $\omega\epsilon\beta$ /SS), and the least of their genera is the “genus” of the knot.



**Claim.** The knots of genus  $\leq 2$  are precisely the images of 4-component tangles via




**A Bit about Ribbon Knots.** A “ribbon knot” is a knot that can be presented as the boundary of a disk that has “ribbon singularities”, but no “clasp singularities”. A “slice knot” is a knot in  $S^3 = \partial B^4$  which is the boundary of a non-singular disk in  $B^4$ . Every ribbon knot is clearly slice, yet,

**Conjecture.** Some slice knots are not ribbon.

**Fox-Milnor.** The Alexander polynomial of a ribbon knot is always of the form  $A(t) = f(t)f(1/t)$ . (also for slice)

**Theorem.**  $K$  is ribbon iff it is  $\kappa T$  for a tangle  $T$  for which  $\tau T$  is the untangle  $U$ .



**The Gold Standard** is set by the “ $\Gamma$ -calculus” Alexander formulas [BNS, BN]. An  $S$ -component tangle  $T$  has

$$\Gamma(T) \in R_S \times M_{S \times S}(R_S) = \left\{ \begin{vmatrix} \omega & S \\ S & A \end{vmatrix} \right\} \text{ with } R_S := \mathbb{Z}\langle\langle T_a : a \in S \rangle\rangle;$$

$$\left( a \nearrow b, b \nearrow a \right) \rightarrow \begin{vmatrix} 1 & a & b \\ a & 1 & 1 - T_a^{-1} \\ b & 0 & T_a^{\pm 1} \end{vmatrix} \quad T_1 \sqcup T_2 \rightarrow \begin{vmatrix} \omega_1 \omega_2 & S_1 & S_2 \\ S_1 & A_1 & 0 \\ S_2 & 0 & A_2 \end{vmatrix}$$

$$\begin{vmatrix} \omega & a & b & S \\ a & \alpha & \beta & \theta \\ b & \gamma & \delta & \epsilon \\ S & \phi & \psi & \Xi \end{vmatrix} \xrightarrow[m_c^{ab}]{T_a, T_b \rightarrow T_c} \begin{pmatrix} (1 - \beta)\omega & c & S \\ c & \gamma + \frac{\alpha\delta}{1 - \beta} & \epsilon + \frac{\delta\theta}{1 - \beta} \\ S & \phi + \frac{\alpha\psi}{1 - \beta} & \Xi + \frac{\psi\theta}{1 - \beta} \end{pmatrix}$$

For long knots,  $\omega$  is Alexander, and that’s the fastest Alexander algorithm I know! Dunfield: 1000-crossing fast.