

Some Hints for Solutions

1 Exercise #1 in handout #2

- (1) Suppose that $\varphi: \mathbb{R} \rightarrow X$ is a diffeomorphism, and let $\psi: X \rightarrow \mathbb{R}$ be its inverse. Suppose that $\varphi(0) = 0$.

Let $j: X \rightarrow \mathbb{R}^2$ be the inclusion map. Then j is smooth (why?).

By the definition of F_X , there exists a smooth map Ψ extending ψ and rendering the following diagram commutative

$$\begin{array}{ccccc}
 \mathbb{R} & \xrightarrow{\varphi} & X & \xrightarrow{\psi} & \mathbb{R} \\
 & \searrow \Phi & \downarrow j & \nearrow \Psi & \\
 & & \mathbb{R} & &
 \end{array}$$

- (2) $\Psi\Phi = \Psi j \varphi = \psi \varphi = \text{id}$ implies $d\Psi\Phi \neq 0$, namely $\Phi'(0) \neq 0$.
- (3) Connectedness consideration of $X - \{0\}$ and $\mathbb{R} - \{0\}$ implies that we may assume $\Phi((-\infty, 0)) \subseteq (-\infty, 0) \times \mathbb{R}$ and $\Phi((0, \infty)) \subseteq (0, \infty) \times \mathbb{R}$. Therefore (explain this)

$$\begin{aligned}
 \Phi'(0) &= \lim_{t \uparrow 0} \frac{\Phi(t) - \Phi(0)}{t} = (a, -a) \\
 \Phi'(0) &= \lim_{t \downarrow 0} \frac{\Phi(t) - \Phi(0)}{t} = (b, b).
 \end{aligned}$$

It follows that $a = b = 0$ hence $\Phi'(0) = 0$, a contradiction.

2 Exercise #2 in handout #3

The following argument is due to Yariv. Use paths to represent derivations. The inclusion $j: U \rightarrow M$ induces $j_*: T_p U \rightarrow T_p M$ via $j_*(D_\gamma) = D_{j\gamma}$.

Define $\pi: T_p M \rightarrow T_p U$ as follows. For $\gamma: (-a, a) \rightarrow M$ there exists ϵ such that $\gamma((-\epsilon, \epsilon)) \subset U$. Define

$$\pi(D_\gamma) = D_{\gamma|_{(-\epsilon, \epsilon)}}.$$

This is independent in the choice of ϵ and of γ (why?). It is almost trivial that j_* and π are inverses of one another.

Another easy proof can be obtained using the definition of the tangent space as in exercise #1 in this handout. Surjectivity of j_* is obvious and dimensional argument concludes.

What I said in class was rubbish. It will make sense when we talk about derivations $C^\infty(M) \rightarrow C^\infty(M)$ which correspond to vector fields. The best is yet to come.

3 Exercise #3 in handout #3

Identify $T_p\mathbb{R}^n$ with \mathbb{R}^n via

$$\frac{\partial}{\partial x_i} \mapsto e_i.$$

Under this identification, if a path $\gamma: (-a, a) \rightarrow \mathbb{R}^n$ represents a tangent vector in $T_p\mathbb{R}^n$, then γ corresponds to $\gamma'(0)$ in \mathbb{R}^n . To see this recall that using the trivial chart on \mathbb{R}^n ,

$$D_\gamma = \sum_{i=1}^n \frac{d\gamma^i}{dt}(0) \frac{\partial}{\partial x_i}.$$

In our example

$$\begin{aligned} \Psi_*\left(\frac{\partial}{\partial \theta}\right) &\equiv D_{(\sin \varphi \cos(\theta+t), \sin \varphi \sin(\theta+t), \cos \varphi) \subset S^2} \xrightarrow{j_*} \\ D_{(\sin \varphi \cos(\theta+t), \sin \varphi \sin(\theta+t), \cos \varphi) \subset \mathbb{R}^3} &\sim (-\sin \varphi \sin \theta, \sin \varphi \cos \theta, 0) \end{aligned}$$

A similar computation shows that

$$\Psi_*\left(\frac{\partial}{\partial \varphi}\right) \sim (\cos \varphi \cos \theta, \cos \varphi \sin \theta, -\sin \varphi).$$

One checks directly that these are nonzero orthogonal vectors and both are orthogonal to the vector $\Psi(\theta, \varphi)$.

This argument also shows that the inclusion map $i: S^2 \rightarrow \mathbb{R}^3$ is an immersion because S^2 is a 2-dimensional manifold and i_* has a 2 dimensional space as its image. Of course, you may (and better) use the theorem proved in class about the inverse image of a regular value.