

DEPARTMENT OF MATHEMATICS  
University of Toronto

**Analysis Exam (3 hours)**

*January 1997*

No aids.

Do all questions.

Each question is worth 20 marks.

1. A Banach space  $X$  is called uniformly convex if  $\lim_n \|x_n - y_n\| = 0$  whenever  $\{x_n\}$  and  $\{y_n\}$  are two sequences in  $X$  such that  $\|x_n\| = \|y_n\| = 1$  for each  $n$  and such that  $\lim_n \|x_n + y_n\| = 2$ . Let  $X$  be a uniformly convex Banach space.
  - a) Let  $\{x_n\}$  be a sequence in  $X$  and  $x \in X$  such that  $\lim_n \|x_n\| = \|x\|$  and such that  $\lim_n f(x_n) = f(x)$  for each  $f \in X^*$ , the dual space of  $X$ . Show that  $\lim_n \|x - x_n\| = 0$ .
  - b) Let  $K$  be a non-empty closed and convex subset of  $X$ . ( $K$  is convex means that  $ax + (1 - a)y \in K$  whenever  $x \in K$ ,  $y \in K$ , and  $a$  is a real number, such that  $0 \leq a \leq 1$ .) Show that there is a unique vector  $x_0 \in K$  such that  $\|x_0\| \leq \|x\|$  for all  $x \in K$ .
2. A bi-infinite sequence of vectors  $\{x_n\}_{n \in \mathbb{Z}}$  in a Hilbert space  $\mathcal{H}$  is called stationary if  $\langle x_n, x_m \rangle = \langle x_{n-m}, x_0 \rangle \forall m, n \in \mathbb{Z}$ . Show that  $\{x_n\}$  is stationary if and only if there is a unitary operator  $U$  on  $\mathcal{H}$  such that  $x_n = U^n x_0 \forall n \in \mathbb{Z}$ . (Recall that  $U$  unitary means  $U^*U = UU^* = \text{id}$ .) If  $\{x_n\}$  is stationary show that there is a Borel measure  $\mu$  on the unit circle  $\mathbb{T}$  such that

$$\int z^n d\mu = \langle x_n, x_0 \rangle .$$

3. Suppose  $f \in L^2(\mathbb{T})$  and let  $a_n = \hat{f}(n)$ , the  $n$ th Fourier coefficient. Prove that if  $f$  is  $C^1$  (i.e. continuously differentiable) then  $\sum_{n \in \mathbb{Z}} n^2 |a_n|^2 < \infty$ . Prove that if  $\sum_{n \in \mathbb{Z}} |na_n| < \infty$  then  $f$  is  $C^1$ .

4. a) Let  $\mu$  be Lebesgue measure on  $[0, 1]$ . Let  $\{\varphi_n\} \subset L^1(\mu)$  be an  $L^1$ -bounded sequence (i.e.  $\|\varphi_n\|_1 < C \forall n$ ) and suppose that  $\lim_{n \rightarrow \infty} \int_{[0, a]} \varphi_n(x) dx = 0$  for each  $a \in [0, 1]$ . Show that  $\lim_{n \rightarrow \infty} \int f(x) \varphi_n(x) dx = 0$  for each  $f \in C[0, 1]$ .
- b) Give an example of a uniformly bounded sequence  $\{\varphi_n\}$  such that

$$\lim_{n \rightarrow \infty} \int_{[0, a]} \varphi_n(x) dx = 0 \text{ for each } a \in [0, 1],$$

for which  $\lim_{n \rightarrow \infty} \int_{[0, 1]} |\varphi(x)| dx$  diverges and  $\varphi_n(x)$  diverges a.e. with respect to Lebesgue measure.

- c) Show that the conclusion of (a) will not hold if the assumption of  $L^1$ -boundedness is dropped.

5. Evaluate via residues  $\int_{-\infty}^{\infty} \frac{e^{itx}}{x^2+a^2} dx$ , where  $t \in \mathbb{R}$  and  $a > 0$ .

6. Let  $F$  be the family of holomorphic functions on the unit disc  $\Delta$  such that  $f(0) = 1$  and  $\operatorname{Re} f > 0$ .

- a) Show that  $\frac{1-|z|}{1+|z|} \leq |f(z)| \leq \frac{1+|z|}{1-|z|}$  for all  $f \in F$  and  $z \in \Delta$ .
- b) Deduce that  $F$  is a normal family.
- c) How large can  $|f'(0)|$  be?

Total = 120  
 $120 \times \frac{5}{6} = 100$ .