

THE RATE OF CONVERGENCE OF THE WALK ON SPHERES ALGORITHM

ILIA BINDER AND MARK BRAVERMAN

ABSTRACT. In this paper we examine the rate of convergence of one of the standard algorithms for emulating exit probabilities of Brownian motion, the Walk on Spheres (WoS) algorithm. We obtain the complete characterization of the rate of convergence of WoS in terms of the local geometry of a domain.

1. INTRODUCTION

Brownian Motion (BM) is the most important model of randomized motion in \mathbb{R}^d . It is the simplest (but, in a sense, generic) example of a continuous diffusion process. BM has found an astonishing number of application to diverse areas of Mathematics and Science, including Biomathematics, Finance, Partial Differential Equations, and Statistical Physics [EKM97, KS98, Maz02, Nel67, Szn98].

Because of the ubiquity of BM, its effective simulation provides a way to efficiently solve a variety of problems, such as computation of Conformal Maps, Tomography, and Stochastic PDEs. One of the main ways in which simulations of BM are used is to study its first hitting probabilities with respect to some stopping conditions. A particularly interesting stopping condition is of hitting the boundary of some topologically open bounded connected domain Ω . For any starting point x , the *harmonic measure* h_x on $\partial\Omega$ is given by the hitting point distribution. In many of the BM's applications, it is enough to obtain information about the harmonic measure, more specifically, to efficiently sample from it.

One of the immediate applications of the ability to sample from harmonic measures is solving the Dirichlet problem in \mathbb{R}^d . The Dirichlet problem on a domain $\Omega \subset \mathbb{R}^d$ with continuous boundary condition $f : \partial\Omega \rightarrow \mathbb{R}$ is the problem of finding a C^2 -smooth function $u : \Omega \rightarrow \mathbb{R}$ satisfying

$$(1) \quad \begin{cases} \Delta u(x) = 0 & x \in \Omega \\ u(x) = f(x) & x \in \partial\Omega \end{cases}$$

In other words, finding a harmonic function u subject to the boundary conditions f . By the celebrated Kakutani's Theorem [Kak44, GM04], the value of u at $x \in \Omega$ is exactly the expected value of f with respect to the harmonic measure h_x on $\partial\Omega$: $u(x) = \mathbf{E}_{h_x(z)}[f(z)]$.

In the present paper we study the amount of time it takes to sample from the harmonic measure with precision ε using the Walk on Spheres algorithm – the simplest and most commonly used algorithm for sampling from the harmonic measure.

2000 *Mathematics Subject Classification.* 60G42, 65C05, 31B25, 31B05.

Key words and phrases. Walk on Spheres algorithm, Harmonic measure, Potential Theory.

The first author was partially supported by NSERC Discovery grant 5810-2004-298433. This research was partially conducted during the period the second author was employed by the Clay Mathematics Institute as a Liftoff Fellow.

¹A preliminary version of this work appeared in the proceedings of the 20th ACM-SIAM Symposium on Discrete Algorithms (SODA09)

1.1. **The Walk on Spheres algorithm.** Let us now define the Walk on Spheres algorithm (WoS). We would like to simulate a BM in a given bounded domain Ω until it gets ε -close to the boundary $\partial\Omega$. Of course one could simulate it using jumps of size δ in a random direction on each step, but this would require $O(1/\delta^2)$ steps. Since we must take $\delta = O(\varepsilon)$, this would also mean that the process may take $O(1/\varepsilon^2)$ steps to converge.

The idea of the WoS algorithm is very simple: we do not care about the path the BM takes, but only about the point at which it hits the boundary. Thus if we are currently at a point $X_n \in \Omega$ and we know that

$$d(X_n) := d(X_n, \partial\Omega) \geq r,$$

i.e. that X_n is at least r -away from the boundary, then we can just jump $r/2$ units in a random direction from X_n to a point X_{n+1} . To justify the jump we observe that a BM hitting the boundary would have to cross the sphere

$$S_n = \{x : |x - X_n| = r/2\}$$

at some point, and the first crossing location X_{n+1} is distributed uniformly on the sphere. There is nothing special about a jump of $d(X_n)/2$ and it can be replaced with any $\beta d(X_n)$ where $0 < \beta < 1$.

Let $\{\gamma_n\}$ be a sequence of i.i.d. random variables each being a vector uniformly distributed on the unit sphere in \mathbb{R}^d . We could take, for example, $\gamma_n = \Gamma_n^d / |\Gamma_n^d|$, where Γ_n^d is a normally distributed d -dimensional Gaussian variable. Then, schematically, the Walk on Spheres algorithm can be presented as follows:

```

WalkOnSpheres( $X_0, \varepsilon$ )
 $n := 0$ ;
while  $d(X_n) = d(X_n, \partial\Omega) > \varepsilon$  do
  compute  $r_n$ : a multiplicative estimate on  $d(X_n)$  such that  $\beta \cdot d(X_n) < r_n < d(X_n)$ ;
   $X_{n+1} := X_n + (r_n/2) \cdot \gamma_n$ ;
   $n := n + 1$ ;
endwhile
return  $X_n$ 

```

Thus at each step of the algorithm we jump at least $\beta/2$ and at most $1/2$ -fraction of the distance to the boundary in a random direction. An example of running the WoS algorithm in 2-d is illustrated on Figure 1.




FIGURE 1. An illustration of the WoS algorithm for $d = 2$: one step jump (a), and a possible run of the algorithm for several steps (b)

As mentioned earlier, it is clear that the algorithm is correct. Moreover, it is not hard to see that it converges in $O(1/\varepsilon^2)$ steps. However, in many situations, this rate of convergence is unsatisfactory.

In particular, if we wanted to get 2^{-n} -close to the boundary, it would take us a number of steps *exponential* in n . As it turns out, in many natural situations, the rate of convergence is *polynomial* or even *linear* in n (i.e. logarithmic in $1/\varepsilon$). The object of the paper is to prove that this is the case, and give precise condition on when the faster convergence occurs.

While an actual implementation of the WoS would involve round-off errors introduced through an imperfect simulation, we will ignore those to simplify the presentation as they do not affect any of the main results. Thus the problem becomes purely that of analyzing the family of stochastic processes $\{X_t\}$ and their convergence speed to $\partial\Omega$.

Providing the domain Ω to the algorithm. It is worth noting that the algorithm needs access to the input domain Ω in a very weak sense. We need an oracle $\text{dist}_\Omega(x)$ that satisfy the following:

$$(2) \quad \text{dist}_\Omega(x) \in \begin{cases} (\beta d(x), d(x)) & \text{if } x \in \Omega, d(x) > \beta\varepsilon \\ [0, \beta\varepsilon) & \text{if } x \in \Omega, d(x) \leq \beta\varepsilon \\ 0 & \text{if } x \notin \Omega \end{cases}$$

for some $0 < \beta < 1$. Note that dist_Ω would also allow us to decide both the size of the jump on step n and whether X_n is sufficiently close to $\partial\Omega$ for the algorithm to terminate.

If Ω is given to the algorithm as a union of squares on a ε -fine grid, then dist_Ω can be computed in time $\text{poly}(1/\varepsilon)$. In many applications, however, this function can be computed in time $\text{poly}(\log 1/\varepsilon)$, because we only need to estimate the distance within a *multiplicative* error of β . The precise condition for this is that the complement set Ω^c is poly-time computable as a subset of \mathbb{R}^d in the sense of Computable Analysis. See for example [BW99, Wei00, BC06] for more details on poly-time computability of real sets. The vast majority of domains in applications satisfy this condition.

Thus, in cases when the domain Ω is sufficiently nice for Ω^c to be poly-time computable, the rate of convergence of the WoS becomes the crucial component in the running time of its execution. In particular, depending on whether the rate of convergence is $\text{poly}(1/\varepsilon)$ or $\text{poly}(\log 1/\varepsilon)$ it could take time that is either exponential or polynomial in n to sample points that are 2^{-n} -away from $\partial\Omega$.

1.2. Prior work. The WoS algorithm was first proposed in 1956 by M. Muller in [Mul56]. As we have mentioned earlier, it is not hard to see that it takes at most $O(1/\varepsilon^2)$ steps for the WoS process to reach an ε -neighborhood of $\partial\Omega$. It is also obvious that the the algorithm takes $\Omega(\log 1/\varepsilon)$ steps to finish, since the distance to the boundary is at most halved at each step. It turns out that in many situations this simple lower bound is tight.

Logarithmic rate of convergence of the process X_t to the boundary was established for *convex* domains by M. Motoo in [Mot59]. It was later generalized by G.A. Mikhailov in [Mih79] to planar domains satisfying any *cone condition* (i.e. at every point of the boundary there is a cone of certain fixed opening in the complement of the domain), as well for 3-dimensional domains satisfying a cone condition with large enough surface angle. See also [EKMS80] and [Mil95] for additional historical background and the use of the algorithm for solving various types of boundary value problems.

In our earlier work [BB07], we established polylogarithmic, but not logarithmic, upper bounds on the rate of convergence of WoS for planar domains, and for a restricted class of higher-dimensional domains. Unfortunately, the techniques of [BB07] do not generalize well to general domains in higher dimensions.

Our present results subsume all prior work on the rate of convergence of the WoS. We introduce an easily verified dimensional condition on the domain which provides tight bounds for the rates of convergence. We also show that the condition is tight. For example, we exhibit planar domains for which the rate of convergence is $\Omega(\log^2 1/\varepsilon)$.

1.3. Results. Let $H^\beta(K)$ denote the β -dimensional Hausdorff content of K , which is given by the following formula:

$$H^\beta(K) = \inf_{K \subset \cup B(x_j, r_j)} \sum r_j^\beta.$$

Intuitively, $H^\beta(K)$ measures the volume of the set K , when it is viewed as a β -dimensional set. For example, if K is the two dimensional unit disc, then $H^3(K) = 0$, but $H^2(K) > 0$.

We use Hausdorff content to introduce the notion of α -thickness.

Definition 1. A domain $\Omega \subset \mathbb{R}^d$ is said to be α -thick $0 \leq \alpha \leq d$ if there exists a constant $C > 0$ such that for every $x \in \partial\Omega$

$$H^{d-\alpha}(B(x, r) \setminus \Omega) \geq Cr^{d-\alpha}, \quad r < 1.$$

Roughly speaking, α -thick domains have complements of codimension α , which are uniformly large at every scale at every boundary point.

We call the constant c the thickness of the domain Ω . It is not hard to see that the property of α -thickness is monotone: an α -thick domain is α' -thick for $\alpha < \alpha' \leq d$.

Let us list some examples of α -thick domains.

- (1) All d -dimensional domains are d -thick;
- (2) all bounded d -dimensional domains Ω such that the complement Ω^c is connected are $d - 1$ -thick.
- (3) all convex domains and all domains satisfying *any* cone condition are 0-thick;
- (4) all domains Ω that are bounded by a smooth hypersurface $\partial\Omega$ are 0-thick.

For example, if Ω is a domain such that the complement Ω^c is connected, then each point $x \in \partial\Omega$ is connected to ∞ via a 1-dimensional curve, thus implying $d - 1$ -thickness.

It turns out that the α -thickness of the domain is responsible for the rate of convergence of the WoS algorithm. This idea is formulated precisely in our Main Theorem.

Theorem 2. Let Ω be a bounded α -thick domain in \mathbb{R}^d . Then the expected rate of convergence of the WoS from any $x \in \Omega$ until termination at distance $< \varepsilon$ to the boundary is given by the following table:

	Rate of convergence
(3) $\alpha < 2$	$O(\log 1/\varepsilon)$
$\alpha = 2$	$O(\log^2 1/\varepsilon)$
$\alpha > 2$	$O((1/\varepsilon)^{2-4/\alpha})$

The $O(\cdot)$ in the expressions above depends on the dimension d , on α , on the thickness constant C from Definition 1 and on $\beta > 0$ from the definition of the WoS. It does not depend directly on Ω .

Moreover, the rates of convergence above are tight. That is, for each α there is a family of α -thick domains Ω_n^α with some thickness C , such that the rate of convergence with $\varepsilon = 1/n$ on Ω_n^α is asymptotically given by the formulas in (3).

Recall that the rate of convergence cannot be better than $O(\log 1/\varepsilon)$. An intuitive explanation to the phase transition phenomenon occurring at $\alpha = 2$, is that a BM in \mathbb{R}^d almost surely “misses” sets of co-dimension > 2 , while hitting sets of co-dimension < 2 with positive probability.

It is worth noting that the main result in [BB07] is the special case $\alpha = d = 2$ of the theorem.

The following corollaries are implied directly by the Theorem 2.

- Corollary 3.**
- (1) Since any planar domain is 2-thick, the WoS converges in $O(\log^2 1/\varepsilon)$ steps;
 - (2) since any planar domain with connected exterior is 1-thick, the WoS converges in $O(\log 1/\varepsilon)$ steps;
 - (3) since any domain in \mathbb{R}^d is d -thick, for $d \geq 3$ the WoS converges in $O((1/\varepsilon)^{2-4/d})$ steps;

- (4) since any 3-dimensional domain with connected exterior is 2-thick, the WoS converges in $O(\log^2 1/\varepsilon)$ steps;
- (5) since for any $d \geq 4$, any d -dimensional domain with connected exterior is $d - 1$ -thick, the WoS converges in $O((1/\varepsilon)^{2-4/(d-1)})$ steps;
- (6) since any domain bounded by a smooth hypersurface is 0-thick, the WoS converges in $O(\log 1/\varepsilon)$ steps.

The rest of the paper is organized as follows. In Section 2 we construct the auxiliary boundary barrier measures and the energy functions. Using these functions, we prove the upper estimates of Theorem 2. More technical estimates on the energy function are done in Section 3. Finally, in Section 4, we present examples of α -thick domains with the slow rate of convergence of the WoS process.

Acknowledgments. We would like to thank Stephen Cook and Greg Lawler for their valuable suggestions.

2. UPPER BOUNDS: ENERGY FUNCTIONS

2.1. Construction of an auxiliary measure. In this section we will construct a family of measures near boundary points of an α -thick domain. These measures will be used to construct energy functions, which, in turn, play crucial role in the proof of Theorem 2.

Lemma 4. *There exists a constant $c = c(\alpha, d, C)$ such that for any α -thick domain Ω with thickness C in \mathbb{R}^d and for any $x \in \partial\Omega$, one can find a Borel measure μ_x which satisfies the following conditions:*

- (1) $\text{supp}(\mu_x) \cap \Omega = \emptyset$, or, equivalently, $\mu_x(\Omega) = 0$;
- (2) for any $y \in \mathbb{R}^d$ and $r > 0$, $\mu_x(B(y, r)) \leq r^{d-\alpha}$;
- (3) for any $r < 1$, $\mu_x(B(x, r)) \geq c \cdot r^{d-\alpha}$.

With a slight abuse of notation, we will also refer to the constant c from this lemma as the *thickness* of the domain.

Proof. The proof of the Lemma follows the standard reasoning that can be found in, say, Chapter II of [Car67].

Let us consider the dyadic grid selected so that the point x has coordinates $(1/3, 1/3, \dots, 1/3)$. For an integer d -multi-index $\gamma = (\gamma_1, \dots, \gamma_d)$, let $D_{k,\gamma}$ be the cube

$$\{(x_1, \dots, x_d : \gamma_n 2^{-k} \leq x_n < (\gamma_n + 1)2^{-k}, n = 1, \dots, d)\}.$$

Let $D_k(x)$ be the unique dyadic cube of the size 2^{-k} which contains x . Note that x is always at distance $2^{-k}/3$ from the boundary of $D_k(x)$.

We will construct inductively the sequence of measures ν_n . They will satisfy the following properties:

- (a) $\text{supp} \nu_n \cap \Omega = \emptyset$.
- (b) $\nu_n(D_{k,\gamma}) \leq H^{d-\alpha}(D_{k,\gamma} \setminus \Omega)$ for $1 \leq k \leq n$
- (c) $\nu_n(D_k(x)) = H^{d-\alpha}(D_k(x) \setminus \Omega)$ for $1 \leq k \leq n$

Let ν_1 be a delta measure in a point of $D_1(x) \setminus \Omega$ with the total mass $H^{d-\alpha}(D_1(x))$. It clearly satisfies all of our assumptions.

Assume now that the measure ν_n has already been constructed. The measure ν_{n+1} will be a sum of delta-measures on the points outside of Ω lying in the cubes from the $n + 1$ -st dyadic generation, such that $\nu_{n+1}(D_{n,\gamma}) = \nu_n(D_{n,\gamma})$ for all γ (so ν_{n+1} will be obtained from ν_n by re-distributing the latter over the cubes of the $(n + 1)$ -st generation). Thus the measure ν_{n+1} would automatically satisfy the second and the third condition for $k \leq n$.

To construct ν_{n+1} , we use the following rule.

First, we set $\nu_{n+1}(D_{n+1}(x)) = H^{d-\alpha}(D_{n+1}(x) \setminus \Omega)$. The measure ν_{n+1} clearly satisfies our condition (c) on $D_{n+1}(x)$.

Second, for any other dyadic cubes $D_{n+1,\gamma} \subset D_n(x)$, we assign the mass

$$(4) \quad \nu_{n+1}(D_{n+1,\gamma}) = H^{d-\alpha}(D_{n+1,\gamma} \setminus \Omega) \frac{(\nu_n(D_n(x)) - \nu_{n+1}(D_{n+1}(x)))}{\sum_{D_{n+1,\delta} \subset D_n(x), D_{n+1,\delta} \neq D_{n+1}(x)} H^{d-\alpha}(D_{n+1,\delta} \setminus \Omega)},$$

so that $\nu_n(D_n(x)) = \nu_{n+1}(D_n(x))$. By sub-additivity of the Hausdorff content,

$$\sum_{D_{n+1,\delta} \subset D_n(x)} H^{d-\alpha}(D_{n+1,\delta} \setminus \Omega) \geq H^{d-\alpha}(D_n(x)),$$

and hence

$$(5) \quad \nu_{n+1}(D_{n+1,\gamma}) \leq H^{d-\alpha}(D_{n+1,\gamma} \setminus \Omega)$$

for $D_{n+1,\gamma} \subset D_n(x)$.

Finally, for any other dyadic cubes from $(n+1)$ -st generation, we set

$$(6) \quad \nu_{n+1}(D_{n+1,\gamma}) = H^{d-\alpha}(D_{n+1,\gamma} \setminus \Omega) \frac{\nu_n(D_n)}{\sum_{D_{n+1,\delta} \subset D_n} H^{d-\alpha}(D_{n+1,\delta} \setminus \Omega)},$$

where D_n is the unique cube from the n -th dyadic generation containing $D_{n+1,\gamma}$. Using the sub-additivity of the Hausdorff content, as above, we get the estimate (5) for all cubes of the $(n+1)$ -st generation. The construction again satisfies $\nu_{n+1}(D_n) = \nu_n(D_n)$.

Let now ν be any weak* limit point of the sequence ν_n . ν is still supported outside of Ω . By the second property of the measures ν_n ,

$$(7) \quad \nu(D_{k,\gamma}) \leq \frac{\sum_{D_{k,\delta} \cap D_{k,\gamma} \neq \emptyset} H^{d-\alpha}(D_{k,\delta} \setminus \Omega)}{D_{k,\delta} \cap D_{k,\gamma} \neq \emptyset} \leq 3^d H^{d-\alpha}(D_{k,\gamma}) < 3^d (\sqrt{d})^d 2^{-k(d-\alpha)}$$

for all k . Using the third property of the measures ν_n , the α -thickness of Ω , and the fact that $D_k(x)$ contains the ball of the radius $2^{-k}/3$, we get that

$$(8) \quad \nu(D_k(x)) \geq H^{d-\alpha}(D_k(x) \setminus \Omega) \geq c 3^{-d} 2^{-k(d-\alpha)}$$

for any k .

Every ball can be covered by certain (d -dependent) number of dyadic cubes of comparable size, so (7) implies that $\nu(B(y, r)) \lesssim r^{d-\alpha}$. Every ball centered at x also contains a dyadic cube of comparable (again, d -dependent) size, hence by (8), $\nu(B(x, r)) \gtrsim r^{d-\alpha}$. Now we can set μ_x to be an appropriately normalized measure ν . □

2.2. Energy Function of optimal growth. The heart of the proof of the upper bounds in Theorem 2 is the construction of a subharmonic function with optimal growth at the boundary, the *Energy Function* U on Ω . We will construct $U(x)$ so that it is “small” in the interior of Ω , and grows to ∞ as x approaches the boundary $\partial\Omega$. The α -thickness of the domain allows us to establish that the value of $U(X_t)$ grows in expectation as the WoS progresses. Thus after a certain number of steps $U(X_t)$ will be large in expectation which would imply that X_t is close to $\partial\Omega$ with high probability.

The construction of the function is based on the notion of a *Riesz potential*. For a finite Borel measure μ on \mathbb{R}^d , and $\alpha < d$, the α -Riesz potential of the measure μ is defined by

$$U_\alpha^\mu(x) = \frac{1}{d-\alpha} \int \frac{d\mu(z)}{|z-x|^{d-\alpha}}.$$

For $\alpha = d$, the d -Riesz potential is defined by

$$U_\alpha^\mu(x) = \int \log \frac{1}{|z-x|} d\mu(z).$$

The value $U_\alpha^\mu(x) = \infty$ is allowed when the integral diverges.

An important special case is the case of $\alpha = 2$, the so-called Newton potential. We will denote U_2^μ simply by U^μ . In this case the expression under the integral is harmonic in \mathbb{R}^d . It is well known (e.g. see [Lan72]) that the function U^μ is superharmonic on \mathbb{R}^d , and harmonic outside of $\text{supp } \mu$.

More generally, outside of the $\text{supp } \mu$, we have the identity

$$(9) \quad \Delta U_\alpha^\mu(y) = (d - \alpha + 2)(2 - \alpha)U_{\alpha-2}^\mu(y).$$

It shows that for $0 < \alpha < 2$, the function U_α^μ is subharmonic outside of $\text{supp } \mu$.

The following important technical identity, which easily follows from Fubini's Theorem and substitution, relates the local behavior of the measure μ and the growth of its potential U_α^μ . For $\alpha < d$, we have

$$(10) \quad U_\alpha^\mu(y) = \frac{1}{d - \alpha} \int_0^\infty \mu(B(y, t^{-1/(d-\alpha)})) dt = \int_0^\infty \frac{\mu(B(y, r))}{r^{d-\alpha+1}} dr,$$

and for $\alpha = d$,

$$(11) \quad U_\alpha^\mu(y) = \int_{-\infty}^\infty \mu(B(y, e^{-t})) dt = \int_0^\infty \frac{\mu(B(y, r))}{r} dr$$

Let us now fix an α -thick domain $\Omega \subset B(0, 1) \subset \mathbb{R}^d$. Let us consider the set \mathcal{M} of all Borel measures μ supported inside $\overline{B(0, 2)}$ and outside of Ω (i.e. $\mu(\Omega) = 0$), satisfying the following condition:

$$(12) \quad \text{for any } y \in \mathbb{R}^d \text{ and } r > 0, \mu(B(y, r)) \leq r^{d-\alpha}$$

Let us now introduce the *Energy Function* $U(y)$. Recall that $U^\mu(y) := U_2^\mu(y)$.

$$(13) \quad U(y) := \begin{cases} \sup_{\mu \in \mathcal{M}} U_\alpha^\mu(y), & \text{when } \alpha \leq 2 \\ \sup_{\mu \in \mathcal{M}} U^\mu(y), & \text{when } \alpha \geq 2. \end{cases}$$

Since the set \mathcal{M} is weakly*-compact, for every $y \in \partial\Omega$ there exists a measure maximizing the potential in (13) at the point y .

Let us summarize the properties of $U(y)$ in the following claim. The proof uses the identities (10) and (11). Recall that $d(y) = \text{dist}(y, \partial\Omega)$.

Claim 5. *Let Ω be an α -thick domain. Then*

- (1) $U(y)$ is subharmonic in Ω .
- (2) For $\alpha \leq 2$, $U(y) \leq \log \frac{2}{d(y)}$ for all $y \in \Omega$.
- (3) For $\alpha > 2$, $U(y) \leq \frac{1}{\alpha - 2} d(y)^{2-\alpha}$ for all $y \in \Omega$.

Proof. Let $\alpha \leq 2$, $y \in \Omega$ and $\mu \in \mathcal{M}$. Equations (10) and (11) imply that

$$(14) \quad U_\alpha^\mu(y) \leq \int_{d(y)}^2 \frac{1}{t} dt = \log \frac{2}{d(y)}.$$

Similarly, for $\alpha > 2$, we will use the harmonic potential U^μ . Let $\alpha > 2$, $y \in \Omega$ and $\mu \in \mathcal{M}$. Once again, (12), $\text{supp } \mu \cap \Omega = \emptyset$, and the equations (10) imply that

$$(15) \quad U^\mu(y) \leq \int_{d(y)}^2 \frac{1}{t^{\alpha-1}} dt \leq \frac{1}{(\alpha - 2)d(y)^{\alpha-2}}.$$

By equations (14) and (15), $U(y)$ is a supremum of a locally bounded family of subharmonic functions. Thus $U(y)$ is subharmonic.

The second and third statements of the claim follow directly from (14) and (15) respectively. \square

Let X_t be the WoS process initiated at some point $X_0 = y \in \Omega$. Let us define a new process $U_t = U(X_t)$, the value of the energy function at the t -th step of the process. Note that because U is subharmonic, U_t is a submartingale, that is $\mathbf{E}[U_{t+1}|U_t] \geq U_t$.

For the rest of the section let $n = 1/\varepsilon$. Claim 5 immediately implies that a large value of U_t will guarantee the closeness to the boundary. More specifically,

Claim 6. *For $\alpha \leq 2$, if $U_t > \log 2n$ then $d(X_t) < 1/n$.*

For $\alpha > 2$, $U_t > (\alpha - 2)n^{\alpha-2}$ implies $d(X_t) < 1/n$.

The proof of Theorem 2 relies on finer lower bounds on the function U , which would guarantee the optimal rate of boundary convergence. We prove the bounds in the next section. These bounds depend heavily on the value of α . We first give a probabilistic proof of the upper bounds in Theorem 2, and then prove the finer estimates on U in Section 3.

2.3. Logarithmic convergence: the case $\alpha < 2$. In the heart of the proof for this case lies the following strong estimate on the behavior of the Riesz potentials near the boundary.

Lemma 7. *For any $\alpha < 2$ and $c > 0$, there exist two constants δ and η , such that the following holds.*

Let Ω be an α -thick domain in \mathbb{R}^d with thickness c . Let $y \in \Omega$ and $x \in \partial\Omega$ be the closest point to y . Let $\mu \in \mathcal{M}$.

Then either

$$(16) \quad U(z) > U_\alpha^\mu(z) + 1 \text{ whenever } \delta/4 \cdot d(y) < |z - x| < \delta \cdot d(y).$$

or

$$(17) \quad \mu(B(y, 2d(y))) \geq \eta d(y)^{d-\alpha}$$

The lemma is established in Section 3.1.

Note that after $k = O(|\log \delta|)$ steps of the WoS process,

$$(18) \quad \delta/4 \cdot d(X_t) < |X_{t+k} - x| < \delta \cdot d(X_t) \text{ with a certain probability } p,$$

where x is the point of $\partial\Omega$ that is closest to X_t , and $p > 0$ depends only on β and the dimension d .

Let us fix X_t and take the measure $\mu \in \mathcal{M}$ maximizing the value of $U_\alpha^\mu(X_t)$. By the preceding observation, in the first case in Lemma 7, the subharmonicity of U implies that the expectation of U_{t+k} , conditioned on U_t , will increase by some definite constant.

On the other hand, using the identity (9) and the α -thickness of Ω , one can see that the Laplacian of U_α^μ is large near the point X_t in the second case of Lemma 7. Thus, since large Laplacian leads to a fast build-up of mean values, we have the above-mentioned increase by a constant after the first step. We arrive at the following estimate, which shows that U_t grows at least linearly in expectation.

Lemma 8. *There are constants L and k , depending only on c , β , and α , such that*

$$\mathbf{E}[(U_{t+k} - U_t)|U_t] > L.$$

A detailed proof of the lemma can be found in Section 3.2.

Lemma 8 implies that $\mathbf{E}[U_t] > tL/k + U_0$. Since $d(X_t) \geq (1 - \beta)^t d(X_0)$, Claim 6 implies that $U_t \leq U_0 + t|\log(1 - \beta)| + \log 2$. This implies that $U_t > U_0 + tL/2k$ with probability at least P , where P depends only on β . This, together with Claim 6 implies the necessary upper bound in the case $\alpha < 2$.

2.4. Polylogarithmic convergence: the case $\alpha = 2$. In the case $\alpha = 2$ the steady linear growth of U_t given by Lemma 8 no longer holds. In fact, the only thing that generally holds in this case is the submartingale property $\mathbf{E}[U_{t+1}|U_t] \geq U_t$. We are able to overcome this difficulty by showing that the submartingale process $\{U_t\}$ has a deviation bounded from below by a constant at every step. To this end it suffices to show that U_t can grow by some η with a non-negligible probability. We use the following estimate on the energy function (established in Section 3.1).

Lemma 9. *There exists a constant δ , dependent only on the thickness c , such that the following holds. Let Ω be a 2-thick domain. Let $y \in \Omega$ and $x \in \partial\Omega$ be the closest point to y . Then*

$$(19) \quad U(z) > U(y) + 1 \text{ whenever } |z - x| < \delta \cdot d(y).$$

Since the function U is subharmonic, observation (18) implies the following estimate (see Section 3.2 for a proof).

Lemma 10. *Let Ω be a 2-thick domain in \mathbb{R}^d . There are constants k and L , depending only on the thickness c , the jump ratio β , and the dimension d , such that*

$$\mathbf{E}[(U_{t+k} - U_t)^2|U_t] > L.$$

We can now use Lemma 10 to prove the upper bounds on the rate of convergence for the case $\alpha = 2$. Let us replace the submartingale U_t by a stopped submartingale

$$V_t = \begin{cases} U_t, & t < T_n \\ U_{T_n}, & t \geq T_n \end{cases}$$

By the optional stopping time theorem (see [KS91]), V_t is also a positive submartingale; $V_t \leq \log \frac{4}{n}$. This implies, in particular, that

$$(20) \quad \mathbf{E}[V_t(V_{t+k} - V_t)] = \mathbf{E}[\mathbf{E}[V_t(V_{t+k} - V_t)|V_t]] \geq \mathbf{E}[\mathbf{E}[V_t(V_t - V_t)|V_t]] = 0$$

Lemma 10 implies that

$$(21) \quad \mathbf{E}[(V_{t+k} - V_t)^2] > L \cdot \mathbf{P}[T_n > t + k].$$

We are now in a position to establish the upper bounds for $\alpha = 2$.

Proof of the upper bound from Theorem 2 for $\alpha = 2$. Assume first that for some M ,

$$\mathbf{P}[T_n > M \log^2 n] \geq 1/2.$$

It means that for all $t \leq M \log^2 n - k$, $\mathbf{P}[T_n \geq t + k] \geq 1/2$. This implies

$$\mathbf{E}[V_{t+k}^2] = \mathbf{E}[(V_{t+k} - V_t + V_t)^2] = \mathbf{E}[V_t^2] + \mathbf{E}[(V_{t+k} - V_t)^2] + 2\mathbf{E}[V_t(V_{t+k} - V_t)] \geq \mathbf{E}[V_t^2] + L/2.$$

The last inequality follows from (20) and (21). Hence $\mathbf{E} \left[V_{M \log^2 n}^2 \right] \geq \frac{LM \log^2 n}{2k}$. Since $V_t \leq \log \frac{4}{n}$, this leads to a contradiction for large enough M . \square

2.5. Polynomial convergence: the case $\alpha > 2$. For the case $\alpha > 2$, the required converse to Claim 6 is relatively simple.

Lemma 11. *For $\alpha > 2$, and an α -thick domain Ω in \mathbb{R}^d with the thickness c ,*

$$U(y) \geq K \cdot d(y)^{2-\alpha}$$

for all $y \in \Omega$. Here the constant $K = K(c, \alpha)$ depends only on c and α .

The lemma is established in Section 3.1.

The idea of the proof of Theorem 2 in this case is now as follows. When the WoS is far from the boundary $\partial\Omega$ it makes fairly big steps and when it is close it makes small steps. There are not too many big steps because the number of big steps of length $> \varepsilon$ confined to $B(0, 1)$ is bounded by $O(1/\varepsilon^2)$. On the other hand, there are not too many small steps, because a small step means that the WoS is very close to $\partial\Omega$, and should converge before an opportunity to make many more steps.

More precisely, the number of “big” jumps is bounded by the following Claim.

Claim 12. *Let $N(\varepsilon, T)$ be the number of the jumps in the WoS process before the time t which are bigger than ε , i.e.*

$$N(\varepsilon, T) = \#\{t \leq T \mid |X_t - X_{t-1}| \geq \varepsilon\}.$$

Then

$$\mathbf{P} \left[N(\varepsilon, T) > \frac{4}{\varepsilon^2} \right] < 1/4.$$

Proof. Note now that because X_t is a martingale, we have

$$(22) \quad 1 \geq \mathbf{E}[X_T^2] - X_0^2 = \sum_{k=1}^T (\mathbf{E}[X_k^2] - \mathbf{E}[X_{k-1}^2]) = \sum_{t=1}^T \mathbf{E}[(X_t - X_{t-1})^2] =$$

$$\mathbf{E} \left[\sum_{t=1}^T (X_t - X_{t-1})^2 \right] \geq \varepsilon^2 \mathbf{E}[N(\varepsilon, T)]$$

The last equation implies the statement of the claim, by Tschebyshev inequality. \square




FIGURE 2. The regions R_k from the proof of the upper bounds for $\alpha > 2$

To bound the number of small jumps, we denote by $R_0 \subset \Omega$ the $1/n$ -neighborhood of $\partial\Omega$, and more generally, by

$$R_k := \{x \in \Omega : 2^{k-1}/n < d(x, \partial\Omega) \leq 2^k/n\}$$

(see Figure 2). Note that by Lemma 11, we have

$$(23) \quad 2^{(k-1)(2-\alpha)} n^{\alpha-2} \geq U(y) \geq K 2^{k(2-\alpha)} n^{\alpha-2}$$

for $y \in R_k$. Using this fact we prove the following.

Claim 13. Denote by v_k the number of visits of X_t to R_k before the time T when X_t first hits the $1/n$ -neighborhood of the boundary $\partial\Omega$,

$$v_k = \#\{t < T : X_t \in R_k\}.$$

Then

$$\mathbf{P}[v_k > C_2 \cdot 2^{k(\alpha-2)} M] < 1/4^M,$$

for some constant $C_2 = C_2(c, d, \alpha, \beta)$ and for any $M > 1$.

Proof. Suppose that at some point t , $X_t \in R_k$. We estimate from below the probability that this is the *last* time the WoS visits R_k .

First of all, with some probability $p = p(\alpha, \beta, c) > 0$ and for some constant η , $X_{t+\eta} \in R_{k+2\log c}$, i.e. the series of first jump brings us much closer to $\partial\Omega$. Consider the subharmonic function

$$\Phi(y) = (2n)^{2-\alpha} U(y) - 2^{k(2-\alpha)}.$$

in Ω . Then the process $\Phi(X_{t+j})$ is a submartingale. We stop it at time $t + \tau$, $\tau \geq \eta$, when either the WoS terminates or when $X_{t+\tau} \in R_k$ (i.e. the process gets back to R_k), whichever comes first. If $X_{t+\tau}$ is $1/n$ -close to $\partial\Omega$ (but not closer than $1/2n$), then $\Phi(X_{t+\tau}) \leq 1$, by (23). If $X_{t+\tau} \in R_k$, then, again by (23), $\Phi(X_{t+\tau}) \leq 0$. Another application of (23) implies that if $y \in R_{k+2\log c}$, then $\Phi(y) > \gamma$ for some constant γ . Thus the probability that the WoS terminates at $X_{t+\tau}$ (i.e. we never visit R_k again) is at least

$$\mathbf{P}[X_{t+\tau} \notin R_k] \geq \mathbf{E}[\Phi(X_{t+\tau})] \geq \Phi(X_{t+\eta}) \geq p\gamma.$$

Thus the probability that the visit X_t to R_k is the last one is at least $p \cdot \gamma$. The claim now follows from an estimate of the probability of having at least v_k returns to R_k , each of them *not* being the last one. \square

Claims 12 and 13 together imply the upper bounds on the rate of convergence for $\alpha > 2$.

Proof of the upper bounds from Theorem 2 for $\alpha > 2$. By Claim 13, for any k , we have that

$$\mathbf{P}[v_{k-s} > C_2 \cdot 2^{(k-s)(\alpha-2)} \cdot (3/2 + s/2)] < 1/4^{3/2+s/2} = (1/8) \cdot 2^{-s}.$$

Hence, by union bound $v_{k-s} \leq C_2 \cdot 2^{(k-s)(\alpha-2)} \cdot (3/2 + s/2)$ for all $s \geq 0$ with probability at least $3/4$. Let k be such that $2^k \approx n^{2/\alpha}$. Then, with the probability at least $3/4$, we have the total number of jumps smaller than $2^k/n$ bounded by

$$(24) \quad \sum_{s=0}^k v_{k-s} \leq \sum_{s=0}^k C_2 \cdot 2^{(k-s)(\alpha-2)} \cdot (3/2 + s/2) < 4C_2 \cdot 2^{k(\alpha-2)} \approx 4C_2 \cdot n^{2-4/\alpha}.$$

If we take $N = (C_1 + 8C_2)n^{2-4/d}$ steps of the WoS, (24) implies that at least half the steps would be of magnitude at least $2^k/n \approx n^{2/d-1}$, except with probability $< 1/4$. Applying the estimate from Claim 12 with $\varepsilon = 2^k/n$, we see that with the probability at least $3/4$, $N(2^k/n, t) \leq 4n^2/2^{2k} \approx 4n^{2-4/\alpha}$. Hence with probability $\geq 1/2$ the WoS terminates after $O(n^{2-4/\alpha})$ steps. \square

3. BOUNDARY BEHAVIOR OF THE ENERGY FUNCTION

In this section we prove the analytical estimates on the behavior of the energy function that have been used in Section 2.

3.1. Estimating boundary growth. We start with the easiest case $\alpha > 2$.

Lemma 11 (Section 2.5): For $\alpha > 2$, and an α -thick domain Ω in \mathbb{R}^d with the thickness c ,

$$U(y) \geq K \cdot d(y)^{2-\alpha}$$

for all $y \in \Omega$. Here the constant $K = K(c, \alpha)$ depends only on c and α .

Proof. Let x be the closest to y point at $\partial\Omega$, and let $\mu = \mu_x$ be the corresponding measure from the definition of the α -thick domains. Then, by the identity (10) and since $B(x, r) \subset B(y, r + d(y))$,

$$(25) \quad U(y) \geq U_2^\mu(y) = \int_{d(y)}^2 \frac{\mu(B(y, r))}{r^{d-1}} dr \geq \int_{2d(y)}^2 \frac{\mu(B(x, r - d(x)))}{r^{d-1}} dr \geq \int_{2d(y)}^2 \frac{\mu(B(x, r/2))}{r^{d-1}} \geq c2^{\alpha-d} \int_{2d(y)}^2 t^{1-\alpha} \geq K \cdot d(y)^{2-\alpha}.$$

□

Unfortunately, in the case $\alpha \leq 2$ lower bounds of the type established in the proof of Lemma 11 are insufficient, and we will use finer estimates provided by the following construction.

Let y be a point in Ω , x be the point of $\partial\Omega$ that is the closest to y , and μ be a measure in the class \mathcal{M} . We construct a new measure $\nu \in \mathcal{M}$, which we call the *amalgamation* of μ at the point y in the following way.




FIGURE 3. Construction of the amalgamation $\nu = \sum_k \mu_k$

Let measure μ_1 be the measure μ_x from Lemma 4 restricted to $B(y, 2d(y))$, $\mu_2 = \mu_3 = 0$, and for $k \geq 4$, let μ_k be the measure μ restricted to the d -dimensional annulus

$$A_k = \{w : 2^{k-1}d(y) \leq |w - y| \leq 2^k d(y)\}$$

scaled by the factor $1 - \gamma_k := 1 - 2^{(4-k)(d-\alpha)}$. Let us also put $\gamma_1 = \gamma_2 = \gamma_3 = 1$. We define

$$\nu := \sum_k \mu_k.$$

The ingredients of the construction are illustrated on Figure 3.

Let us now prove that $\nu \in \mathcal{M}$. Consider any disk $B(w, r)$. Let K be the largest number such that $B(w, r)$ intersects A_K . If $B(w, r)$ does not intersect $B(y, 2d(y))$, the measure ν is no greater than μ on $B(w, r)$, and thus $\nu(B(w, r)) \leq r^{d-\alpha}$. If $K \leq 3$, $\nu(B(w, r)) \leq \mu_x(B(w, r)) \leq r^{d-\alpha}$. For all other cases, $r \geq 2^{K-3}d(y)$, which, by the choice of γ_K , implies that $\gamma_K r^{d-\alpha} \geq (2d(y))^{d-\alpha}$. Thus

$$\begin{aligned} \nu(B(w, r)) &\leq \mu_x(B(y, 2d(y))) + \mu(B(w, r)) - \sum_{k=1}^K \gamma_k \mu(B(w, r) \cap A_k) \leq \\ &(2d(y))^{d-\alpha} + (1 - \gamma_K)\mu(B(w, r)) \leq \gamma_K r^{d-\alpha} + (1 - \gamma_K)r^{d-\alpha} = r^{d-\alpha}. \end{aligned}$$

The second inequality follows from the fact that the sequence $\{\gamma_k\}$ is non-increasing. We first apply the amalgamation construction to the case $\alpha = 2$.

Lemma 9 (Section 2.4): There exists a constant δ , dependent only on the thickness c , such that the following holds. Let Ω be a 2-thick domain. Let $y \in \Omega$ and $x \in \partial\Omega$ be the closest point to y . Then

$$(26) \quad U(z) > U(y) + 1 \text{ whenever } |z - x| < \delta \cdot d(y).$$

Proof. Since \mathcal{M} is a compact set, $U(y) = U^\mu(y)$ for some $\mu \in \mathcal{M}$.

Let μ_0 be the restriction of the measure μ to $B(0, 2) \setminus B(y, 2d(y))$. By (10) and (12)

$$(27) \quad U^{\mu_0}(y) \geq U^\mu(y) - \log 2.$$

Let ν be the amalgamation of μ at y . Next, we will show that

$$(28) \quad U^\nu(z) \geq U^{\mu_0}(z) - C_1 + c \cdot 2^{2-d} \log \frac{1}{\delta}$$

and

$$(29) \quad U^{\mu_0}(z) \geq U^{\mu_0}(y) - C_2$$

whenever $|z - x| < \delta d(y)$ for some constants C_1 and C_2 depending only on d and c . These inequalities, together with (27), imply the statement of the lemma whenever δ is sufficiently small (namely, when $\log 1/\delta > 2^{d-2}(C_1 + C_2 + 1 + \log 2)/c$).

To establish (28), let us note that for any k we have

$$\mu(A_1) + \mu(A_2) + \cdots + \mu(A_k) = \mu(B(y, 2^k d(y))) \leq (2^k d(y))^{d-2}.$$

By the Abel summation formula,

$$\sum_k \gamma_k 2^{k(2-d)} \mu(A_k) \leq \sum_k d(y)^{d-2} (2^{d-2} \gamma_{k-1} - \gamma_k) \cdot 2^{2-d} \leq 5(d(y))^{d-2}.$$

This implies

$$(30) \quad \frac{1}{d-2} \sum_{k \geq 2} \gamma_k \int_{A_k} \frac{1}{|w-z|^{d-2}} d\mu_0(w) \leq \sum_{k \geq 2} \gamma_k \mu(A_k) (2^{k-2} d(y))^{2-d} \leq 5 \cdot 4^{d-2}.$$

Thus we obtain

$$\begin{aligned}
U^\nu(z) &\geq \int_{2\delta d(y)}^{d(y)} \frac{\mu_x(B(z, r))}{r^{d-1}} dr + \sum_{k \geq 2} \int_0^\infty \frac{\mu_k(B(z, r))}{r^{d-1}} dr \geq \\
&\int_{2\delta d(y)}^{d(y)} \frac{\mu_x(B(x, r - \delta d(y)))}{r^{d-1}} dr + \int_0^\infty \frac{\mu_0(B(z, r))}{r^{d-1}} dr - \frac{1}{d-2} \sum_{k \geq 1} \gamma_k \int_{A_k} \frac{d\mu_0(w)}{|w-z|^{d-2}} \geq \\
&\int_{2\delta d(y)}^{d(y)} c \cdot \left(\frac{r - \delta d(y)}{r} \right)^{d-2} \frac{dr}{r} + \int_0^\infty \frac{\mu_0(B(z, r))}{r^{d-1}} dr - 5 \cdot 4^{d-2} \geq c \cdot 2^{2-d} \log \frac{1}{2\delta} + U^{\mu_0}(z) - 5 \cdot 4^{d-2}.
\end{aligned}$$

which implies (28).

To obtain (29), note that for any point $w \in [y, z]$, for $d > 2$ we have, by the estimate (12)

$$\begin{aligned}
(31) \quad |\nabla U^{\mu_0}(w)| &\leq \frac{1}{d-2} \int \left| \nabla_w \frac{1}{|\xi - w|^{d-2}} \right| d\mu_0(\xi) = \int \frac{1}{|\xi - w|^{d-1}} d\mu_0(\xi) = \\
&(d-1) \int_0^\infty \frac{\mu_0(B(w, r))}{r^d} dr = (d-1) \int_{d(y)}^\infty \frac{\mu_0(B(w, r))}{r^d} dr \leq (d-1) \int_{d(y)}^\infty \frac{r^{d-2}}{r^d} dr \leq \frac{A}{d(y)}
\end{aligned}$$

for some constant A depending only on d and c . The same inequality is derived similarly in the case $d = 2$. This implies that

$$U^{\mu_0}(z) - U^{\mu_0}(y) = \int_{[y, z]} \nabla U^{\mu_0}(w) \cdot dw \geq -|z - y| \frac{A}{d(y)} \geq -A,$$

which is exactly the equation (29). \square

Another application of the amalgamation construction will establish the lower bounds required in the case $\alpha < 2$.

Lemma 7 (Section 2.3): For any $\alpha < 2$ and $c > 0$, there exist two constants δ and η , such that the following holds.

Let Ω be an α -thick domain in \mathbb{R}^d with thickness c . Let $y \in \Omega$ and $x \in \partial\Omega$ be the closest point to y . Let $\mu \in \mathcal{M}$.

Then either

$$(32) \quad U(z) > U_\alpha^\mu(z) + 1 \text{ whenever } \delta/4 \cdot d(y) < |z - x| < \delta \cdot d(y).$$

or

$$(33) \quad \mu(B(y, 2d(y))) \geq \eta d(y)^{d-\alpha}$$

Proof. Let $\eta = \delta^{d-\alpha+2}$. Assume that $\mu(B(y, 2d(y))) < \eta d(y)^{d-\alpha}$. Let μ_0 be the restriction of μ to $B(0, 2) \setminus B(y, 2d(y))$, as in the previous lemma. We have, by (10),

$$(34) \quad U_\alpha^{\mu_0}(z) \geq U_\alpha^\mu(z) - \int_{d(z)}^{2d(y)} \frac{\mu(B(z, r))}{r^{d-\alpha+1}} dr \geq U_\alpha^\mu(z) - 2d(y)\delta.$$

On the other hand, the same reasoning as the proof of (28) above, gives

$$(35) \quad U(z) \geq U_\alpha^\nu(z) \geq U_\alpha^{\mu_0}(z) - C_1 + c \cdot 2^{\alpha-d} \log \frac{1}{\delta}$$

for some constant C_1 depending only on d , α , and c . Here, as in (28), ν is the amalgamation of μ at y .

Estimates (34) and (35) together imply the statement of the lemma. \square

3.2. The boundary drift of the WoS process: $\alpha \leq 2$. First we establish that the process U_t has the drift toward the boundary in the case $\alpha < 2$.

Lemma 8 (Section 2.3): There are constants L and k , depending only on c , β , and α , such that

$$\mathbf{E}[(U_{t+k} - U_t)|U_t] > L.$$

Proof. Let us fix X_t . By weak*-compactness of the set \mathcal{M} , there exists a measure μ such that $U_\alpha^\mu(X_t) = U(X_t) = U_t$. By Lemma 7, either

$$(36) \quad U(z) > U_\alpha^\mu(z) + 1 \text{ whenever } \delta/2 \cdot d(X_t) < |z - x| < \delta \cdot d(X_t).$$

where x is the closest to X_t point on $\partial\Omega$, or

$$(37) \quad \mu(B(X_t, 2d(X_t))) \geq \eta d(X_t)^{d-\alpha}$$

Let us start with the first case.

For some $p > 0$ dependent only on d and β ,

$$(1 - \beta/2)^{-k} d(X_t)/2 < \mathbf{P}[|X_{t+k} - x| < (1 - \beta/2)^{-k} d(X_t)] > p^k.$$

Hence, for sufficiently large k ,

$$(38) \quad \mathbf{P}[\delta/2 \cdot d(X_t) < |X_{t+k} - x| < \delta \cdot d(X_t)] > p^k.$$

Let us now observe that by subharmonicity of the functions U and U_α^μ , the previous estimate, the fact that $U \geq U_\alpha^\mu$ and the assumption (34),

$$(39) \quad \mathbf{E}[(U_{t+k} - U_t)|X_t] = \mathbf{E}[(U_\alpha^\mu(X_{t+k}) - U_\alpha^\mu(X_t))|X_t] + \mathbf{E}[U(X_{t+k}) - U_\alpha^\mu(X_{t+k})|X_t] \geq \\ \mathbf{E}[U(X_{t+k}) - U_\alpha^\mu(X_{t+k})|X_t \text{ and } \delta/2 \cdot d(X_t) < |X_{t+k} - x| < \delta \cdot d(X_t)] > p^k.$$

Since the value of X_t determines the value of U_t , this establishes the statement of Lemma in the first case (with $L = p^k$).

Now let us consider the second case. By the Green formula, for a C^2 -smooth function u ,

$$(40) \quad \mathbf{E}[u(X_{t+1})|X_t] - u(X_t) = \int_{\beta d(X_t) S^d} u(y) dS(y) - u(X_t) = \int_0^{\beta d(X_t)} r^{1-d} \int_{B(X_t, r)} \Delta u(y) dV(y) dr$$

where S^d is the unit sphere in \mathbb{R}^d with the normalized Lebesgue measure S , and dV is the volume element in \mathbb{R}^d .

Note that by (10) and (9), for $|y - X_t| \leq \beta d(X_t)$ we have

$$(41) \quad \Delta U_\alpha^\mu(y) = (d - \alpha + 2)(2 - \alpha)U_{\alpha-2}^\mu(y) = \\ (d - \alpha + 2)(2 - \alpha) \int_0^\infty \frac{\mu(B(y, r))}{r^{d-\alpha+3}} dr \geq \\ (d - \alpha + 2)(2 - \alpha) \mu(B(X_t, 2d(X_t))) \int_{(2+\beta)d(X_t)}^\infty \frac{1}{r^{d-\alpha+3}} dr \geq \\ \eta(2 - \alpha) \frac{\mu(B(X_t, 2d(X_t)))}{((2 + \beta)d(X_t))^{d-\alpha+2}} \geq C_1(d(X_t))^{-2}$$

for some constant C_1 depending only on d , α , and β .

So, using (40), applied to $u = U^\mu$ and (41), we get

$$(42) \quad \mathbf{E}[U_{t+k}|X_t] - U_t \geq \mathbf{E}[U_{t+1}|X_t] - U_t \geq \mathbf{E}[U^\mu(X_{t+1})|X_t] - U^\mu(X_t) \geq \\ \int_0^{\beta d(X_t)} r^{1-d} (C_2(d(X_t))^{-2} r^d) dr = C_2 ((d(X_t))^2) (d(X_t))^{-2} \beta^2/2 = L$$

for some constants C_2 and L depending only on d , α , and β . Since again the value of X_t determines the value of U_t , the Lemma follows. \square

Let us now turn to the case $\alpha = 2$.

Lemma 10 (Section 2.4): Let Ω be a 2-thick domain in \mathbb{R}^d . There are constants k and L , depending only on the thickness c , the jump ratio β , and the dimension d , such that

$$\mathbf{E}[(U_{t+k} - U_t)^2 | U_t] > L.$$

Proof. Fix X_t . By Lemma 9, there exists a constant δ , dependent only on d and c , such that

$$(43) \quad U(y) > U(X_t) + 1 \text{ whenever } |y - x| < \delta \cdot d(X_t).$$

This implies that $\|U_{t+k} - U_t\|^2 > 1$ whenever $|X_{t+k} - x| < \delta d(X_t)$. Note that for some $p > 0$ dependent only on d and β ,

$$\mathbf{P}[|X_{t+k} - x| < (1 - \beta/2)^{-k} d(X_t)] > p^k.$$

Hence, for sufficiently large k , $\mathbf{P}[|X_{t+k} - x| < \delta d(X_t)] > p^k$, which, in turn, implies the statement of the lemma. \square

4. LOWER BOUNDS: EXAMPLES

In this section we construct examples of α -thick domains for which the bounds in Theorem 2 are tight. The main idea of the construction is as follows. We take a domain A in \mathbb{R}^d , such as the unit ball or a cylinder. We remove a “thin” subset of points C from A to obtain $\Omega = A \setminus C$. The set C can be thought of as the subset of the grid $(\gamma\mathbb{Z})^d$, for some small $\gamma > 0$. The set C will be chosen so that it “separates” the origin from the boundary of A . We set $n = 1/\varepsilon$. We choose γ so that the probability of the WoS originated at 0 hitting a $1/n$ -neighborhood of C before hitting the boundary of A is $< 1/2$ (this means that C is “thin”). Hence, with high probability, the WoS will reach ∂A before terminating. However, in this case the WoS will have to “pass through” the set C , where its step magnitudes are bounded by γ . This will, in turn, yield an $\Omega(1/\gamma^2)$ bound on the convergence time. The analysis is more intricate in the case when $\alpha = 2$. In the case when $\alpha > 2$ is not an integer, a slight modification to this construction is needed, as will be described below.

4.1. Proof of the lower bound in the case $\alpha > 2$. In this section we will give an example of a “thin” α -thick domain Ω_α for which the WoS will likely take $\Omega(n^{2-4/\alpha})$ steps to converge within $\varepsilon = 1/n$ from the boundary $\partial\Omega_\alpha$. The domain Ω_α will reside in \mathbb{R}^d , where $d = \lceil \alpha \rceil \geq 3$. It is easy to see that the examples in higher dimensions $d' > d$ can be constructed from Ω_α by simply multiplying Ω_α by $[-1, 1]^{d'-d}$.

The set

$$\Omega_\alpha := \left(B(0, 1)_{d-1} \times [-1, 1] \right) \setminus S$$

is comprised of a d -dimensional cylinder with a set of points S removed. Here $B(0, 1)_{d-1}$ denotes the unit ball in \mathbb{R}^{d-1} . We take A to be the “middle $1/3$ ” shell of the d -dimensional cylinder:

$$A = \{z \in \mathbb{R}^{d-1} : 1/3 < |z| < 2/3\} \times \{x \in [-1, 1] : 1/3 < |x| < 2/3\}.$$

Let $0 < \gamma \ll 1$ be the grid size that will be selected later. We consider the set A_γ of gridpoints in A .

$$A_\gamma = (\gamma\mathbb{Z})^d \cap A.$$

Let $0 \leq \eta := d - \alpha < 1$. Denote by C_η the η -dimensional Cantor set in the interval $[0, 1]$. It is obtained by removing the middle λ -fraction of the interval, then removing the middle λ -fraction of each subinterval etc. For the set C_η to be η -dimensional, we choose λ so that

$$\eta = \frac{\log 2}{\log 2 - \log(1 - \lambda)}.$$

In the special case when $\eta = 0$, we set $C_0 = \{0\}$. We now define the set S :

$$S := A_\gamma + \{0\} \times \gamma C_\eta.$$

In other words, S is obtained by attaching a γ -scaled copy of C_η to each gridpoint of A_γ . This completes the definition of the set $\Omega_\alpha = (B(0, 1)_{d-1} \times [-1, 1]) \setminus S$. Each point in $\partial\Omega_\alpha$ has an η -dimensional Cantor set in Ω_α^c attached to it is captured by the following claim. Thus there is a universal constant $C \geq 1/16$ such that for every γ , the set Ω_α is α -thick with the thickness C .

The following two claims assert that for an appropriately chosen γ , the WoS originated at the origin $0 \in \mathbb{R}^d$ and terminated at the $1/n$ neighborhood of $\partial\Omega_\alpha$ is likely to hit the boundary of the external cylinder (as opposed to the neighborhood of S), and is likely to spend $\Omega(n^{2-4/\alpha})$ steps getting there.

Claim 14. *If $\gamma > 8n^{2/\alpha-1}$ then a WoS originated at 0 and terminated at the $1/n$ -neighborhood of the boundary $\partial\Omega_\alpha$ will hit the boundary of the cylinder $B(0, 1)_{d-1} \times [-1, 1]$ with probability at least $3/4$.*

Proof. It is not hard to see that we can choose a finite subset P of points in S such that $|P| < 2\gamma^{-\alpha} \cdot n^\beta$, and for every x such that $d(x, S) < 1/n$ there is a $p \in P$ such that $|x - p| < 2/n$. Consider the harmonic function

$$(44) \quad \Phi(x) := \sum_{y \in P} \frac{1}{|x - y|^{d-2}} > 0.$$

Since the function Φ is harmonic, its application to the WoS process X_t gives a martingale. Hence if T is the stopping time of the process,

$$\mathbf{E}[\Phi(X_T)] = \Phi(X_0) = \Phi(0) < 3^{d-2} \cdot |P| < 6\gamma^{-\alpha} \cdot n^\eta.$$

On the other hand, if $d(X_T, S) < 1/n$, then there is a $y \in P$ with $|X_T - y| < 2/n$, and

$$\Phi(X_T) \geq 1/|y - X_T|^{d-2} > (n/2)^{d-2}.$$

Hence the probability of X_T being near S is bounded by

$$\frac{\mathbf{E}[\Phi(X_T)]}{(n/2)^{d-2}} < \frac{6\gamma^{-\alpha} \cdot n^\eta}{(n/2)^{d-2}} < \frac{2^{d+1}\gamma^{-\alpha}}{n^{\alpha-2}} < \frac{8^\alpha n^2}{4(\gamma n)^\alpha} < 1/4.$$

The last inequality follows from the condition on γ . □

Claim 15. *There is a universal constant $\delta > 0$ such that for γ as above, with probability at least $1/2$ the WoS takes at least $\delta(1/\gamma)^2$ steps to reach the boundary of the cylinder $B(0, 1)_{d-1} \times [-1, 1]$.*

Proof. The proof is done analogously to the proof of Claim 18 below. □

Hence the expected number of steps is at least

$$\frac{\delta}{2} \cdot \left(\frac{1}{8n^{2/\alpha-1}} \right)^2 = \Omega(n^{2-4/\alpha}),$$

which completes the proof of the lower bound for Theorem 2 in the case when $\alpha > 2$.

4.2. Proof of the lower bound in the case $\alpha = 2$. We will now give an example of a two dimensional domain Ω such that the expected convergence time of the WoS to a $O(1/n)$ -neighborhood of $\partial\Omega$ is $\Omega(\log^2 n)$. By taking the d -dimensional domain $\Omega_d = \Omega \times \mathbb{R}^{d-2}$ for $d > 2$, we obtain a lower bound of $\Omega(\log^2 n)$ for 2-thick domains in \mathbb{R}^d , proving the lower bound for $\alpha = 2$ in Theorem 2.

The domain Ω will consist of the unit disc in \mathbb{R}^2 with $O(\log n)$ holes “poked” out of it in a grid formation. More specifically, let $\gamma = 4/\log^{1/2} n$. We consider the grid $\Gamma = \gamma\mathbb{Z} \times \gamma\mathbb{Z} \subset \mathbb{R}^2$. We take Ω to be the unit disc with points from Γ removed from the “middle third” annulus of the disc.

$$(45) \quad \Omega = B(0, 1) \setminus ((B(0, 2/3) \setminus B(0, 1/3)) \cap \Gamma).$$

The set Ω is illustrated on Fig. 4(a).

We will show that a WoS originated at the origin $X_0 = 0$ would require an expected time of $\Omega(\log^2 n)$ to converge. It is immediate to see that the same lower bound holds for any point $X_0 \in B(0, 1/3)$. We first observe the following:

Claim 16. *With probability at least $7/8$, a WoS originated at $X_0 = 0$ that runs until $d(X_t) < 1/n$ terminates near the unit circle (and not near one of the holes).*

Proof. Let $\{a_i\}_{i=1}^k = B(0, 1) \setminus \Omega$ be the set of holes in Ω . Define the harmonic function

$$\Phi(z) = \sum_{i=1}^k \log(2/|z - a_i|).$$

It is clear the $\Phi(z) > 0$ for all $z \in B(0, 1)$. For any point u in the $1/n$ -neighborhood of any of the holes, $\Phi(u) > \log n$. On the other hand,

$$\Phi(0) < k \cdot \log 6 < 2/\gamma^2 = (\log n)/8.$$

If X_t is the WoS process with $X_0 = 0$ terminated at time T when $d(X_T, \partial\Omega) < 1/n$, then $\Phi(X_t)$ is a martingale. Hence,

$$(\log n)/8 > \Phi(X_0) = \mathbf{E}[\Phi(X_t)] > \mathbf{P}[X_t \text{ near a hole}] \cdot \log n.$$

Hence the probability that the WoS terminates near a hole is less than $1/8$. □

For simplicity, we will assume that at every step for the process the WoS jumps exactly half way to the boundary $\partial\Omega$.

To facilitate the analysis we replace the WoS process X_t on Ω with the following process Y_t . It evolves in exactly the same fashion as X_t , except when Y_t is closer than $1/n$ to one of the holes in Ω . In this case, instead of terminating, the process makes a jump of $1/n$ in a direction selected uniformly at random. The process Y_t is guaranteed to terminate near the unit circle. We denote the termination time by T . Further, we set $Y_t = Y_T$ for $t > T$. Note that if the process X_t does not terminate near one of the holes, then the process Y_t coincides with X_t . Claim 16 implies that this happens with probability at least $7/8$:

Claim 17. $\mathbf{P}[X_t \text{ does not coincide with } Y_t] < 1/8$.

We define two regions A and B , $B \subset A \subset \Omega$. We take A to be the union of discs with radius $r = \gamma/4$ around the holes in Ω . We take B to be the union of discs with radius $r/2$ around the same holes. The sets Ω , A and B are illustrated on Fig. 4(a).

Let time t_0 be the first time with $|Y_t| > 1/2$. Let t' be the first time afterward with either $|Y_t| > 2/3$ or $|Y_t| < 1/3$. Our goal is to show that with probability at least $3/4$, $|t_0 - t'| = \Omega(\log^2 n)$. We define a subprocess Z_t of Y_t as follows. Let $\{s_i\}_{i=0}^k$ be a subsequence of times s between t_0 and t' such that $Y_s \notin A$. We set $Z_i = Y_{s_i}$. We further define $\Delta_i = Z_i - Z_{i-1}$. An instance of the process




FIGURE 4. An illustration the sets Ω , A and B (a), and a possible sequence of jumps in the processes $\{Y_t\}$ and $\{Z_t\}$ (b)

Z_i is illustrated on Fig. 4(b). Since Y_t is a martingale, and Z_i is defined by a stopping rule on Y_t , Z_i is also a martingale, and

$$(46) \quad \mathbf{E}[\Delta_i \mid \Delta_1, \Delta_2, \dots, \Delta_{i-1}] = 0.$$

In addition, it is not hard to see from the definition of Y_t that $|\Delta_i| < 4/\log^{1/2} n$ for all i . Our first claim is that the number k of steps Z_i is $\Omega(\log n)$.

Claim 18. $\mathbf{P}[k < 10^{-4} \log n] < 1/8$.

Proof. Denote $\ell = 10^{-4} \log n$. Then, by (46),

$$\begin{aligned} \mathbf{E}[(Z_0 - Z_\ell)^2] &= \mathbf{E}[(\Delta_1 + \Delta_2 + \dots + \Delta_\ell)^2] = \sum_{j=1}^{\ell} \mathbf{E}[\Delta_j^2] + \sum_{1 \leq i < j \leq \ell} \mathbf{E}[\Delta_i \Delta_j] = \\ &= \sum_{j=1}^{\ell} \mathbf{E}[\Delta_j^2] + \sum_{1 \leq i < j \leq \ell} \mathbf{E}[\Delta_i \cdot \mathbf{E}[\Delta_j \mid \Delta_i]] = \sum_{j=1}^{\ell} \mathbf{E}[\Delta_j^2] < \ell \cdot 16/\log n < 1/288. \end{aligned}$$

On the other hand, by definition, $|Z_0 - Z_k| > 1/6$, and $(Z_0 - Z_k)^2 > 1/36$. Thus,

$$\mathbf{P}[k \leq \ell] = \mathbf{P}[Z_\ell = Z_k] < (1/288)/(1/36) = 1/8.$$

□

Thus the number of steps the process Z_t takes is at least $10^{-4} \log n$ w.p. $> 7/8$. The process Y_t consists of the steps of the process Z_t plus, in addition, steps the process takes within the region

A. We claim that once the process Y_t enters the region A , it is expected to spend $\Omega(\log n)$ steps there. Moreover, the following holds.

Claim 19. *Let $\eta > 2$. Then there is a $\theta > 0$ such that whenever $Y_t \in A$, if $s > t$ is the first time, conditioned on Y_t such that $Y_s \notin A$, then*

$$(47) \quad \mathbf{P}[s - t > \theta \log^2 n] > \eta / \log n,$$

for sufficiently large n .

Proof. Denote the hole in Ω that is closest to Y_t by x . Given that $Y_t \in A$, there is some fixed probability $p > 0$ that $Y_{t+1} \in B$. In other words, $|Y_{t+1} - x| < r/2 = \gamma/8$. Consider the harmonic function

$$\Phi(z) = \log(r/|x - z|).$$

Let $t' > t + 1$ be the first time such that either $Y_{t'} \notin A$ (and thus $t' = s$), or $|Y_{t'} - x| < n^{-p/(5\eta)}$. If $Y_{t'} \notin A$, then $\Phi(Y_{t'}) < 0$. In the other case, $\Phi(Y_{t'}) < p \log n / (4\eta)$. Since t' is a stopping time, the optional stopping time theorem applied to the martingale $\Phi(Y_{t+\tau})$ combined with the estimate $\Phi(Y_t) > 1/2$, gives

$$\mathbf{P}[|Y_{t'} - x| < n^{-p/(5\eta)}] > (1/2) / (p \log n / (4\eta)) > 2\eta / (p \log n).$$

To complete the argument, we claim that assuming $|Y_{t'} - x| < n^{-p/(5\alpha)}$, it will take the process another $\Omega(\log^2 n)$ steps to escape A with probability at least $1/2$. We consider the process $\phi_\tau = \Phi(Y_{t'+\tau})$ stopped at time τ_0 when either $Y_{t'+\tau_0}$ escapes A , or gets closer than distance $1/n$ from x . ϕ_τ is a martingale. Moreover, it is not hard to see that $|\phi_0 - \phi_{\tau_0}| > p \log n / (6\eta)$, and $|\phi_i - \phi_{i+1}| < 1$ for all i . These two facts imply that

$$\mathbf{E}[\tau_0] > \sum_{i=1}^{\tau_0} \mathbf{E}[(\phi_i - \phi_{i-1})^2] = \mathbf{E}[(\phi_{\tau_0} - \phi_0)^2] > (p \log n / (6\eta))^2 = p^2 \log^2 n / (36\eta^2).$$

Tschebyshev inequality implies that $\theta = p^2 / (72\eta^2)$ satisfies the statement of the claim. \square

By Claim 18 we know that except with probability $< 1/8$ the walk will contain at least $\Omega(\log n)$ visits to A . It remains to use Claim 19 to show that at least one of these stays must be $\Omega(\log^2 n)$ long. Recall that T is the stopping time of the process Y_T , and k is the number of steps Y_t takes outside of A .

Claim 20. *Let $\alpha_1 = 10^{-4}$ from Claim 18. There is a constant $\alpha_2 > 0$ such that*

$$(48) \quad \mathbf{P}[k > \alpha_1 \log n \text{ and } T < \alpha_2 \log^2 n] < 1/8.$$

Proof. For every t such that $1/3 < |Y_t| < 2/3$ and $Y_t \notin A$, there is a probability $p_1 > 0$ such that either $Y_{t+1} \in A$ or $Y_{t+2} \in A$. By Claim 19 we can choose $\alpha_2 > 0$ such that whenever $Y_{t'} \in A$, the process $Y_{t'+\tau}$ does not escape A for at least $\alpha_2 \log^2 n$ with probability at least $p_2 = 6 / (\alpha_1 p_1 \log n)$. Hence for each $1/3 < |Y_t| < 2/3$ with $Y_t \notin A$, the probability that Y_{t+1} or Y_{t+2} enters A , and stays there for at least $\alpha_2 \log^2 n$ steps is at least $p_1 \cdot p_2 = 6 / (\alpha_1 \log n)$. Since there are at least $k = \alpha_1 \log n$ Y_t 's satisfying $1/3 < |Y_t| < 2/3$ and $Y_t \notin A$, the probability that for neither one of them does Y_{t+1} or Y_{t+2} enter A , and stay there for at least $\alpha_2 \log^2 n$ steps is at most

$$(1 - 6 / (\alpha_2 \log n))^{k/2} < (1 - 6 / (\alpha_1 \log n))^{\alpha_1 \log n / 2} < e^{-(6 / (\alpha_1 \log n)) \cdot (\alpha_1 \log n / 2)} = e^{-3} < 1/8.$$

\square

Claims 17, 18 and 20 imply the following.

Claim 21. Let X_t be the WoS process on the set Ω with $X_0 = 0$. Let T' be its termination time. Then

$$\mathbf{P}[T' > \alpha_2 \log^2 n] > 5/8,$$

where $\alpha_2 > 0$ is the constant from Claim 20. In particular, this implies that $\mathbf{E}[T'] = \Omega(\log^2 n)$.

Proof. We know that $T' > \alpha_2 \log^2 n$ if the following three conditions hold: (C1) the process X_t coincides with the process Y_t ; (C2) the process Y_t makes at least $k > \alpha_1 \log n$ steps outside of A in the $\{z : 1/3 < |z| < 2/3\}$ annulus; and (C3) the stopping time T of Y_t satisfies $T > \alpha_2 \log^2 n$. In fact conditions (C1) and (C3) suffice. We have $\mathbf{P}[\overline{C1}] < 1/8$ by Claim 17, $\mathbf{P}[\overline{C2}] < 1/8$ by Claim 18, and $\mathbf{P}[C2 \cap \overline{C3}] < 1/8$ by Claim 20. Here \overline{C} denotes the complement of an event C . Hence

$$\mathbf{P}[\overline{C1} \cup \overline{C2} \cup \overline{C3}] \leq \mathbf{P}[\overline{C1}] + \mathbf{P}[\overline{C2}] + \mathbf{P}[C2 \cap \overline{C3}] < 3/8,$$

and

$$\mathbf{P}[T' > \alpha_2 \log^2 n] \geq \mathbf{P}[C1 \cap C2 \cap C3] = 1 - \mathbf{P}[\overline{C1} \cup \overline{C2} \cup \overline{C3}] > 5/8.$$

□

Claim 21 gives the lower bound for Theorem 2 in the case $\alpha = 2$.

REFERENCES

- [BB07] I. Binder and M. Braverman. Derandomization of Euclidean Random Walks. *LNCS*, 4627:353–365, 2007.
- [BC06] M. Braverman and S. Cook. Computing over the reals: Foundations for scientific computing. *Notices of the AMS*, 53(3):318–329, 2006.
- [BW99] V. Brattka and K. Weihrauch. Computability of subsets of Euclidean space I: Closed and compact subsets. *Theoretical Computer Science*, 219:65–93, 1999.
- [Car67] Lennart Carleson. *Selected problems on exceptional sets*. Van Nostrand Mathematical Studies, No. 13. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967.
- [EKM97] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling Extremal Events for Insurance and Finance*. Springer, 1997.
- [EKMS80] B. S. Elepov, A. A. Kronberg, G. A. Mihaïlov, and K. K. Sabel'fel'd. *Reshenie kraevykh zadach metodom Monte-Karlo*. “Nauka” Sibirsk. Otdel., Novosibirsk, 1980.
- [GM04] J. B. Garnett and D. E. Marshall. *Harmonic Measure*. Cambridge Univ Press, 2004.
- [Kak44] Shizuo Kakutani. Two-dimensional Brownian motion and harmonic functions. *Proc. Imp. Acad. Tokyo*, 20:706–714, 1944.
- [KS91] I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus*. Springer Verlag, 2 edition, 1991.
- [KS98] I. Karatzas and S.E. Shreve. *Methods of Mathematical Finance*. Springer, 1998.
- [Lan72] N. S. Landkof. *Foundations of modern potential theory*. Translated from the Russian by AP Doohovskoy, volume 180. 1972.
- [Maz02] R.M. Mazo. *Brownian Motion: Fluctuations, Dynamics, and Applications*. Oxford University Press, 2002.
- [Mih79] G. A. Mihaïlov. Estimation of the difficulty of simulating the process of “random walk on spheres” for some types of regions. *Zh. Vychisl. Mat. i Mat. Fiz.*, 19(2):510–515, 558–559, 1979.
- [Mil95] G. N. Milstein. *Numerical Integration of Stochastic Differential Equations*. Kluwer Academic Publishers, Dodrecht, 1995.
- [Mot59] Minoru Motoo. Some evaluations for continuous Monte Carlo method by using Brownian hitting process. *Ann. Inst. Statist. Math. Tokyo*, 11:49–54, 1959.
- [Mul56] M. E. Muller. Some continuous Monte Carlo methods for the Dirichlet problem. *Ann. Math. Statist.*, 27:569–589, 1956.
- [Nel67] E. Nelson. *Dynamical Theories of Brownian Motion*. Princeton University Press Princeton, NJ, 1967.
- [Szn98] A.S. Sznitman. *Brownian Motion, Obstacles and Random Media*. Springer, 1998.
- [Wei00] K. Weihrauch. *Computable Analysis*. Springer-Verlag, Berlin, 2000.

ILIA BINDER, DEPT. OF MATHEMATICS, UNIVERSITY OF TORONTO.

MARK BRAVERMAN, MICROSOFT RESEARCH, NEW ENGLAND