

Let  $z_2, z_3, z_4 \in \hat{\mathbb{C}}$  - three different points.

Theorem  $\exists!$  Möbius  $S: Sz_2=1, Sz_3=0, Sz_4=\infty$

Proof. If  $z_2 \neq \infty, z_3 \neq \infty, z_4 \neq \infty$ :

$$S(z) = \frac{z-z_3}{z-z_4} \cdot \frac{z_2-z_4}{z_2-z_3}$$

$$\begin{array}{lll} z_2 = \infty: & z_3 = \infty & z_4 = \infty \\ S(z) = \frac{z-z_3}{z-z_4} & S(z) = \frac{z_2-z_4}{z-z_4} & S(z) = \frac{z-z_3}{z_2-z_3} \end{array}$$

Uniqueness:  $S_1$  - another such transformation.

$$\text{Then } SS_1^{-1}(0)=0, SS_1^{-1}(1)=1, SS_1^{-1}(\infty)=\infty$$

$$\begin{array}{ccc} \Downarrow & & \Downarrow \\ b=0 & \implies & a=1 \iff c=\infty \end{array}$$

$$\Downarrow \\ SS_1^{-1}(z)=z$$

Def. The cross ratio  $(z_1, z_2, z_3, z_4) = S(z_1)$ , where  $S$  - Möbius map with  $Sz_2=1, Sz_3=0, Sz_4=\infty$ .

Theorem. Let  $T$  be a Möbius map. Then

(Cross ratio is invariant)  $(Tz_1, Tz_2, Tz_3, Tz_4) = (z_1, z_2, z_3, z_4)$  for any four distinct  $z_1, z_2, z_3, z_4 \in \hat{\mathbb{C}}$ .

Proof. Let  $Sz_2=1, Sz_3=0, Sz_4=\infty$ .

$$\text{Then } ST^{-1}(Tz_2)=1, ST^{-1}(Tz_3)=0, ST^{-1}(Tz_4)=\infty$$

$$\text{So } (Tz_1, Tz_2, Tz_3, Tz_4) = ST^{-1}(Tz_1) = Sz_1 = (z_1, z_2, z_3, z_4)$$

Theorem The cross ratio  $(z_1, z_2, z_3, z_4) \in \mathbb{R} \iff$

$z_1, z_2, z_3, z_4$  lie on the same circle or line.

Proof.  $(z_1, z_2, z_3, z_4) \in \mathbb{R} \iff (Sz_1, Sz_2, Sz_3, Sz_4) \in \mathbb{R} \iff (Sz_1, 1, 0, \infty) = Sz_1 \in \mathbb{R}$

$$\iff Sz_1 \text{ lies on the line generated by } \begin{array}{c} \text{Take the} \\ \text{right s} \end{array} \infty = Sz_4, 1 = Sz_3, 0 = Sz_2$$

lines and circles are Möbius invariant  $\iff z_1$  lies on the same line or circle as  $z_2, z_3, z_4$

## Symmetry.

Symmetry with respect to  $\mathbb{R}$ :  $z \mapsto \bar{z}$

$$(z, 1, 0, \infty) \mapsto (\bar{z}, 1, 0, \infty) = \overline{(z, 1, 0, \infty)}$$

Def. The points  $z$  and  $z^*$  are symmetric with respect to the line or the circle generated by  $z_1, z_2, z_3, z_4$  if  $(z, z_1, z_2, z_3, z_4) = \overline{(z^*, z_1, z_2, z_3, z_4)}$ .  
(or  $Sz = \overline{Sz^*}$ ).

Theorem. Does not depend on the choice of  $z_1, z_2, z_3, z_4$  on the same line or circle.

Symmetric wrt a line: the usual symmetry.

Symmetric wrt a circle  $|z-a|^2 = r^2$ :  $(z^*-a)(\bar{z}-a) = r^2$ .



$$\frac{|z^*-a|}{r} = \frac{r}{|z-a|}$$

$$\arg(z^*-a) = \arg(\bar{z}-a)$$

Proof. First, let us observe that

if  $z_1, z_2, z_3, z_4 \in \mathbb{R}$ , then  $z^* = \bar{z}$  (the map  $S$  has real coefficients,

$$\text{so } \frac{a\bar{z}+b}{c\bar{z}+d} = (\bar{z}, z_1, z_2, z_3, z_4) = \overline{\left(\frac{az+b}{cz+d}\right)} = \overline{(z, z_1, z_2, z_3, z_4)})$$

if  $z_1, z_2, z_3, z_4$  lie on the same line, then  $\exists Tz = az+b$ , such that  $Tz_1, Tz_2, Tz_3, Tz_4 \in \mathbb{R}$ .  $T$  preserves symmetry and crossratio.

Finally, if  $z_1, z_2, z_3, z_4 \in \{ |z-a|=r \}$  i.e.  $(z_i-a)(\bar{z}_i-a) = r^2$ , then

$$(\bar{z}, z_1, z_2, z_3, z_4) = \overline{(z-a, z_1-a, z_2-a, z_3-a, z_4-a)} = (\bar{z}-a, \bar{z}_1-a, \bar{z}_2-a, \bar{z}_3-a, \bar{z}_4-a)$$

$$(\bar{z}-a, \frac{r^2}{z_1-a}, \frac{r^2}{z_2-a}, \frac{r^2}{z_3-a}, \frac{r^2}{z_4-a}) = \left( \frac{r^2}{\bar{z}-a}, z_1-a, z_2-a, z_3-a, z_4-a \right) =$$

$$\left( \frac{r^2}{\bar{z}-a} + a, z_1, z_2, z_3, z_4 \right)$$

$z^*$

$$(z^*-a)(\bar{z}-a) = r^2 = (\bar{z}^*-a)(z-a)$$

Remark. With respect to  $\{ |z-a|=r \}$ ,  $a^* = \infty$

Proof Take  $z_2 = a+r$ ,  $z_3 = a-r$ ,  $z_4 = a+ir$

$$\text{Then } (\infty, z_1, z_2, z_3, z_4) = \frac{z_2-z_4}{z_2-z_3} = \frac{r-ir}{2r} = \frac{1-i}{2}$$

$$(a, z_1, z_2, z_3, z_4) = \frac{z_1-z_3}{a-z_4} \cdot \frac{z_2-z_4}{z_2-z_3} = \frac{r}{-ir} \cdot \frac{1-i}{2} = i \cdot \frac{1-i}{2} = \frac{1+i}{2} = \overline{(\infty, z_1, z_2, z_3, z_4)} =$$

Corollary  $T$ -Möbius,  $z, z^*$  - symmetric with respect

a circle or line  $\ell \Rightarrow Tz, |z|^2$ -symmetric wrt  
circle or line  $T\ell$ .

Proof.  $z_1, z_2, z_3, z_4 \in \ell$   $(z_1, z_2, z_3, z_4) \stackrel{\text{sym}}{=} (\overline{z_1}, \overline{z_2}, \overline{z_3}, \overline{z_4})$   
 $(Tz_1, Tz_2, Tz_3, Tz_4) = (T\overline{z_1}, T\overline{z_2}, T\overline{z_3}, T\overline{z_4})$

Theorem.  $T$ -Möbius,  $T(\mathbb{D}) = \mathbb{D}$  ( $\mathbb{D} = \{z, |z| < 1\}$ ).

Then  $Tz = e^{i\theta} \frac{z-a}{1-\overline{a}z}$ , for some  $a \in \mathbb{D}$ ,  $\theta \in \mathbb{R}$ .

Proof. First, for  $|z| = 1$ ,  $z = e^{i\theta}$

$$|Tz| = |e^{i\theta}| \frac{|z-a|}{|1-\overline{a}z|} = \frac{|z-a|}{|z||z-\overline{a}|} = 1. \text{ So the circle is preserved.}$$

So  $\mathbb{D}$  is mapped either to itself, or to  $\mathbb{D}_- = \{z, |z| > 1\}$ .

But  $a \rightarrow 0$ ,  $a \in \mathbb{D}$ , so  $T\mathbb{D} = \mathbb{D}$ .

Let  $T(\mathbb{D}) = \mathbb{D}$ . Then let  $a = T^{-1}0 \in \mathbb{D}$ .

Then  $T(a^*) = T(1/\overline{a}) = 0^* = \infty$ .

so  $T(z) = c \frac{z-a}{z - \frac{1}{\overline{a}}} = \underbrace{-c\overline{a}}_d \frac{z-a}{1-\overline{a}z} = d \frac{z-a}{1-\overline{a}z}$ .

But  $|T1|=1$ , so  $|d| \frac{|1-a|}{|1-\overline{a}|} = 1 \Rightarrow |d|=1 \Rightarrow d = e^{i\theta}$

Theorem  $H = \{Im z > 0\}$ .  $T(H) = H \Leftrightarrow T = \frac{az+b}{cz+d}$   $a, b, c, d \in \mathbb{R}$ .

Proof. Very similar, left as exercise.

$ad - bc > 0$

$Im \frac{ai+b}{ci+d} > 0$