

**MAT 347**  
**Quotient groups**  
**October 7, 2015**

Let  $G$  be a group and let  $S \subseteq G$ . We want to define an equivalence relation in  $G$  that will identify all the elements in  $S$ , and we want to maintain the group operation. Given  $a, b \in G$ , we say that  $a \sim b$  when there exists  $x \in S$  such that  $b = ax$ . In general, this relation will not be an equivalence relation.

1. Find necessary and sufficient conditions for  $\sim$  to be an equivalence relation.

For the rest of this worksheet, let  $G$  be a group and let  $H \leq G$ . We will consider the equivalence relation defined above with  $S = H$ . Given  $a \in G$ , the *left coset* of  $a$  is the equivalence class of this relation, and we denote it  $aH$ . (Why do we use this notation?) The *quotient set*  $G/H$  is the set of all equivalence classes. The *index* of  $H$  on  $G$ , written  $|G : H|$  is the number of equivalence classes (when this number is finite).

2. Prove that  $aH = bH$  iff [there exists  $x \in H$  such that  $b = ax$ ] iff  $a^{-1}b \in H$
3. If  $H$  is finite, what is the cardinality of each coset  $aH$ ? If both  $G$  and  $H$  are finite, what is the relation between  $|G|$ ,  $|H|$ , and  $|G : H|$ ?
4. Consider the group  $G = D_8$  and consider the two subgroups  $H_1 := \langle s \rangle$  and  $H_2 := \langle r^2 \rangle$ . For each of them, write the complete list of cosets, and list which elements are in each coset.

Next, we want to try to use the operation on  $G$  to define an operation on the set  $G/H$ . Given  $aH, bH \in G/H$ , we can try to define their product by

$$(aH) \star (bH) = (ab)H$$

[*Note:* I am using  $\star$  to emphasize that I am defining a new operation. As soon as we make sure this operation works and there is no ambiguity, we will drop the  $\star$ .]

5. In general, the operation  $\star$  is not well-defined. Going back to the example in Question 4, show that with one of those subgroups, the operation is well-defined, but with the other subgroup, the operation is not well-defined.

**Definition:** We say that the subgroup  $H$  is a *normal subgroup* of  $G$  when the operation  $\star$  in  $G/H$  is well-defined. We write it  $H \trianglelefteq G$

6. Assume  $H \trianglelefteq G$ . In this case we know the operation in  $G/H$  is well-defined. What other conditions do we need to impose so that  $G/H$  is a group with this operation?

## The big theorem about normal subgroups

**Notation:** Let  $G$  be a group. Given subsets  $A, B$  and elements  $x, y$  we will use the following notation:

$$\begin{aligned}xA &:= \{xa \mid a \in A\} \\xAy &:= \{xay \mid a \in A\} \\AB &:= \{ab \mid a \in A, b \in B\}\end{aligned}$$

7. Let  $G$  be a group and let  $H \subseteq G$ . Explore the relation between the following statements (which ones imply which ones)?
- (a)  $H \trianglelefteq G$
  - (b)  $aH = Ha$  for all  $a \in G$   
(Notice that this does not mean that  $a$  commutes with the elements of  $H$ . It only means that the sets  $aH$  and  $Ha$  are the same set.)
  - (c)  $aHa^{-1} = H$  for all  $a \in G$
  - (d)  $aHa^{-1} \subseteq H$  for all  $a \in G$
  - (e) There exists some group  $L$  and some group homomorphism  $f : G \rightarrow L$  such that  $H = \ker f$ .
8. Find all normal subgroups of  $D_8$ .