

**International Mathematics
TOURNAMENT OF THE TOWNS**

Senior A-Level Paper

Fall 2007.

1. (a) Each of Peter and Basil thinks of three positive integers. For each pair of his numbers, Peter writes down the greatest common divisor of the two numbers. For each pair of his numbers, Basil writes down the least common multiple of the two numbers. If both Peter and Basil write down the same three numbers, prove that these three numbers are equal to each other.
(b) Can the analogous result be proved if each of Peter and Basil thinks of four positive integers instead?
2. Let K , L , M and N be the midpoints of the sides AB , BC , CD and DA of a cyclic quadrilateral $ABCD$. Let P be the point of intersection of AC and BD . Prove that the circumradii of triangles PKL , PLM , PMN and PNK are equal to one another.
3. Determine all finite increasing arithmetic progressions in which each term is the reciprocal of a positive integer and the sum of all the terms is 1.
4. Attached to each of a number of objects is a tag which states the correct mass of the object. The tags have fallen off and have been replaced on the objects at random. We wish to determine if by chance all tags are in fact correct. We may use exactly once a horizontal lever which is supported at its middle. The objects can be hung from the lever at any point on either side of the support. The lever either stays horizontal or tilts to one side. Is this task always possible?
5. The audience arranges n coins in a row. The sequence of heads and tails is chosen arbitrarily. The audience also chooses a number between 1 and n inclusive. Then the assistant turns one of the coins over, and the magician is brought in to examine the resulting sequence. By an agreement with the assistant beforehand, the magician tries to determine the number chosen by the audience.
 - (a) Prove that if this is possible for some n_1 and n_2 , then it is also possible for n_1n_2 .
 - (b) Determine all n for which this is possible.
6. Let P and Q be two convex polygons. Let h be the length of the projection of Q onto a line perpendicular to a side of P which is of length p . Define $f(P, Q)$ to be the sum of the products hp over all sides of P . Prove that $f(P, Q) = f(Q, P)$.
7. There are 100 boxes, each containing either a red cube or a blue cube. Alex has a sum of money initially, and places bets on the colour of the cube in each box in turn. The bet can be anywhere from 0 up to everything he has at the time. After the bet has been placed, the box is opened. If Alex loses, his bet will be taken away. If he wins, he will get his bet back, plus a sum equal to the bet. Then he moves onto the next box, until he has bet on the last one, or until he runs out of money. What is the maximum factor by which he can guarantee to increase his amount of money, if he knows that the exact number of blue cubes is
 - (a) 1;
 - (b) some integer k , $1 < k \leq 100$.

Note: The problems are worth 2+2, 6, 6, 6, 4+4, 8 and 3+5 points respectively.