

PROBLEMS (due Feb 29)

1. Let $\tau_x = \inf\{t \geq 0 : B_t \geq x\}$. Prove that τ_x is equal in distribution to $x^2(\sup_{0 \leq s \leq 1} B_s)^{-2}$.
2. Let B_t^1 and B_t^2 be independent Brownian motions. Find the one dimensional marginal distribution of $\frac{B_t^1}{|B_t^2|}$ (the one dimensional marginal distribution of X_t means $F(t) = P(X_t \leq x)$, $t \geq 0$).
3. Let $B_t = (B_t^1, B_t^2)$ be a two dimensional Brownian motion starting at $(0, 0)$. Let τ be the first time that B_t hits the line $\{(x, y) : y = a\}$. Compute the distribution of B_τ (Hint: Use problems 1 and 2.)
4. Let $\tau_{\pm x} = \inf\{t \geq 0 : |B_t| = x\}$. Compute the Laplace transform of $\tau_{\pm x}$.
5. Let G be a nice subset of the lattice \mathbf{Z}^d and ∂G be its boundary. Let f be a function defined on the boundary. Let X_n be a symmetric nearest neighbour random walk on the lattice starting at $x \in G$ and let τ_G be the first time that the walk hits the boundary of G . Show that the solution of the discrete Dirichlet problem

$$\Delta u = 0 \quad \text{in } G, \quad u = f \quad \text{on } \partial G$$

is given by $u(x) = E_x[f(X_{\tau_G})]$. The lattice Laplacian $\Delta u(x)$ is given by $\sum_{y \sim x} u(y) - u(x)$ where $y \sim x$ means y is a neighbour of x . (Note that the wording of the problem is purposefully vague and part of the problem is to make words like “subset”, “nice” and “boundary” precise here.)

6. Let \mathcal{O} be an orthogonal transformation of $L^2[0, \infty)$. Let $\mathbf{1}_{[0,t]}$ be the indicator function of $0 \leq s \leq t$. Define

$$\tilde{B}(t) = \int_0^\infty (\mathcal{O}\mathbf{1}_{[0,t]})(s) dB(s).$$

Show that $B \mapsto \tilde{B}$ is a measure preserving transformation of the space of Brownian paths.

7. Let B_t , $t \geq 0$ be a d -dimensional Brownian motion. Show that the one dimensional marginal densities $f(t, x) = \lim_{\epsilon \rightarrow 0} \frac{P(|B_t - x| \leq \epsilon)}{\text{Volume of } \{|x| \leq \epsilon\}}$ satisfies the heat equation $\partial_t f = \frac{1}{2} \sum_{i=1}^d \frac{\partial^2 f}{\partial x_i^2}$. Find the equation satisfied by the one dimensional marginals of the *Bessel process* $r_t = |B_t|$.