

MAT 257Y Term Test 2 Ptactice Test 1

- (1) Let $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be C^1 where $n > m$. Suppose $[df(x_0)]$ has rank m .

Show that there exists $\epsilon > 0$ such that for any $y \in B(f(x_0), \epsilon)$ there exists $x \in \mathbb{R}^n$ such that $f(x) = y$.

- (2) Let A be a rectangle in \mathbb{R}^n and let $S \subset A$ be a set of measure zero which is rectifiable. Show that S has content zero.

Hint: Use that $\int_A \chi_S$ exists and must be equal to zero.

- (3) Let $f: [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be continuous.

Show that

$$\int_0^1 \left(\int_0^x f(x, y) dy \right) dx = \int_0^1 \left(\int_y^1 f(x, y) dx \right) dy$$

- (4) Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^2 .

Prove that $F(x) = \int_0^1 f(x, y) dy$ is C^2 on \mathbb{R} .

- (5) Prove that the union of countably many sets of measure zero has measure 0.

- (6) let $S = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1, y \geq |x|\}$.

Compute $\int_S y$.

- (7) Let $A \subset \mathbb{R}^n, B \subset \mathbb{R}^m$ be rectangles. let $f: A \times B \rightarrow \mathbb{R}$ be integrable.

Prove that there is a set $S \subset A$ of measure 0 such that for any $x \in A \setminus S$ the integral $\int_B f(x, y) dy$ exists.

- (8) Let $f: [-1, 1] \times [-1, 1] \rightarrow \mathbb{R}$ be a continuous function. Suppose $f(-x, y) = -f(x, y)$ for any x, y .

Prove that $\int_{[-1,1] \times [-1,1]} f = 0$.

Hint: use Fubini's theorem.