

PERELMAN'S STABILITY THEOREM

VITALI KAPOVITCH

ABSTRACT. We give a proof of the celebrated stability theorem of Perelman stating that for a noncollapsing sequence X_i of Alexandrov spaces with $\text{curv} \geq k$ Gromov-Hausdorff converging to a compact Alexandrov space X , X_i is homeomorphic to X for all large i .

1. INTRODUCTION

A fundamental observation of Gromov says that the class of complete n -dimensional Riemannian manifolds with fixed lower curvature and upper diameter bounds is precompact in Gromov-Hausdorff topology. The limit points of this class are Alexandrov spaces of dimension $\leq n$ with the same lower curvature and upper diameter bounds. Given a sequence of manifolds M_i in the above class converging to an Alexandrov space X it's interesting to know what can be said about the relationship between topology of the limit and elements of the sequence.

The main purpose of this paper is to give a careful proof of the following Theorem of Perelman which answers this question in the situation when $\dim X = n$.

Stability Theorem 1.1. *Let X^n be a compact n -dimensional Alexandrov space of $\text{curv} \geq \kappa$. Then there exists an $\epsilon = \epsilon(X) > 0$ such that for any n -dimensional Alexandrov space Y^n of $\text{curv} \geq \kappa$ with $d_{G-H}(X, Y) < \epsilon$, Y is homeomorphic to X .*

To prove the Stability Theorem, it's clearly enough to show that if X_i^n is a sequence of compact n -dimensional Alexandrov spaces with $\text{curv} \geq \kappa$, $\text{diam} \leq D$ converging in Gromov-Hausdorff topology to a space X of dimension n , then Hausdorff approximations $X_i \rightarrow X$ can be approximated by homeomorphisms for all large i . It is this statement that will be proved in the present paper.

A proof of the Stability Theorem was given in [Per91]. However, that paper is very hard to read and is not easily accessible. We aim to provide a comprehensive and hopefully readable reference for Perelman's result.

Perelman also claims to have a proof of the Lipschitz version of the Stability Theorem which says that the stability homeomorphisms can be chosen to be bi-Lipschitz. However, the proof of that result has never been written down and the author has never seen it (although he very much wants to).

It is also worth pointing out that the Stability Theorem in dimension 3 plays a key role in the classification of collapsing of 3-manifolds with a lower curvature bound by Shioya and Yamaguchi [SY00, SY05] which in turn plays a role in Perelman's work on the geometrization

2000 AMS Mathematics Subject Classification: 53C20. Keywords: nonnegative curvature, nilpotent.
The author was supported in part by an NSF grant.

conjecture. However, as was communicated to the author by Kleiner, for that particular application, if one traces through the proofs of [SY00, SY05] carrying along the additional bounds arising from the Ricci flow, then one finds that in each instance when a 3-dimensional Alexandrov space arises as a Gromov-Hausdorff limit of smooth manifolds, it is in fact smooth, and after passing to an appropriate subsequence, the convergence will also be smooth to a large order. For such convergence the stability theorem is very well known and easily follows from Cheeger-Gromov compactness. Therefore Perelman's stability theorem is unnecessary for the application to geometrization.

The Stability Theorem immediately implies the following finiteness theorem due to Grove-Petersen-Wu which was originally proved using controlled homotopy theory techniques.

Theorem 1.2. [GPW91] *The class of n -dimensional Riemannian manifolds ($n \neq 3$) with sectional curvature $\geq k$, diameter $\leq D$ and volume $\geq v$ has only finitely many topological (differentiable if $n \neq 3, 4$) types of manifolds.*

The restriction $n \neq 3$ in this theorem comes from the use of controlled homotopy results which do not hold in dimension 3. Using the Stability theorem one gets that homeomorphism finiteness holds in all dimensions including dimension 3. Alternatively, the homeomorphism finiteness in dimension 3 follows from an earlier result of Grove-Petersen [GP88] that the above class has finitely many homotopy types and the fact that in dimension 3, a fixed homotopy type of closed manifolds contains only finitely many homeomorphism types (this a direct consequence of the geometrization conjecture).

Let us mention that the proof of Stability Theorem 1.1 presented here is fundamentally the same as the one given in [Per91]. However, as was pointed out by Perelman in [Per93], the proof can be simplified using the constructions developed in [Per93] and [Per95]. We carry out these simplifications in the present paper.

In [Per95] Perelman introduced the following notion:

A function $f: \mathbb{R}^k \rightarrow \mathbb{R}$ is called DC if it can be locally represented as a difference of two concave (or, equivalently, semiconcave) functions. It is easy to see that DC is an algebra and $\frac{f}{g}$ is DC if both f and g are DC and g is never zero. A map $F: \mathbb{R}^k \rightarrow \mathbb{R}^l$ is called DC if it has DC coordinates. It is also easy to see (see [Per95]) that if $F: \mathbb{R}^k \rightarrow \mathbb{R}^l$ and $G: \mathbb{R}^l \rightarrow \mathbb{R}^m$ are DC then so is $G \circ F$. This allows for an obvious definition of a DC geometric structure on a topological manifold. Despite the ease and naturality of its definition, to the best of the author's knowledge, this type of geometric structure has never been studied. In particular, the relationship between DC structures and classical geometric structures such as TOP, PL, smooth or Lipschitz is not at all understood. It is easy to see that a PL-manifold is DC and a DC manifold is Lipschitz but that's all one can say at the moment.

The notion of DC functions and DC homeomorphisms also makes sense on Alexandrov spaces since Alexandrov spaces admit an abundance of semiconcave (and hence DC) functions coming from distance functions. In [Per95], Perelman showed that the set of regular points of an Alexandrov space has a natural structure of a DC manifold.

The above discussion naturally leads to the following question.

Question 1.3. *Let X_i^n be a noncollapsing sequence of Alexandrov spaces with $\text{curv} \geq \kappa$, $\text{diam} \leq D$ Gromov-Hausdorff converging to an Alexandrov space X^n . Is it true that X_i is DC-homeomorphic to X for all large i ?*

Or more weakly. Suppose M_i^n is a noncollapsing sequence of Riemannian manifolds with $\text{sec} \geq \kappa$, $\text{diam} \leq D$ Gromov-Hausdorff converging to an Alexandrov space X^n .

Is it true that M_i are DC homeomorphic to each other for all large i ?

Let us say a few words about the proof of the stability theorem. One of the main ingredients is the Morse theory for functions on Alexandrov spaces. The starting point is based on the following simple observation:

Given $k + 1$ nonzero vectors in \mathbb{R}^n with pairwise angles bigger than $\pi/2$, any k of them are linearly independent.

Motivated by this, we'll say that a map $f = (f_1, \dots, f_k)$ (with coordinates given by distance functions $f_i = d(\cdot, a_i)$) from an Alexandrov space to \mathbb{R}^k is regular at a point p if there exists a point a_0 such that the comparison angles at p for the triangles $\Delta a_i p a_j$ are bigger than $\pi/2$ for all $0 \leq i \neq j \leq k$.

The fibration theorem [Per91, Per93] shows that just like for regular points of smooth functions on differentiable manifolds, a map from an Alexandrov space X to \mathbb{R}^k is a topological submersion on the set of its regular points. In particular, if $k = \dim X$ then it's a local homeomorphism.

It's well known that for any point p in an Alexandrov space, $d(\cdot, p)$ has no critical points in a sufficiently small punctured ball $B(p, \epsilon) \setminus \{p\}$. Therefore, by the fibration theorem, $B(p, \epsilon)$ is homeomorphic to the cone over the metric sphere $S(p, \epsilon)$.

The fibration theorem (and, perhaps, even more importantly, its proof) plays a key role in the proof of the stability theorem. In particular it implies that Alexandrov spaces are stratified topological manifolds. The k -dimensional strata of X is basically equal to the set of points p in X such there exist $k + 1$ (but not $k + 2$) vectors in $T_p X$ with pairwise angles bigger than $\pi/2$.

Another important tool in the proof of stability is the gluing theorem derived from deformations of homeomorphisms results of Siebenmann [Sie72]. Given a noncollapsing sequence of Alexandrov spaces $X_i \rightarrow X$, it says that one can glue Hausdorff close stability homeomorphisms defined on a fixed open cover of the limit space X to make a global nearby homeomorphism, i.e it reduces the stability theorem to the local situation.

To construct local stability homeomorphisms near a point $p \in X$ one argues by reverse induction on the dimension k of the strata passing through p . The base of induction follows from the fact mentioned above that a map $f: X^n \rightarrow \mathbb{R}^n$ is a local homeomorphism near a regular point. The general induction step is quite involved and can not be readily described in an introduction. To show some of its flavor, we will however try to say a few words about the last step of induction from $k = 1$ to $k = 0$.

Given a point p in the zero strata (i.e. with the diameter of the space of directions at p at most $\pi/2$), look at a ball $B(p, r)$ where again r is so small that $d(\cdot, p)$ is regular in $B(p, r) \setminus \{p\}$. One can show that the same holds true for appropriately chosen lifts $p_i \in X_i$ of p for all large i provided r is sufficiently small. This means that both $B(p, r) \setminus \{p\}$ and $B(p_i, r) \setminus \{p_i\}$ consist of points lying in the union of strata of dimensions ≥ 1 .

Fixing a small $\delta \ll r$, the induction assumption implies the existence of homeomorphisms of the annuli $\bar{B}(p, r) \setminus B(p, \delta)$ and $\bar{B}(p_i, r) \setminus B(p_i, \delta)$ close to the original Hausdorff approximations. In particular, metric spheres $S(p, \delta)$ and $S(p_i, \delta)$ are homeomorphic. On the

other hand, the fibration theorem implies that $B(p, \delta)$ is homeomorphic to the cone over $S(p, \delta)$ and $B(p_i, \delta)$ is homeomorphic to the cone over $S(p_i, \delta)$ for all large i . Gluing these homeomorphisms we obtain homeomorphisms $B(p, r)$ onto $B(p_i, r)$ which will be close to the original Hausdorff approximations since $\delta \ll r$.

The general induction step is a rather nontrivial fibered version of the above argument. It is carried out in Local Stability Lemma 7.9. with the main geometric ingredient provided by Lemma 6.9. For technical reasons, one has to work with more general semiconcave functions than just distance functions. An important role here is played by a technical construction from [Per93] of strictly concave functions obtained by manipulating distance functions.

Let us briefly describe the structure of this paper.

In section 3 we give a simplified proof of the stability theorem in the special case of limits of closed Riemannian manifolds using controlled homotopy theory techniques. In section 4 we give the background on necessary topological results on stratified spaces and some geometric constructions on Alexandrov spaces. In sections 5 and 6 we define admissible maps and their regular points and show that they satisfy similar properties to regular points of smooth maps between manifolds. Section 7 contains the proof of the stability theorem. In section 8 we use the Stability Theorem to prove a finiteness theorem for Riemannian submersions. In section 9 we generalize the Stability Theorem to show that stability homeomorphisms can be chosen to respect stratification of Alexandrov spaces into extremal subsets.

Throughout this paper we will assume some knowledge of Alexandrov geometry (see [BGP92] as a basic reference). For a more recent treatment we strongly recommend [Pet06] in the same volume. We will also rely on the Morse theory results and constructions from [Per93] in particular the fibration theorem. Perelman later generalized these results to a larger class of Morse functions in the unpublished preprint [Per95] (cf. also [Pet06]). However, these generalizations are not needed for the proof of the Stability theorem presented here. Various references to [Per95] throughout this paper are only given for bibliographical completeness.

Acknowledgements: The author is profoundly grateful to Anton Petrunin and Fred Wilhelm for numerous conversations and suggestions regarding the preparation of this paper. The author would like to thank Alexander Lytchak for discussions and suggestions regarding the controlled homotopy theory proof of the stability theorem for manifolds.

2. NOTATIONS

For an open subset U in a space X and a subset $A \subset U$ we'll write $A \Subset U$ if $\bar{A} \subset U$.

We'll denote by $\mathbf{Alex}^n(D, \kappa, v)$ the class of n -dimensional Alexandrov spaces of curv $\geq \kappa$, diam $\leq D$, vol $\geq v$. Similarly, we'll denote by $\mathbf{Alex}^n(D, \kappa)$ the class of n -dimensional Alexandrov spaces of curv $\geq \kappa$, diam $\leq D$ and by $\mathbf{Alex}^n(\kappa)$ the class of n -dimensional Alexandrov spaces of curv $\geq \kappa$.

For a point $v = (v_1, \dots, v_k) \in \mathbb{R}^k$ and $r > 0$ we'll denote by $I_r^k(v)$ the cube $\{x \in \mathbb{R}^k \mid |x_i - v_i| \leq r \text{ for all } i\}$.

For an Alexandrov space Σ of curv ≥ 1 we'll often refer to its points as vectors and to distances between its points as angles.

For a metric space X with diam $\leq \pi$, we'll denote by CX the Euclidean cone over X . Let o be the vertex of CX . For $u \in CX$ we'll denote $|u| = d(o, u)$.

We'll call a function $h: CX \rightarrow \mathbb{R}$ 1-homogeneous if $h(t \cdot x) = t \cdot h(x)$ for any $t \in \mathbb{R}, x \in X$.

For any space X we'll denote by KX the open cone on X and by $\bar{K}X$ the closed cone on X (i.e. the join of X and a point).

For two points p, q in an Alexandrov space, we'll denote by \uparrow_p^q an element of $\Sigma_p X$ tangent to a shortest geodesic connecting p to q . We'll denote by \uparrow_p^q the set of *all* such directions.

For three points x, p, y in an Alexandrov space X of $\text{curv} \geq \kappa$, we'll denote by $\tilde{\angle} xpy$ the comparison angle at p , i.e the angle $\angle \tilde{x}\tilde{p}\tilde{y}$ in the triangle $\tilde{x}\tilde{p}\tilde{y}$ in the complete simply connected space of constant curvature κ with $d(x, y) = d(\tilde{x}, \tilde{y}), d(p, y) = d(\tilde{p}, \tilde{y}), d(x, p) = d(\tilde{x}, \tilde{p})$,

We will also often use the following convention. In the proofs of various theorems we'll denote by c or C various constants depending on the dimension and the lower curvature bound present and which sometimes will depend on additional parameters present. When important this dependence will be clearly indicated.

We will denote by \varkappa various continuous increasing functions $\varkappa: R_+ \rightarrow \mathbb{R}_+$ satisfying $\varkappa(0) = 0$.

By $o(i)$ we will denote various positive functions on \mathbb{Z}_+ such that $o(i) \xrightarrow{i \rightarrow \infty} 0$.

We'll write $o(i|c)$ to indicate a function which depends on an extra parameter c and satisfies $o(i|c) \xrightarrow{i \rightarrow \infty} 0$ for any fixed c . Sometimes we'll use the same convention for $\varkappa(\delta|c)$.

3. SIMPLIFIED PROOF OF THE STABILITY THEOREM FOR LIMITS OF MANIFOLDS

The author is grateful to A. Lytchak for bringing to his attention the fact, that using today's knowledge of the local structure of Alexandrov spaces, a simple proof of the stability theorem for $n \geq 4$ can be given for the special case of limits of Riemannian manifolds.

The proof uses controlled homotopy theory techniques employed in [GPW91].

Let us briefly describe the argument the general outline of which was suggested to the author by A. Lytchak.

It is now well-known that the class $\mathfrak{Aler}^n(D, \kappa, v)$ has a common contractibility function (see e.g. [PP93] or [Per93]). Therefore, by [GPW91, Lemma 1.3], if $X_i^n \xrightarrow{i \rightarrow \infty} X^n$ is a convergent sequence in $\mathfrak{Aler}^n(D, \kappa, v)$, then X_i^n is $o(i)$ -homotopy equivalent to X for all large i (see [GPW91]). This means that there are homotopy equivalences $f_i: X_i \rightarrow X$ with homotopy inverses $h_i: X \rightarrow X_i$ such that $f_i \circ h_i \simeq \text{Id}_X$ and $h_i \circ f_i \simeq \text{Id}_{X_i}$ through homotopies $F_i: X \times [0, 1] \rightarrow X$ and $G_i: X_i \times [0, 1] \rightarrow X_i$ such that all the point trajectories of F_i and $f_i \circ G_i$ have $o(i)$ -small diameters in X . In fact, in our case, one can make f_i, h_i to be $o(i)$ -Hausdorff approximations.

It is relatively easy to show (see [GPW91]) that if $M_i \in \mathfrak{Aler}^n(D, \kappa, v)$ is a sequence of closed Riemannian manifolds Gromov-Hausdorff converging to a space X , then X is a homology manifold.

However, at the time of the writing of [GPW91] the local structure of Alexandrov spaces was not well understood and it was therefore not known if X is an actual manifold. This made the application of controlled topology techniques employed in [GPW91] fairly tricky.

We first show that X is a manifold.

We'll need the following result from [Kap02] (cf. [GW97]).

Lemma 3.1. *Suppose $M_i \in \mathfrak{Aler}^n(D, \kappa, v)$ is a sequence of Riemannian manifolds Gromov-Hausdorff converging to a space X . Let $p \in X$ be any point.*

Then there exists a $\delta > 0$, a 1-Lipschitz function $h: X \rightarrow \mathbb{R}$, strictly convex on $B(p, \delta)$ such that $h(p) = 0$ is a strict local minimum of h and a sequence of smooth 1-Lipschitz functions $h_i: M_i \rightarrow \mathbb{R}$ uniformly converging to h such that h_i is strictly convex on $B(p_i, \delta)$ (where $p_i \in M_i$ converges to p) for all large i .

This lemma together with the fibration theorem of Perelman [Per93] easily implies that a noncollapsing limit of Riemannian manifolds with lower sectional curvature bound is a topological manifold.

Lemma 3.2. *Let $M_i \in \mathfrak{Aler}^n(D, \kappa, v)$ be a sequence of closed Riemannian manifolds converging to an Alexandrov space X . Then X is a topological manifold.*

Proof. We argue by induction on dimension. The cases of $n \leq 2$ are easy and are left to the reader as an exercise.

Suppose $n \geq 3$. Let $p \in X$ be any point and let h be the function provided by Lemma 3.1. Then for some small $\epsilon > 0$ the set $\{h \leq \epsilon\}$ is a compact convex subset of X and $\{h_i \leq \epsilon\}$ is compact convex in M_i for large i . Obviously, $\{h_i \leq \epsilon\} \xrightarrow[i \rightarrow \infty]{G-H} \{h \leq \epsilon\}$. By [Pet97, Theorem 1.2], we have that $\{h_i = \epsilon\} \xrightarrow[i \rightarrow \infty]{G-H} \{h = \epsilon\}$ with respect to their induced inner metrics. By the Gauss Formula, $\text{sec}(\{h_i = \epsilon\}) \geq \kappa$ for all large i and hence $\{h = \epsilon\}$ is an $(n-1)$ -dimensional Alexandrov space of $\text{curv} \geq \kappa$ which is a manifold by induction assumption. Moreover, since h_i is smooth, strictly convex with unique minimum, $\{h_i = \epsilon\}$ is diffeomorphic to S^{n-1} for large i and hence $\{h = \epsilon\}$ is a homotopy $(n-1)$ -sphere. Since h is strictly convex in $\{h \leq \epsilon\}$, it has no critical points in $\{h \leq \epsilon\} \setminus \{p\}$. Therefore, by [Per93, Theorem 1.4], $\{h < \epsilon\}$ is homeomorphic to the open cone over $\{h = \epsilon\}$.

If $n = 3$ then $\{h = \epsilon\}$ is obviously homeomorphic to S^2 which means that $\{h < \epsilon\}$ is homeomorphic to \mathbb{R}^3 . If $n = 4$ then $\{h = \epsilon\}$ is a homotopy 3-sphere and a manifold. By the work of Freedman [Fre82, Corollary 1.3], this implies that the cone over $\{h = \epsilon\}$ is homeomorphic to \mathbb{R}^4 . If $n \geq 5$ then $\{h = \epsilon\}$ is a homotopy S^{n-1} and a manifold and hence is homeomorphic to S^{n-1} by the Poincare conjecture. □

Remark 3.3. As was suggested to the author by A. Lytchak, a different proof of Lemma 3.2 can be given for $n \geq 5$ by verifying that X satisfies the disjoint disk property and thus is a manifold by a result of Edwards [Dav00]. This method is used in [LN] to prove a version of stability for limits of manifolds with curvature bounded above. However, the author prefers his own argument given above.

Now that we know that X is a manifold and $f_i: M_i \rightarrow X$ are $o(i)$ -homotopy equivalences, we can apply the results from [CF79] for $n \geq 5$ and [Qui82] together with [CF79] for $n = 4$, which say that under these conditions f_i s can be $o(i)$ -approximated by homeomorphisms for all large i .

The same holds in dimension 3 by [Jak88] but only modulo the Poincare conjecture.

However, it is possible (and would certainly be a lot more preferable) that one can use the fact that all the subsets $\{h_i \leq \epsilon\}$ are actually topological balls and not merely contractible to give a proof in dimension 3 which does not rely on the Poincare conjecture.

Remark 3.4. By using relative versions of controlled homotopy theory results from [Qui82] and [CF79] mentioned above, it should be possible to generalize the above proof to the case of pointed Gromov-Hausdorff convergence of manifolds. This would amount to the manifold case of Theorem 7.11 below. Alternatively, one can handle the pointed case as follows.

Suppose we have a pointed convergence $(M_i^n, q_n) \rightarrow (X^n, q)$ where M_i^n are (possibly noncompact) Riemannian manifolds of $\text{sec} \geq \kappa$. The proof of Lemma 3.2 is obviously local and hence X is a topological manifold.

Let p be any point in X . Let h and h_i be the functions constructed in the proof of Lemma 3.2. Let Y_i be the double of $\{h_i \leq \epsilon\}$ and Y be the double of $\{h \leq \epsilon\}$. Obviously $Y_i \xrightarrow[i \rightarrow \infty]{G-H} Y$ and Y_i is homeomorphic to S^n for large i . While the metric on Y_i is not smooth along the boundary of $\{h_i \leq \epsilon\}$, it's easy to see that the proof of Lemma 3.2 still works for the convergence $Y_i \rightarrow Y$ and hence Y is a closed topological manifold. By the same controlled homotopy theory results used earlier, we conclude that Hausdorff approximations $Y \rightarrow Y_i$ can be $o(i)$ -approximated by homeomorphisms $g_i: Y \rightarrow Y_i$ for all large i . Restricting g_i to $\{h < \epsilon/2\}$ we obtain an open embedding of $\{h < \epsilon/2\}$ into M_i , which is $o(i)$ -close to the original Hausdorff approximation $(M, q) \rightarrow (M_i, q_i)$.

Finally, by using topological gluing theorem 4.11 below, for any fixed $R > 0$ we can glue finitely many such local homeomorphisms to get an open embedding $B(q, R) \rightarrow Y_i$, which is $o(i)$ -Hausdorff close to the original Hausdorff approximation $(M, q) \rightarrow (M_i, q_i)$.

4. BACKGROUND

4.1. Stratified spaces. Most of the material of this section is taken with almost no changes from [Per91] as no significant simplifications or improvements of the exposition seem to be possible.

Definition 4.1. *A metrizable space X is called an MSC-space (space with multiple conic singularities) of dimension n if every point $x \in X$ has a neighborhood pointed homeomorphic to an open cone over a compact $(n-1)$ -dimensional MCS space. Here we assume the empty set to be the unique (-1) -dimensional MCS-space.*

Remark 4.2. A compact 0-dimensional MCS-space is a finite collection of points with discrete topology.

Remark 4.3. An open conical neighborhood of a point in an MCS-space is unique up to pointed homeomorphism [Kwu64].

It easily follows from the definition that an MCS space has a natural topological stratification.

We say that a point $p \in X$ belongs to the l -dimensional strata X_l if l is the maximal number m such that the conical neighbourhood of p is pointed homeomorphic to $R^m \times K(S)$ for some MCS-space S . It is clear that X_l is an l -dimensional topological manifold.

We will need two general topological results which hold for spaces more general than Alexandrov spaces and follow from the general theory of deformations of homeomorphisms developed by Siebenmann [Sie72].

Theorem 4.4. [Sie72, Theorem 5.4, Corollary 6.14, 6.9]

Let X be a metric space and $f: X \rightarrow \mathbb{R}^k$ be a continuous, open, proper map such that for each $x \in X$ we have

- (1) $f^{-1}(f(x))$ is a compact MCS-space;
- (2) x admits a product neighborhood with respect to f , i.e. there exists an open neighbourhood U_x of x and a homeomorphism $F_x: U_x \rightarrow U_x \cap f^{-1}(f(x)) \times f(U_x)$ such that $f_x = p_2 \circ F_x$ where $p_2: U_x \cap f^{-1}(f(x)) \times f(U_x) \rightarrow \mathbb{R}^k$ is the coordinate projection onto the second factor.

Then f is a locally trivial fiber bundle.

Moreover, suppose we have in addition that $f(U_x) = I^k$. Let $K \subset U_x$ be a compact subset. Then there exists a homeomorphism $\varphi: f^{-1}(I^k) \rightarrow f^{-1}(f(x)) \times I^k$ respecting f (i.e. such that $f = p_2 \circ \varphi$. and such that $\varphi|_K = F_x|_K$).

The next gluing theorem is the key topological ingredient in the proof of the Stability Theorem. It says that for MCS spaces close local homeomorphisms given on a finite open covering can be glued to a nearby global homeomorphism under some mild (but important!) geometric assumptions.

First we need a technical definition.

Definition 4.5. A metric space X is called \varkappa -connected if for any two points $x_1, x_2 \in X$ there exists a curve connecting x_1 and x_2 of $\text{diam} \leq \varkappa(d(x_1, x_2))$.

Gluing Theorem 4.6. Let X be a compact MCS-space, $\{U_\alpha\}_{\alpha \in \mathfrak{A}}$ be a finite covering of X . Given a function \varkappa_0 , there exists $\varkappa = \varkappa(X, \{U_\alpha\}_{\alpha \in \mathfrak{A}}, \varkappa_0)$ such that the following holds:

Given a \varkappa_0 -connected MCS-space \tilde{X} , an open cover of \tilde{X} $\{\tilde{U}_\alpha\}_{\alpha \in \mathfrak{A}}$, a δ -Hausdorff approximation $\varphi: X \rightarrow \tilde{X}$ and a family of homeomorphisms $\varphi_\alpha: U_\alpha \rightarrow \tilde{U}_\alpha$, δ -close to φ , then there exists a homeomorphism $\tilde{\varphi}: X \rightarrow \tilde{X}$, $\varkappa(\delta)$ -close to φ .

Proof of Theorem 4.6. This proof of Theorem 4.6 is taken verbatim from [Per91].

We'll need two lemmas.

Lemma 4.7 (Deformation Lemma). Let X be a compact metric MCS-space, $W \Subset V \Subset U \subset X$ be open subsets.

Let $\varphi: U \rightarrow X$ be an open embedding δ -close to the inclusion i .

Then there exists an open embedding $\psi: U \rightarrow X$, $\varkappa(\delta)$ -close to i and such that $\psi \equiv \varphi$ on W and $\psi \equiv i$ on $U \setminus V$.

(Here \varkappa depends on W, U, V, X but not on φ).

Proof. Consider the open embedding $\varphi|_{U \setminus \bar{W}}: U \setminus \bar{W} \rightarrow X$. By the deformation theorem of Siebenmann [Sie72, Theorem 5.4], it can be perturbed to an open embedding $\varphi_1: U \setminus \bar{W} \rightarrow X$ which is $\varkappa(\delta)$ -close to i , coincides with i on some neighborhood of ∂V and is equal to φ outside some compact subset of $U \setminus \bar{W}$. Now let

$$\psi(x) = \begin{cases} \varphi(x), & \text{for } x \in W \\ \varphi_1(x), & \text{for } x \in U \setminus \bar{W} \\ x, & \text{for } x \in U \setminus V \end{cases}$$

It's clear that ψ satisfies the conclusion of the Lemma. □

Lemma 4.8. *Under the assumptions of Theorem 4.6, let $x \in X, \tilde{x} \in \tilde{X}$ satisfy $d(\varphi(x), \tilde{x}) < \delta$. Let $V \subset X$ be an open set containing $B(x, \varkappa_0(\delta) + 10\delta)$. Suppose $\psi: V \rightarrow \tilde{X}$ be an open embedding, δ -close to φ .*

Then $\tilde{x} \in \psi(V)$.

Proof. Let $\tilde{\gamma}: [0, 1] \rightarrow \tilde{X}$ be a curve of diam $\leq \varkappa_0(\delta)$ with $\tilde{\gamma}(0) = \psi(x), \tilde{\gamma}(1) = \tilde{x}$.

We'll show that $\tilde{\gamma}$ has a lift $\gamma: [0, 1] \rightarrow V$ with respect to ψ . Since ψ is an open embedding we can lift $\tilde{\gamma}$ on some interval $[0, \epsilon)$. Observe that given a lift γ of $\tilde{\gamma}$ on $[0, t)$ for some $t \leq 1$ it can always be extended to $[0, t]$ provided the closure of $\gamma([0, t))$ is contained in $B(x, \varkappa_0(\delta) + 10\delta)$. The fact that ψ is δ -close to φ and φ is a δ -Hausdorff approximation assures that this is always the case. Therefore the lift γ can be extended to $[0, 1]$ with $\psi(\gamma(1)) = \tilde{x}$. □

Remark 4.9. The proof of Lemma 4.8 is the only place in the proof of Theorem 4.6 where we use the assumption that \tilde{X} is \varkappa_0 -connected.

We can now continue with the proof of Theorem 4.6.

Suppose $U_{\alpha_1} \cap U_{\alpha_2} \neq \emptyset$. Let $U_1^4 \Subset U_1^3 \Subset U_1^2 \Subset U_1^1 \Subset U_{\alpha_1}$ and $U_2^4 \Subset U_2^3 \Subset U_2^2 \Subset U_2^1 \Subset U_{\alpha_2}$ be open subsets such that U_1^4, U_2^4 still cover $X \setminus \cup_{\alpha \in \mathfrak{A} \setminus \{\alpha_1, \alpha_2\}} U_\alpha$.

By Lemma 4.8, we have $\varphi_{\alpha_1}(U_1^1 \cap U_2^1) \subset \varphi_{\alpha_2}(U_{\alpha_2})$ provided δ is small enough.

Therefore we can consider the open embedding $\varphi_{\alpha_2}^{-1} \circ \varphi_{\alpha_1}: U_1^1 \cap U_2^1 \rightarrow U_{\alpha_2}$. Clearly, it is 2δ -close to the inclusion i . By Lemma 4.7, there exists an open embedding $\psi: U_1^1 \cap U_2^1 \rightarrow U_{\alpha_2}$, $\varkappa(\delta)$ -close to i and such that $\psi \equiv \varphi_{\alpha_2}^{-1} \circ \varphi_{\alpha_1}$ on $U_1^3 \cap U_2^3$ and $\psi \equiv i$ on $U_1^1 \cap U_2^1 \setminus U_1^3 \cap U_2^3$. We can extend ψ to U_2^1 by setting $\psi \equiv i$ on $U_2^1 \setminus U_1^1 \cap U_2^1$ and define $\varphi'_{\alpha_2} = \varphi_{\alpha_2} \circ \psi$.

Now we define $\varphi': U_1^4 \cup U_2^4 \rightarrow \tilde{X}$ by the formula

$$\varphi'(x) = \begin{cases} \varphi_{\alpha_1}(x) & \text{for } x \in U_1^4 \\ \varphi'_{\alpha_2}(x) & \text{for } x \in U_2^4 \end{cases}$$

It's clear that φ' is an open immersion and it's actually an embedding provided δ is small enough.

Moreover, by Lemma 4.8 we have

$$\tilde{X} \setminus \bigcup_{\alpha \in \mathfrak{A} \setminus \{\alpha_1, \alpha_2\}} \tilde{U}_\alpha \subset \varphi'(U_1^4 \cup U_2^4).$$

Now the statement of the theorem immediately follows by induction on the number of elements in \mathfrak{A} . □

In fact, we will need a somewhat stronger version of this theorem which assures that the gluing can be done relative to a fiber bundle structure on all the limit and approximating spaces.

Theorem 4.10 (Strong Gluing Theorem). *Under the assumptions of Gluing Theorem 4.6 we are given in addition continuous maps $f: X \rightarrow \mathbb{R}^k$, $\tilde{f}: \tilde{X} \rightarrow \mathbb{R}^k$, $h: X \rightarrow \mathbb{R}$, $\tilde{h}: \tilde{X} \rightarrow \mathbb{R}$ and a compact set $K \subset X$ such that the following holds*

- (1) *for any U_α with $U_\alpha \cap K \neq \emptyset$ we have $(\tilde{f}, \tilde{h}) \circ \varphi_\alpha = (f, h)$*
- (2) *for any U_α with $U_\alpha \cap K = \emptyset$ we have $\tilde{f} \circ \varphi_\alpha = f$*
- (3) *for any U_α with $U_\alpha \cap K \neq \emptyset$, U_α is contained in a product neighbourhood with respect to (f, h)*
- (4) *for any U_α with $U_\alpha \cap K = \emptyset$, U_α is contained in a product neighbourhood with respect to f*

Then the gluing homeomorphism $\bar{\varphi}$ can be chosen to respect f on X and (f, h) on K (i.e. $(\tilde{f}, \tilde{h}) \circ \bar{\varphi} = (f, h)$ on K and $\tilde{f} \circ \bar{\varphi} = f$ on X).

Proof. The proof of Theorem 4.6 can be trivially adapted to prove Strong Gluing Theorem 4.10 once we observe that the theorem of Siebenmann quoted in the proof of Lemma 4.7 has a stronger version respecting products with \mathbb{R}^k [Sie72, Theorem 6.9] so that the deformation ψ given by Lemma 4.7 can be made to respect the product structure $X \cong X_1 \times \mathbb{R}^k$ if $\varphi: U \rightarrow X$ respects that product structure. □

For applications to pointed Gromov-Hausdorff convergence we will need the following local version of the Gluing theorem for which the requirement that the approximated space be \varkappa -connected can be slightly weakened. For simplicity, we only state the unparameterized version.

Theorem 4.11. *Let $U \Subset V \Subset W \subset X$ be relatively compact open subsets in an MCS-space X . Let $\{U_\alpha\}_{\alpha \in \mathfrak{A}}$ be a finite covering of \bar{W} with the property that if $U_\alpha \cap \bar{V} \neq \emptyset$ then $U_\alpha \Subset W$.*

Then given a function \varkappa_0 , there exists $\varkappa = \varkappa(X, U, V, W, \{U_\alpha\}_{\alpha \in \mathfrak{A}}, \varkappa_0)$ such that the following holds:

Given a \varkappa_0 -connected MCS-space X' , and subsets $U' \Subset V' \Subset W' \Subset X'$, an open cover $\{U'_\alpha\}_{\alpha \in \mathfrak{A}}$ of \bar{W}' , a δ -Hausdorff approximation $(\bar{W}, \bar{V}, \bar{U}) \rightarrow (\bar{W}', \bar{V}', \bar{U}')$ and a family of homeomorphisms $\varphi_\alpha: U_\alpha \rightarrow U'_\alpha$, δ -close to φ ,

then there exists an open embedding $\varphi': V \rightarrow X'$, $\varkappa(\delta)$ -close to φ such that $\varphi(V) \supset U'$ if δ is sufficiently small.

Proof. The proof is exactly the same as the proof of Theorem 4.6 except in the induction procedure we only glue the embeddings of those U_α which intersect \bar{V} . □

Definition 4.12. *A map $f: X \rightarrow Y$ between two metric spaces is called ϵ -co-Lipschitz if for any $p \in X$ and all small R we have $f(B(p, R)) \supset B(f(p), \epsilon R)$.*

We will often make use of the following simple observation the proof of which is left to the reader as an exercise.

Lemma 4.13. *Let $f: X \rightarrow Y$ is ϵ -co-Lipschitz where X is compact. Let $p \in X$ and $\gamma: [0, 1] \rightarrow Y$ be a rectifiable curve with $\gamma(0) = f(p)$. Then there exists a lift $\tilde{\gamma}: [0, 1] \rightarrow X$ of γ such that $\tilde{\gamma}(0) = p$ and $L(\tilde{\gamma}) \leq \frac{1}{\epsilon} L(\gamma)$.*

4.2. Polar vectors.

Definition 4.14. Let $\text{curv}\Sigma \geq 1$. Given elements $u, v \in C\Sigma$ we define $\langle u, v \rangle$ by the formula

$$\langle u, v \rangle = |v| \cdot |u| \cdot \cos \angle uv$$

Definition 4.15. Let $\text{curv}\Sigma \geq 1$. Two vectors $u, v \in C\Sigma$ are called **polar** if for any $w \in C\Sigma$ we have

$$\langle v, w \rangle + \langle u, w \rangle \geq 0$$

More generally, u is called **polar** to a set $V \subset C\Sigma$ if for any $w \in C\Sigma$ we have

$$\sup_{v \in V} \langle v, w \rangle + \langle u, w \rangle \geq 0.$$

It is known (see [PP96] or [Pet06, Lemma 1.3.9]), that for any $v \in \Sigma$ there exists u , polar to v .

A function $f: \Sigma \rightarrow \mathbb{R}$ is called *spherically concave* if for any y lying on a shortest geodesic connecting x and $z \in \Sigma$ we have

$$f(y) \sin d(x, z) \geq f(z) \sin d(x, y) + f(x) \sin d(z, y).$$

As with ordinary concave functions, for a space with boundary we demand that the canonical extension of f to the doubling of Σ be spherically concave.

It's easy to see that f is spherically concave iff its 1-homogeneous extension to $C\Sigma$ is concave.

We will need the following property of polar vectors [Pet06, Section 1.3.8]:

Let $f: C\Sigma \rightarrow \mathbb{R}$ be concave and 1-homogeneous. Suppose $u \in C\Sigma$ is polar to $V \subset C\Sigma$. Then

$$(4.1) \quad f(u) + \inf_{v \in V} f(v) \leq 0.$$

Finally, we'll make use of the following fact [Pet06, Section 1.3.8]:

Given any two distinct points p, q in an Alexandrov space we have

$$(4.2) \quad \nabla_p d(\cdot, q) \text{ is polar to } \uparrow_p^q.$$

4.3. Gradient flows of semiconcave functions.

It was shown in [PP96] (cf. [KPT05]) that semiconcave functions on Alexandrov spaces admit well defined forward gradient flows.

Moreover, one can bound the Lipschitz constant of the gradient flows as follows.

Suppose $f: X \rightarrow \mathbb{R}$ is λ -concave. Let α and β are two f -gradient curves. Then

$$(4.3) \quad d(\alpha(t_1), \beta(t_1)) \leq d(\alpha(t_0), \beta(t_0)) \exp(\lambda(t_1 - t_0)) \text{ for all } t_1 \geq t_0$$

In particular, if f is concave then its gradient flow φ_t is 1-Lipschitz for any $t > 0$.

5. ADMISSIBLE FUNCTIONS AND THEIR DERIVATIVES

Definition 5.1. *Let X be an Alexandrov space of curv $\geq \kappa$. Let $f: X \rightarrow \mathbb{R}$ have the form $f = \sum_{\alpha} \lambda_{\alpha} \varphi_{\alpha}(d(\cdot, A_{\alpha}))$ where each A_{α} is a closed subset of X , $\lambda_{\alpha} \geq 0$, $\sum_{\alpha} \lambda_{\alpha} \leq 1$ and each $\varphi_{\alpha}: \mathbb{R} \rightarrow \mathbb{R}$ is a twice differentiable function with $0 \leq \varphi'_{\alpha} \leq 1$. We say that such f is admissible on $U = X \setminus \cup_{\alpha} A_{\alpha}$.*

With a slight abuse of notations we'll sometimes simply say that $f: X \rightarrow \mathbb{R}$ of the above form is admissible.

It's obvious from the definition that an admissible function is 1- Lipschitz and semi-concave on U . More precisely, f is λ -concave near $p \in U$ where λ depends on κ and $d(p, \cup_i A_i)$.

By the first variation formula, $df_p: \Sigma_p X \rightarrow \mathbb{R}$ has the form

$$df_p = \sum_{\alpha} -a_{\alpha} \cos(d(\cdot, \uparrow_p^{A_{\alpha}})) \text{ where } a_{\alpha} = \lambda_{\alpha} \varphi'_{\alpha}(d(\cdot, A_{\alpha})).$$

Note that $a_{\alpha} \geq 0$, $\sum_{\alpha} a_{\alpha} \leq 1$.

In view of this, following [Per93], we call a function $h: \Sigma \rightarrow \mathbb{R}$ where $\text{curv} \Sigma \geq 1$ a function of class DER if it has the form

$$\sum_{\alpha} -a_{\alpha} \cos(d(\cdot, A_{\alpha})) \text{ where } a_{\alpha} \geq 0, \quad \sum_{\alpha} a_{\alpha} \leq 1.$$

for some finite collection $\{A_{\alpha}\}_{\alpha \in \mathfrak{A}}$ of subsets of Σ .

DER functions are spherically concave on Σ and concave when radially extended to $C\Sigma$.

We define the scalar product of two functions in DER by the formula: for

$$(5.1) \quad h = \sum_{\alpha} -a_{\alpha} \cos(d(\cdot, A_{\alpha})), \quad g = \sum_{\beta} -b_{\beta} \cos(d(\cdot, B_{\beta}))$$

put

$$\langle h, g \rangle = \sum_{\alpha, \beta} a_{\alpha} b_{\beta} \cos(d(A_{\alpha}, B_{\beta})).$$

Note that this definition depends on the representations of h, g given by (5.1) and not just the values of f, g at every point.

It is shown in [Per93] that the scalar product satisfies the following properties:

- (i) $\langle d_p h, d_p g \rangle \leq \langle h, g \rangle - h(p)g(p)$ for any $p \in \Sigma$;
- (ii) $\langle h, h \rangle \geq (\inf_{g \in DER(\Sigma)} \langle h, g \rangle)^2 \geq 0$;
- (iii) For any $h \in DER(\Sigma)$ there is a point $\hat{A} \in \Sigma, 0 \leq \hat{a} \leq 1$ such that for $\hat{h} = -\hat{a} \cos(d(\cdot, \hat{A}))$ we have $\langle h, g \rangle \geq \langle \hat{h}, g \rangle$.

Notice that an admissible function f on an Alexandrov space X can be naturally lifted to a nearby Alexandrov space \tilde{X} by lifting the sets A_{α} from the definition of f to nearby sets in \tilde{X} and defining \tilde{f} by the same formula as f .

We will need the following simple lemma:

Lemma 5.2. *Let $X_i \xrightarrow{G-H}_{i \rightarrow \infty} X$ where $\text{curv} X_i \geq \kappa$ and let $f, g: X \rightarrow \mathbb{R}$ be admissible at $p \in X$. Let $f_i, g_i: X_i \rightarrow \mathbb{R}$ be natural lifts of f, g . Let $X_i \ni p_i \xrightarrow{i \rightarrow \infty} p$. Then*

$$\langle d_p f, d_p g \rangle \geq \limsup_i \langle d_{p_i} f_i, d_{p_i} g_i \rangle.$$

Proof. By linearity it's easy to see that it's enough to prove the lemma for f, g of the form $f = d(\cdot, A), g = d(\cdot, B)$. For functions of this form the statement easily follows from Toponogov comparison by an argument by contradiction. \square

6. ADMISSIBLE MAPS AND THEIR REGULAR POINTS

Definition 6.1. *A map $g: X \rightarrow \mathbb{R}^k$ is called admissible on an open set $U \subset X$ if it admits a representation $g = G \circ f$ where all the components of $f: U \rightarrow \mathbb{R}^k$ are admissible on U and G is a bi-Lipschitz homeomorphism between open sets in \mathbb{R}^k .*

Remark 6.2. The inclusion of the bi-Lipschitz map G in the definition of an admissible map might seem strange at this point. However, it will significantly simplify certain steps in the proof of the Stability theorem as well as prove useful in applications.

To unburden the exposition we will employ the following convention. If an admissible map is denoted by f (with any indices) we'll automatically assume that in the above definition $G \equiv \text{Id}$.

Definition 6.3. *An admissible map $g: U \rightarrow \mathbb{R}^k$ is called ϵ -regular at $p \in U$ if for some representation $g = G \circ f$ of g the following holds:*

- (1) $\min_{i \neq j} -\langle d_p f_i, d_p f_j \rangle > \epsilon$;
- (2) There exists $v \in \Sigma_p X$ such that $f'_i(v) > \epsilon$ for all i .

We say that g is regular at p if it's ϵ -regular at p for some $\epsilon > 0$.

It is easy to see that $f = (f_1, \dots, f_k)$ is ϵ -regular at p iff there is a point q such that for $f_0 = d(\cdot, q)$ we have

$$\langle d_p f_i, d_p f_j \rangle < -\epsilon \text{ for all } 0 \leq i \neq j \leq k.$$

Example 6.4. As was mentioned in the introduction, a basic example of a regular map is as follows. Suppose $\{f_i = d(\cdot, A_i)\}_{i=0, \dots, k}$ satisfy

$$\angle \uparrow_p^{A_i} \uparrow_p^{A_j} > \pi/2 \text{ for all } i \neq j.$$

Then $f = (f_1, \dots, f_k)$ is regular at p .

This example shows that regular points of admissible maps naturally generalize regular points of smooth maps because of the following simple observation:

Given $k + 1$ non-zero vectors in \mathbb{R}^n with all pairwise angles $> \pi/2$, any k of them are linearly independent.

As an obvious corollary of Lemma 5.2 we obtain

Corollary 6.5. *Let $f: X \rightarrow \mathbb{R}^k$ be admissible and ϵ -regular at $p \in X$. Let $0 < \epsilon' < \epsilon$. Suppose $X_i \xrightarrow{G-H}_{i \rightarrow \infty} X$ where $X_i \in \mathfrak{A}lcr^n(D, \kappa)$. Let $X_i \ni p_i \xrightarrow{i \rightarrow \infty} p$ and let $f_i: X_i \rightarrow \mathbb{R}^k$ be natural lifts of f . Then there exists $\delta > 0$ such that f_i is ϵ' -regular on $B(p_i, \delta)$ for all large i .*

In particular, the set of ϵ -regular (regular) points of an admissible map is open.

Remark 6.6. In [Per95], Perelman generalized all the Morse theory results from [Per93] to the more general and much more natural class of admissible functions consisting of 1-Lipschitz semiconcave functions. Corollary 6.5 is one of the main reasons why we restrict the class of admissible functions to the rather special semiconcave functions constructed from distance functions. Various definitions of regularity are possible for maps with semiconcave coordinates (see [Per95, Pet06]). While all these different definitions allow for relatively straightforward generalization of the results from [Per93], the author was unable to prove the analogue of Corollary 6.5 using any of these definitions.

Another (perhaps more serious) reason why we are forced to work with a small class of admissible functions is that there is currently no known natural way of lifting general semiconcave functions from the limit space to the elements of the sequence.

The following Lemma is due to Perelman.

Lemma 6.7. [Per93, Lemma 2.3], [Per95, Lemma 2.2], [Pet06, Lemma 8.1.4]

Suppose Σ^{n-1} has $\text{curv} \geq 1$ and let $f_0, \dots, f_k: C\Sigma \rightarrow \mathbb{R}$ be 1-homogeneous concave functions such that $\epsilon = \min_{i \neq j} -\langle f_i, f_j \rangle > 0$. Then

- (1) $k \leq n$;
- (2) There exists $w \in \Sigma$ such that $f_i(w) > \epsilon$ for all $i \neq 0$.
- (3) There exists $v \in \Sigma$ such that $f_0(v) > \epsilon$, $f_1(v) < -\epsilon$ and $f_i(v) = 0$ for $i = 2, \dots, k$.

Let $g: X \rightarrow \mathbb{R}^k$ be an admissible map. Let $X_{reg}^\epsilon(g) \subset X$ ($X_{reg}(g) \subset X$) be the set of ϵ -regular (regular) points of g .

Then the following properties hold [Per93, Per95]

- a) $X_{reg}^\epsilon(g)$ (and hence also $(X_{reg}(g))$) is open for any $\epsilon > 0$ (see Corollary 6.5).
- b) If g is ϵ -regular on an open set $U \subset X$ then g is ϵ -co-Lipschitz on U . (This is an easy corollary of Lemma 6.7. See [Per93, Per95] for details.)
- c) $g: X_{reg}(g) \rightarrow \mathbb{R}^k$ is open.
- d) $k \leq \dim X$ if $X_{reg}(g) \neq \emptyset$. This immediately follows from part (1) of Lemma 6.7.

Theorem 6.8 (Local Fibration Theorem). [Per93, Per95] *Let $g: X \rightarrow \mathbb{R}^k$ be an admissible map. Then $g|_{X_{reg}(g)}: X_{reg}(g) \rightarrow \mathbb{R}^k$ is locally a topological bundle map. This means that any point $p \in X_{reg}(g)$ possesses an open product neighborhood with respect to g with an MSC-space as a fiber.*

We will use the original Local Fibration Theorem without a proof. However, we will prove a more general version of it (see Theorem 9.7) in Section 9 in order to prove the Relative Stability Theorem.

We will need the following technical Lemma due to Perelman which plays a key role in the proofs of both the Stability Theorem and the Fibration Theorem above.

Key Lemma 6.9. [Per93, Section 3] *Let p be a regular point of $f: X \rightarrow \mathbb{R}^k$. Suppose f is incompressible at p , i.e. for any admissible function $f_1: X \rightarrow \mathbb{R}$, the point p is not regular for $(f, f_1): X \rightarrow \mathbb{R}^{k+1}$.*

Then there exists an admissible function $h: X \rightarrow \mathbb{R}$ with the following properties

- (i) $h(p) = 0$.
- (ii) h is strictly concave on $B(p, R)$ for some $R > 0$.
- (iii) There are $r > 0, A > 0$ such that $h < A$ on $f^{-1}(\bar{I}^k(f(p), r))$ and $f^{-1}(\bar{I}^k(f(p), r)) \cap \{h \geq -A\}$ is compact in $B(p, R)$.
- (iv) h has a unique maximum in $B(p, R) \cap f^{-1}(v)$ for all $v \in \bar{I}^k(f(p), r)$. Let S denote the set of such maximum points.
- (v) (f, h) is regular on $(\bar{I}^k(f(p), r) \cap B(p, R)) \setminus S$.

As an immediate corollary we obtain

Corollary 6.10. *Let $f: X^n \rightarrow \mathbb{R}^n$ be an admissible map. Then f is locally bi-Lipschitz on the set of its regular points $X_{reg}(f)$.*

Proof of Corollary 6.10. Let p be a regular point of f . By part a) of Lemma 6.7, f is incompressible at p . Let h be the function in $B(p, R)$ provided By lemma 6.9. By part (v) of Lemma 6.9, $(f, h): B(p, R) \setminus S \rightarrow \mathbb{R}^{n+1}$ is regular. However, as was just mentioned, by part (1) of Lemma 6.7, a map from X^n to \mathbb{R}^{n+1} can not have any regular points. Therefore $B(p, R) \setminus S = \emptyset$. By part (iv) of Lemma 6.9, this is equivalent to saying that $B(p, R) \cap f^{-1}(v)$ consists of a single point for all $v \in \bar{I}^k(f(p), r)$ which means that f is injective near p . Finally, recall that being regular, f is both Lipschitz and co-Lipschitz near p , which together with local injectivity means that it's locally bi-Lipschitz on $X_{reg}(f)$. \square

Proof of Key Lemma 6.9. Since a complete proof is given in [Per93] we don't include as many details.

For simplicity we assume that all components f_j of f are actually concave near p (the proof can be easily adapted to the general case of semiconcave f_j s). Since f is ϵ -regular at p there is a point q near p such that $f_j(q) > f_j(p) + \epsilon d(p, q)$ for all $j = 1, \dots, k$. Since f_j 's are 1-Lipschitz, for all r near p and all $x \in B(q, \epsilon d(p, q)/4)$ we have

$$(6.1) \quad f_j(x) > f_j(r) + \epsilon \frac{d(x, r)}{4}$$

Fix a small positive $\delta \ll \epsilon d(p, q)$. Choose a maximal δ -net $\{q_\alpha\}_{\alpha \in \mathfrak{A}}$ in $S(q, \epsilon d(p, q)/4)$. A standard volume comparison argument shows that $|\mathfrak{A}| = N \geq c\delta^{1-n}$ where $n = \dim X$.

Consider the function $h = N^{-1} \sum_{\alpha} h_{\alpha}$ where $h_{\alpha} = \varphi_{\alpha}(d(\cdot, q_{\alpha}))$ and $\varphi_{\alpha}: \mathbb{R} \rightarrow \mathbb{R}$ is the unique continuous function satisfying the following properties

- (1) $\varphi'_{\alpha}(t) = 1$ for $t \leq d(p, q_{\alpha}) - \delta$;
- (2) $\varphi'_{\alpha}(t) = 1/2$ for $t \geq d(p, q_{\alpha}) + \delta$;
- (3) $\varphi''_{\alpha}(t) = -\frac{1}{4\delta}$ for $d(p, q_{\alpha}) - \delta < t < d(p, q_{\alpha}) + \delta$;
- (4) $\varphi_{\alpha}(d(p, q_{\alpha})) = 0$

It is clear that h is admissible near p with $h(p) = 0$ and

$$(6.2) \quad \langle d_r h, d_r f_j \rangle < -\epsilon/8 \text{ for all } j = 1, \dots, k \text{ and all } r \text{ near } p$$

By [Per93, Lemma 3.6], h is strictly $c\delta^{-1}$ -concave on $B(p, \delta)$ (see also [Kap02, Lemma 4.2] for a more detailed proof of the same statement).

Denote $\Sigma_p^\epsilon = \{\xi \in \Sigma_p \mid f'_j(\xi) > \epsilon \text{ for all } j = 1, \dots, k\}$.

It easily follows from the definition that f is incompressible at p iff $\text{diam}(\Sigma_p^0) \leq \pi/2$.

Sublemma 6.11. *If $\text{diam}(\Sigma_p^0) \leq \pi/2$ then for all $r \in B_p(\delta)$ we have*

$$(6.3) \quad h(r) \leq h(p) - c \cdot d(p, r) + c \cdot \max_j \{0, f_j(p) - f_j(r)\}$$

Proof. Since h is 1-Lipschitz and f is ϵ -co-Lipschitz, by using Lemma 4.13, it's enough to prove the Sublemma for $r \in B(p, \delta)$ satisfying $f_j(r) \geq f_j(p)$ for all $j = 1, \dots, k$. Then $\uparrow_p^r \subset \Sigma_p^0$. By construction we also have that $\uparrow_p^{q_\alpha} \subset \Sigma_p^0$ which by assumption of the sublemma implies that $\angle \uparrow_p^r \uparrow_p^{q_\alpha} \leq \pi/2$ for all α . (This is the only place in the proof where we use that $\text{diam}(\Sigma_p^0) \leq \pi/2$ i.e that f is incompressible at p). This means that the derivative of h_α at p in the direction of r is ≤ 0 . By semi-concavity of h_α this implies

$$(6.4) \quad h_\alpha(r) \leq h_\alpha(p) - \lambda \cdot d(p, r)^2$$

Moreover, a volume comparison argument (see [Per93] or [Kap02, Lemma 4.2] for details) shows that for most $\alpha \in \mathfrak{A}$ we actually have $\angle \uparrow_p^r \uparrow_p^{q_\alpha} \leq \pi/2 - C$.

Indeed, recall that $N \geq c\delta^{1-n}$.

Fix a small $\mu > 0$. By the first variation formula and semi-concavity of h_α we see that if $\angle \uparrow_p^r \uparrow_p^{q_\alpha} \leq \pi/2 - \mu$ then

$$(6.5) \quad h_\alpha(r) \leq h_\alpha(p) - \lambda \cdot d(p, r)^2 - \frac{1}{2}\mu \cdot d(p, r)$$

A standard volume comparison argument shows that the $n - 1$ -volume of the set

$A_\mu = \{\xi \in \Sigma_p \text{ such that } \pi/2 - \mu \leq \angle \xi \uparrow_p^r \leq \pi/2\}$ is bounded above by $c\mu$. By another standard volume comparison this implies that the maximal number of points in A_μ with pairwise angles $\geq \delta$ is at most $c\mu\delta^{1-n}$.

This means that if $\mu \ll c$ then for the vast majority of q_α we must have $\angle \uparrow_p^r \uparrow_p^{q_\alpha} \leq \pi/2 - \mu$.

A suitable choice of μ now immediately yields $h(r) \leq h(p) - c \cdot d(p, r)$. \square

Sublemma 6.11 obviously implies that inside $B(p, \delta)$ the sets $\{h \geq -c\delta\} \cap f^{-1}(v)$ are compact for all v close to $f(p)$ which proves (iii).

It remains to prove parts (iv) and (v) of Key Lemma 6.9. For any $v \in \mathbb{R}^k$ denote $U_v = B(p, \delta) \cap f^{-1}(v)$ and $U_v^+ = \{x \in B(p, \delta) \mid f_j(x) \geq v_j \text{ for all } j = 1, \dots, k\}$

Sublemma 6.12. *Let $z \in U_v$ be a point of maximum of h on U_v where $|f_j(p) - v_j| \leq \delta^2, j = 1, \dots, k$. Then for every $x \in U_v \cap B(p, \delta/2)$ we have*

$$h(z) \geq h(x) + c\delta^{-1}d(x, z)^2$$

Proof. First we notice that

$$(6.6) \quad \max_{U_v^+} h = \max_{U_v^+ \cap \bar{B}(p, \delta/2)} h = \max_{U_v \cap \bar{B}(p, \delta/2)} h = h(z)$$

The first and the last equalities follow from Sublemma 6.11 and the fact that h is 1-Lipschitz and f is ϵ -co-Lipschitz. If the equality in the middle is violated, then there exists a maximum point $r \in U_v^+ \cap \bar{B}(p, \delta/2)$ such that $f_l > v_l$ for some l . By applying part (2) of Lemma 6.7 to $d_r h, d_r f_j$ in $\Sigma_r X$ we can find a direction $\xi \in \Sigma_r X$ such that $h'(\xi) > 0$ and $f_j'(\xi) > 0$ for all $j \neq l$. This contradicts the fact that r is a point of maximum. This proves (6.6).

Now consider the midpoint y of a shortest curve connecting x and z . By concavity of f and strict concavity of h we get

$$(6.7) \quad h(y) \geq \frac{h(x) + h(z)}{2} + c\delta^{-1}d(x, z)^2, \text{ and}$$

$$f_j(y) \geq f_j(z) \text{ for all } j = 1, \dots, k$$

In particular $y \in U_v^+$ and therefore $h(y) \leq h(z)$. Combining this with (6.7) we immediately get the statement of Sublemma 6.12. □

To complete the proof of Lemma 6.9 it remain to verify (v).

Let $z \in U_v \cap B(p, \delta)$ be the point of maximum of h on U_v . For any other point $x \in U_v$, by Sublemma 6.11 combined with Lemma 4.13, we can find a point s arbitrary close to z and such that

$$f_j(s) > f_j(x) \text{ for all } j \text{ and } h(s) > h(x) + c\delta^{-1}d(x, s)^2$$

Let $\xi = \uparrow_x^s$. By concavity of h and f_j 's it is obvious that $h'(\xi) > 0$ and $f_j'(\xi) > 0$ for all j . Combined with (6.2) this means that (f, h) is regular at x . □

Remark 6.13. Key Lemma 6.9 in conjunction with Theorem 4.4 easily yields Fibration Theorem 6.8 (see [Per93] for details).

We will prove a more general version of it in Section 9.

We will need the following strengthened version of the Key Lemma which is the main geometric ingredient in the proof of the Stability Theorem.

Lemma 6.14. *Under the conditions of Lemma 6.9 suppose we have a noncollapsing converging sequence $X_i^n \xrightarrow[i \rightarrow \infty]{G-H} X^n$ and admissible functions f_i on X_i converging to f . Let $p_i \in X_i$ satisfy $p_i \xrightarrow[i \rightarrow \infty]{} p$.*

Then there exist admissible lifts h_i of the function h provided by Lemma 6.9 such that for all large i , the functions f_i, h_i satisfy the properties (i)-(v) of Lemma 6.9.

Proof. Part (i) is obvious by construction of h and h_i where we might have to shift h_i by a small constant to make it zero at p_i .

The fact that the natural lifts h_i of h are strictly concave in $B(p_i, R)$ is basically the same as the proof of the concavity of h itself (see [Per93, Lemma 3.6]). It is carried out in full detail in [Kap02, Lemma 4.2]. This proves part (ii).

For the proof of (iii) we can not use Sublemma 6.11 as we did in the proof of Lemma 6.9 because f_i might be complementable everywhere near p_i . Nevertheless, part (iii) is obvious because it holds for (f, h) by Lemma 6.9 and $f_i \rightarrow f, h_i \rightarrow h, p_i \rightarrow p$.

Parts (iv) and (v) are proved in exactly the same way as in the proof of Lemma 6.9. \square

Lemma 6.15. *Let $F = (f_1, \dots, f_k): X \rightarrow \mathbb{R}^k$ be ϵ -regular near p . Then the level set $H = \{F = F(p)\}$ is locally \varkappa -connected near p for a linear function \varkappa .*

Proof. In what follows all constants C will depend on ϵ .

Denote $F(p) = v = (v_1, \dots, v_k)$ and let H_- be the set $\cap_i \{f_i \leq v_i\}$.

By definition of a regular point, there exists q near p such that for $\xi = \uparrow_p^q$ we have $f'_i(\xi) > \epsilon$ for all i (and the same holds for all z near p). Let $\eta = \nabla_p d(\cdot, q)$. Then η is polar to ξ by (4.2). Therefore $df_i(\xi) + df_i(\eta) \leq 0$ for all i by (4.1) and hence

$$(6.8) \quad df_i(\eta) \leq -\epsilon \text{ for all } i.$$

Let $x, y \in B(p, r)$ with $r \ll d(p, q)$ be two close points on the level set H . Let γ be a shortest geodesic connecting x and y . Consider the gradient flow φ_t of $d(\cdot, q)$ in $B(p, r)$.

By above, all f_i 's decrease with the speed at least ϵ along φ_t . Since all f_i are 1-Lipschitz we know that along γ we have $f_i \leq f_i(p) + d(x, y)$. By (6.8) this implies that for some $t \leq C \cdot d(x, y)$ we can guarantee that $\gamma_1 = \varphi_t(\gamma)$ lies in H_- .

By shifting f_i 's by constants we can assume that all v_i 's are the same and equal to a . Since $d(\cdot, q)$ is λ -concave in $B(p, r)$ (where λ depends on $d(p, q)$ and the lower curvature bound of X), by the Lipschitz properties of the gradient flows (4.3), we know that $L(\gamma_1) \leq C \cdot d(x, y)$.

Let γ_2 be the concatenation of the gradient curve of $d(\cdot, q)$ through x followed by γ_1 followed by the the gradient curve of $d(\cdot, q)$ through y taken in opposite direction. Then by above we still have that $L(\gamma_2) \leq C \cdot d(x, y)$. We have also shown that $\gamma_2 \subset H_-$.

Let $f = \min(a, \min_i f_i)$. Then f is still semiconcave with the same concavity constants as f_i 's.

Observe that the gradient flow of f takes H_- to H (the points of H do not move under the flow).

By construction we have that $f \geq f(p) - C \cdot d(x, y)$ along γ_2 . By ϵ -regularity of F near p and Lemma 6.7 we see that $|\nabla f| \geq \epsilon$ on $\{f < a\} \cap B(p, r)$ and hence the gradient flow of f pushes γ_2 into a curve $\gamma_3 \subset H$ connecting x and y in uniformly bounded time. Once again applying the Lipschitz properties of gradient flows we obtain that $L(\gamma_3) \leq C \cdot d(x, y)$. \square

Remark 6.16. It easily follows from the proof that the function \varkappa provided by Lemma 6.15 is semi-continuous under Gromov-Hausdorff convergence in the following sense.

Suppose $X_i^n \rightarrow X^n$ be a convergent sequence of *compact* Alexandrov spaces with $\text{curv} \geq \kappa$ and $f_i: X_i \rightarrow \mathbb{R}^k$ be a sequence of admissible maps with λ -concave 1-Lipschitz components converging to $f: X \rightarrow \mathbb{R}^k$. Suppose f is ϵ -regular near p and $p_i \in X_i$ converges to p . Then, by Corollary 6.5, f_i is ϵ -regular on $B(p_i, r)$ for some $r > 0$ for all large i and the level sets $\{f_i = f_i(p_i)\}$ are \varkappa -connected near p_i for all large i with the same $\varkappa(t) = C(\epsilon, \kappa, \lambda, X) \cdot t$.

7. PROOF OF THE STABILITY THEOREM

Definition 7.1. Let p be a point in an Alexandrov space X . Let $g = G \circ f = (g^1, \dots, g^k): X \rightarrow \mathbb{R}^k$ be regular at p . Then an open product neighborhood of p with respect to such g is called a product neighborhood of a p of rank k .

Definition 7.2. We'll call a subset H of \mathbb{R}^n a generalized quadrant if it has the following form

$$H = \bigcap_{i \in I} \{x_i \geq c_i\} \cap \bigcap_{j \in J} \{x_j \leq c_j\}$$

where I, J are some (possibly empty) subsets of $\{1, \dots, n\}$ and (x_1, \dots, x_n) are the standard coordinates on \mathbb{R}^n .

Definition 7.3. A compact subset P in an Alexandrov space X is called k -framed if P can be covered by a finite collection of open sets U_α such that each U_α is a product neighborhood of rank $k_\alpha \geq k$ for some $p_\alpha \in P$ with respect to some

$$g_\alpha = G_\alpha \circ f_\alpha: X \rightarrow \mathbb{R}^{k_\alpha}$$

and $P \cap U_\alpha = g_\alpha^{-1}(H_\alpha) \cap U_\alpha$ where H_α is a generalized quadrant in \mathbb{R}^{k_α} .

We say that a framing $\{U_\alpha, f_\alpha, H_\alpha\}_{\alpha \in \mathfrak{A}}$ respects a map $f: X \rightarrow \mathbb{R}^l$ if the first l functions of f_α coincide with f for all α .

More generally, we'll say that a framing $\{U_\alpha, g_\alpha = G_\alpha \circ f_\alpha, H_\alpha\}_{\alpha \in \mathfrak{A}}$ respects a representation of an admissible map g

$$g = G \circ f: X \rightarrow \mathbb{R}^l$$

if the first l functions of each f_α coincide with f and G_α has the form $G_\alpha(x, y) = (G(x), T_\alpha(x, y))$ where $x \in \mathbb{R}^l, y \in \mathbb{R}^{k_\alpha - l}$, i.e the first l coordinates of G_α are equal to G . (In particular, the first l functions of g_α also coincide with g).

To simplify notations, we'll often simply say that a framing respects a map g to mean that it respects the representation $g = G \circ f$.

Example 7.4. Any compact Alexandrov space has a zero framing.

Example 7.5. For any point $p \in X$ there exists $\epsilon > 0$ such that $d(\cdot, p)$ has no critical points in $B(p, \epsilon) \setminus \{p\}$. Hence, for any positive $r < R < \epsilon$, the annulus $A(r, R, p) = \bar{B}(p, R) \setminus B(p, r)$ is 1-framed respecting $f = d(\cdot, p)$.

Suppose $X_i^n \rightarrow X^n$ is a converging noncollapsing sequence of compact Alexandrov spaces with $\text{curv} \geq \kappa$. Let $\theta_i: X \rightarrow X_i$ be a sequence of $o(i)$ -Hausdorff approximations.

Let $P \subset X$ be a k -framed compact subset of X . We define the corresponding k -framed subsets $P_i \subset X_i$ as follows. Let $g_\alpha = G_\alpha \circ f_\alpha$ be a representation of g_α given by the definition of an admissible map. We lift the defining functions f_α and f to f_i and $f_{\alpha,i}$ in the natural way. Suppose $G_{\alpha,i}$ is a sequence of *uniformly* bi-Lipschitz homeomorphisms of open sets in \mathbb{R}^k converging to G . Put $g_{\alpha,i} = G_{\alpha,i} \circ f_{\alpha,i}$.

Then $g_{\alpha,i}$ will still be admissible and regular on the corresponding subsets of X_i by Corollary 6.5. In particular we get product neighborhoods $U_{\alpha,i}$ with respect to $g_{\alpha,i}$.

We'll say that a compact set $P_i \subset X_i$ is a lifting of P if $P_i \cap U_{\alpha,i} = g_{\alpha,i}^{-1}(H_\alpha)$ for all α .

Remark 7.6. Note that a lifting of P need not exist! However, if it does, it is automatically k -framed. Moreover, if P is k -framed with respect to $f: X \rightarrow \mathbb{R}^l$ and $I_\alpha, J_\alpha \subseteq \{1, \dots, l\}$ for all α then the lifting exists for all large i . In particular, if X is a compact Alexandrov space and $X_i^n \rightarrow X^n$ with $\text{curv} \geq \kappa$ then the lifting of X with respect to a zero framing exists for any large i and is equal to X_i .

Lemma 7.7. *The set's P_i are \varkappa -connected for all large i and the same $\varkappa(t) = Ct$.*

Proof. Let x, y be two close points in $P \cap U_\alpha$ for some α . Since g_α is L -Lipschitz, $|g_\alpha(x) - g_\alpha(y)| \leq L \cdot d(x, y)$. Since H_α is convex, the straight line segment connecting $g_\alpha(x)$ and $g_\alpha(y)$ lies in H_α . Since f_α (and hence g_α) is ϵ -co-Lipschitz, by Lemma 4.13 we can lift it to a curve $\gamma_1: [0, 1] \rightarrow U_\alpha \cap P$ of length $\leq \frac{L}{\epsilon} d(x, y)$ with $\gamma_1(0) = x$. Observe that $\gamma_1(1)$ and y lie in the same fiber of g_α (and hence of f_α) and $d(\gamma_1(1), y) \leq C(L, \epsilon)d(x, y)$ by the triangle inequality. By Lemma 6.15 we can connect $\gamma_1(1)$ and y by a curve γ_2 inside the fiber $\{g_\alpha = g_\alpha(y)\}$ of length $\leq \tilde{C} \cdot d(\gamma_1(1), y)$. The concatenation of γ_1 and γ_2 provides a curve γ in $P \cap U_\alpha$ connecting x to y with $L(\gamma) \leq C \cdot d(x, y)$.

As was observed in Remark 6.16 the constant \tilde{C} in the above argument can be chosen to be the same for all $f_{\alpha,i}$ and hence, since all $G_{\alpha,i}$ are *uniformly* bi-Lipschitz, all P_i are \varkappa -connected for all large i for the same $\varkappa(t) = Ct$. □

The proof of the stability theorem proceeds by reverse induction in framing and, in fact, it requires us to prove the following stronger version of it:

Theorem 7.8 (Parameterized Stability Theorem). *Suppose $X_i^n \rightarrow X^n$ is a converging noncollapsing sequence of Alexandrov spaces with curvature bounded below and diameter bounded above. Let $\theta_i: X \rightarrow X_i$ be a sequence $o(i)$ -Hausdorff approximations.*

Let $P \subset X$ be a k -framed compact subset of X whose framing respects $g: X \rightarrow \mathbb{R}^l$. Let $K \subset P$ be a compact subset such that the framing of P respects g on P and (g, h) on K for some $g: X \rightarrow \mathbb{R}^l$ and $h: X \rightarrow \mathbb{R}$.

Then for all large i there exist homeomorphisms $\theta'_i: P \rightarrow P_i$ such that θ'_i is $o(i)$ -close to θ_i and respects g on P and (g, h) on K .

Proof. We proceed by reverse induction in k .

If $k = n$ then the locally defined maps $g_{\alpha,i}: U_{\alpha,i} \rightarrow \mathbb{R}^n, g_\alpha: U_\alpha \rightarrow \mathbb{R}^n$ are bi-Lipschitz homeomorphisms. By construction, the maps $\theta_{\alpha,i} = g_{\alpha,i}^{-1} \circ g_\alpha: U_\alpha \rightarrow U_{\alpha,i}$ are homeomorphisms Hausdorff close to θ_i . Moreover, by construction, $\theta_{\alpha,i}$ sends $P \cap U_\alpha$ onto $P_i \cap U_{\alpha,i}$.

Thus the statement of the theorem follows from Strong Gluing Theorem 4.10 and Lemma 7.7.
Induction step.

Suppose the theorem is proved for $k + 1 \leq n$ and we need to prove it for k .

Let P be k -framed and let p lie in P . Then $p \in U_\alpha$ for some α . Let $g_\alpha = G_\alpha \circ f_\alpha: U_\alpha \rightarrow \mathbb{R}^k$ be the admissible map regular at p coming from the definition of a k -framed set.

To simplify the notations we will assume that $G_\alpha = \text{Id}$ and $f_\alpha = g_\alpha$. The proof in the general case easily follows from this one with obvious modifications.

Let $p_i = \theta_i(p)$.

By possibly adding more components to f_α we can construct an admissible map to $f_p: X \rightarrow \mathbb{R}^{k_p}$ where $k_p \geq k$ which is incomplementable at p .

Let $h: B(p, R) \rightarrow R$ be a strictly concave function provided by Lemma 6.9. By choosing a sufficiently small $r, A > 0$ we can assume that the set $U_p = f_p^{-1}(I^{k_p}(f_p(p), r) \cap \{h \geq -A\})$ is compact. By reducing r further we can assume that $|h(x)| \leq a \ll A$ for $x \in S \cap U_p$. We will call U_p a special neighborhood of p .

By Lemma 6.9 we have that U_p is a k -framed compact subset of X .

Since h and all the coordinates of f_p are admissible, they have natural admissible lifts h_i and $f_{p,i}$ which define corresponding neighborhoods U_{p_i} of p_i .

The proof of Stability Theorem 7.8 will easily follow from the following

Local Stability Lemma 7.9. *For all large i there exist homeomorphisms $\theta_{p,i}: U_p \rightarrow U_{p_i}$ respecting f_p and $o(i)$ -close to the Hausdorff approximation θ_i .*

Let us first explain how to finish the proof of theorem 7.8 given Lemma 7.9.

Choose a finite cover of P by the interiors of the special neighborhoods U_{p_β} . For all large i , Lemma 7.9 provides homeomorphisms $\theta_{p_\beta,i}: U_{p_\beta} \rightarrow U_{p_\beta,i}$ respecting f_{p_β} and $o(i)$ -close to θ_i .

Observe that since each $\theta_{p_\beta,i}$ respects f_{p_β} , it sends $P \cap U_{p_\beta}$ onto $P_i \cap U_{p_\beta,i}$. Taking into account Lemma 7.7 we can apply Gluing Theorem 4.10 to obtain the desired homeomorphism $\theta'_i: P \rightarrow P_i$.

Proof of Local Stability Lemma 7.9. All throughout the proof of the Lemma we will work only with points in U_p in X and U_{p_i} in X_i .

If $k_p > k$ then the statement follows directly from the induction hypothesis.

Let's suppose $k_p = k$.

First we change the function h to an auxiliary function \tilde{h} by shifting h by a constant on each fiber of f to make it identically zero on S .

More precisely, let $\tilde{h}(x) = h(x) - h(S \cap f^{-1}(f(x)))$. Recall that by Lemma 6.9(iii), $S \cap f^{-1}(f(x))$ consists of a single point so that this definition makes sense. Also by Lemma 6.9(iii), we have $h(S \cap f^{-1}(f(x))) = \max_{y \in U_p \cap f^{-1}(f(x))} h(y)$ and therefore $\tilde{h} = h - H \circ f$ where $H: \mathbb{R}^k \rightarrow \mathbb{R}$ is given by $H(v) = \max_{x \in U_p \cap f^{-1}(v)} h(x)$. Since f is co-Lipschitz and h is Lipschitz, using Lemma 4.13 we easily conclude that H is Lipschitz.

In particular, $(f, \tilde{h}) = \bar{H} \circ (f, h)$ where \bar{H} is a bi-Lipschitz homeomorphism of some open domains in \mathbb{R}^{k+1} given by $\bar{H}(a, b) = (a, b - H(a))$.

Therefore, we still have that (f, \tilde{h}) is regular on $U_p \setminus S$ and hence it's locally a bundle map on $U_p \setminus S$ by Theorem 6.8.

In addition, by construction, $\tilde{h} = 0$ on S and $\tilde{h} < 0$ on $U_p \setminus S$.

We define \tilde{h}_i, H_i and \bar{H}_i in a similar fashion using f_i, h_i . We obviously have that $\tilde{h}_i \rightarrow \tilde{h}, H_i \rightarrow H, \bar{H}_i \rightarrow \bar{H}$. Moreover, since f_i are uniformly co-lipschitz, all H_i are uniformly Lipschitz and hence all \bar{H}_i are uniformly bi-Lipschitz.

Then we again have that $\tilde{h}_i = 0$ on S_i and $\tilde{h} < 0$ on $U_{p_i} \setminus S_i$. By Lemma 6.14 we also have that (f_i, \tilde{h}_i) is regular on $U_{p_i} \setminus S_i$.

Fix a small $\delta \ll A$. Then the set $\{\tilde{h} \leq -\delta\}$ is $(k+1)$ -framed with the corresponding sets in U_{p_i} given by $\{\tilde{h}_i \leq -\delta\}$ and therefore, by induction assumption, for large i there exist homeomorphisms $\theta'_{\delta,i}: \{\tilde{h} \leq -\delta\} \rightarrow \{\tilde{h}_i \leq -\delta\}$ respecting (f, \tilde{h}) and $o(i|\delta)$ -close to θ_i .

In particular, the fiber F_i of (f_i, \tilde{h}_i) is homeomorphic to the fiber F of (f, \tilde{h}) for all large i .

Next consider the set $\{\tilde{h} > -3\delta/2\}$ and consider the map $(f, \tilde{h}): \{\tilde{h} > -3\delta/2\} \setminus S \rightarrow I^k(f(p), r) \times (-3\delta/2, 0)$.

By Lemma 6.9, this map is regular and proper. Therefore, by Theorem 6.8 and Theorem 4.4, it is a fiber bundle.

Hence, $\{\tilde{h} > -3\delta/2\} \setminus S$ is homeomorphic to $I^k(f(p), r) \times (-3\delta/2, 0) \times F$ with the first two coordinates given by (f, \tilde{h}) . By restriction this gives a homeomorphism $\{\tilde{h} \geq \delta\} \setminus S$ to $I^k(f(p), r) \times [-\delta, 0) \times F$ with the first two coordinates still given by (f, \tilde{h}) .

By Lemma 6.9 and construction of \tilde{h} , \tilde{h} has a unique max (equal to zero) on $f^{-1}(v)$ for any $v \in \bar{I}^k(f(p), r)$. Therefore, the above homeomorphism can be uniquely extended to a homeomorphism $\varphi_\delta: \{\tilde{h} \geq -\delta\} \rightarrow \bar{I}^k(f(p), r) \times KF$ respecting (f, \tilde{h}) .

Similarly, for all large i we have homeomorphisms $\varphi_{\delta,i}: \{\tilde{h}_i \geq -\delta\} \rightarrow \bar{I}^k(f(p), r) \times KF_i$ respecting (f_i, \tilde{h}_i) .

Recall that F_i is homeomorphic to F for large i . Modulo that homeomorphism we can take the composition $\varphi_{\delta,i}^{-1} \circ \varphi_\delta$ and obtain a homeomorphism $\theta''_{\delta,i}: \{\tilde{h} \geq -\delta\} \rightarrow \{\tilde{h}_i \geq -\delta\}$ respecting (f, \tilde{h}) .

Since $\theta''_{\delta,i}$ respects f and S_i is $o(i)$ -close to S , it is clear that $\theta''_{\delta,i}$ is $\varkappa(\delta) + o(i|\delta)$ -close to θ_i .

Gluing $\theta''_{\delta,i}$ and $\theta'_{\delta,i}$ we obtain homeomorphisms $\theta_{\delta,i}: U_p \rightarrow U_{p_i}$ respecting f and $\varkappa(\delta) + o(i|\delta)$ -close to θ_i .

Since δ was arbitrary, a standard diagonal argument provides the desired homeomorphisms $\theta'_{p,i}: U_p \rightarrow U_{p_i}$ respecting f and $o(i)$ -close to θ_i . □

To conclude the proof of Theorem 7.8 observe that all the maps f_p above can be chosen to respect f on P and h on K . □

Remark 7.10. It's instructive to point out precisely what's needed to make the proof of the Stability Theorem work in the Lipschitz category.

1. One needs to generalize the deformation of homeomorphisms results of Siebenmann [Sie72] used in the proof of Theorem 4.6 to Lipschitz category. This is probably possible and in fact it is already known in case of Lipschitz manifolds by [Sul79].

2. Another (probably quite difficult) point is to generalize Perelman's Local Fibration Theorem 6.8 to Lipschitz category. To do this one needs to show that under the assumptions of Lemma 6.9, the homeomorphism of the "tubular" neighborhood of S to the product of S and the cone over F can be made to be bi-Lipschitz. The basic (and probably the most important) case of this would be to show that if h is a proper strictly concave function on an Alexandrov space with a unique maximum at a point p then the superlevel set $\{h \geq h(p) - \epsilon\}$ is bi-Lipschitz homeomorphic to the cone over $\{h = h(p) - \epsilon\}$. This is related to 1. and could possibly be proved using an appropriate generalization of Siebenmann's results.

Similar discussion applies to the case of DC rather than Lipschitz stability.

The Stability theorem has a natural generalization to the case of pointed Gromov-Hausdorff convergence. The following application, saying that the stability homeomorphisms can be constructed on arbitrary large compact subsets, seems to be the most useful. For simplicity we only state the unparameterized version.

Theorem 7.11. *Let $(X_i^n, p_i) \xrightarrow{i \rightarrow \infty}^{G-H} (X^n, p)$ where $\text{curv} X_i \geq \kappa$ for all i . Let $R > 0, \epsilon > 0$ and let $\varphi_i: B(p, R + \epsilon) \rightarrow B(p_i, R + \epsilon)$ be $o(i|R)$ -Hausdorff approximations. Then for all large i there exist open embedding $\psi_i: B(p, R + \epsilon/2) \rightarrow X_i$ which are $o(i|R, \epsilon)$ -close to φ_i and such that $\psi_i(B(p, R + \epsilon/2)) \supset B(p_i, R)$ for all large i .*

Proof. Put $U = B(p, R), V = B(p, R + \epsilon/2), W = B(p, R + \epsilon)$. Recall, that for any point $x \in X$ there is $r_x > 0$ such that $d(\cdot, x)$ has no critical points in $B(p, 2r_x) \setminus \{p\}$. Cover \bar{W} by finitely many such balls $B(p_\alpha, r_\alpha)$ with all $r_\alpha < \epsilon/4$. Let $p_{\alpha, i} = \varphi_i(p_\alpha)$. By Stability Theorem 7.8, for all large i and all α there exist homeomorphisms $\varphi_{\alpha, i}: B(p_\alpha, r_\alpha) \rightarrow B(p_{\alpha, i}, r_\alpha)$ which are $o(i|R, \epsilon)$ -close to φ_i . Now the statement of the theorem follows by the direct application of Theorem 4.11. □

Remark 7.12. In the proof of Theorem 7.11 we could not apply the stability theorem directly to $\bar{B}(p, R)$ because in general we have no information on the existence of critical points of $d(\cdot, p)$ outside a small ball around p . In particular, $\bar{B}(p, R)$ need not be an MCS-space or be \varkappa -connected for any large R .

Remark 7.13. It follows from the stability theorem that for any given $n \in \mathbb{Z}_+, k \in \mathbb{R}, D > 0, v > 0$ there exists $\epsilon > 0$ such that any X and Y in $\mathfrak{Aler}^n(D, \kappa, v)$ with $d_{G-H}(X, Y) \leq \epsilon$ are homeomorphic. It's interesting to see if one can give an explicit estimate on ϵ in terms of n, k, D, v .

8. FINITENESS OF SUBMERSIONS

The following generalization of the Grove-Petersen-Wu finiteness theorem was proved by K. Tapp [Tap02, Theorem 2]:

Theorem 8.1. *Given $n, k \in \mathbb{Z}_+, v, D, \lambda \in \mathbb{R}$ with $k \geq 4$, there are at most finitely many topologically equivalence classes of bundles in the set of Riemannian submersions $M^{n+k} \rightarrow B^n$ satisfying $\text{vol}(B) \geq v, |\text{sec}(B)| \leq \lambda; \text{vol}(M) \geq v, \text{diam}(M) \leq D, \text{sec}(M) \geq -\lambda$.*

Here two submersions $\pi_i: M_i \rightarrow B_i$ ($i = 1, 2$) are called topologically equivalent if there exist homeomorphisms $\varphi: M_1 \rightarrow M_2, f: B_1 \rightarrow B_2$ such that $\pi_2 \circ \varphi = f \circ \pi_1$.

This definition can be naturally extended to the class of submetries of Alexandrov spaces. Recall that a map $f: X \rightarrow Y$ is called a submetry if $f(B(x, r)) = B(f(x), r)$ for any $x \in X, r > 0$ (i.e if f is both 1-Lipschitz and 1-co-Lipschitz).

It is obvious that a Riemannian submersion between complete Riemannian manifolds is a submetry. Moreover, converse is also true according to [BG00]. It is also clear that if a compact group G acts on a Riemannian manifold M by isometries, then the projection $M \rightarrow M/G$ is a submetry.

Submetries enjoy many properties of Riemannian submersions. In particular, one can talk about horizontal and vertical tangent vectors and curves. Also, it's easy to see [BGP92] that submetries increase Alexandrov curvature, that is, if $\text{curv}X \geq \kappa$ and $f: X \rightarrow Y$ is a submetry, then $\text{curv}Y \geq \kappa$. For more basic information on submetries see [Lyt02].

Suppose $\pi: X \rightarrow B$ is a submetry between compact Alexandrov spaces. It trivially follows from the definition that if $f: B \rightarrow \mathbb{R}^k$ is admissible then $f \circ \pi$ is admissible on M . Moreover, f is regular at $p \in B$ iff $f \circ \pi$ is regular at any $y \in \pi^{-1}(p)$. In particular, if B is a Riemannian manifold (or more generally, if B is everywhere n -strained) then π is a fiber bundle. Thus the above notion of equivalence of submersions naturally extends to submetries.

Theorem 8.1 generalizes a theorem of J. Y. Wu, [Wu96] which proved the same result under a strong extra assumption that fibers of the submersions are totally geodesic. The proof of Theorem 8.1 relies on the proof of Wu's theorem which just as the proof of Grove-Petersen-Wu finiteness Theorem 1.2 uses techniques of controlled homotopy theory. This explains the assumption $k \geq 4$ in Theorem 8.1. However, this assumption is, in fact, unnecessary as this result follows from the Parameterized Stability Theorem which does not require any dimensional restrictions.

Theorem 8.2. *Given $k \in \mathbb{Z}_+, V, D, \lambda, \kappa \in \mathbb{R}$, there are at most finitely many equivalence classes of submetries $X^{n+k} \rightarrow B^n$ where $X^{n+k} \in \mathfrak{Aler}^{n+k}(D, \kappa, v)$ and B^n is a closed Riemannian manifold satisfying $\text{vol}(B) \geq V, |\text{sec}(B)| \leq \lambda$.*

Proof. We first give a proof in case of fixed B .

Let $\pi: X \rightarrow B$ be a submetry where X is an Alexandrov space of curvature bounded below.

Let $p \in B$ be any point. Choose $n + 1$ unit vectors $v_0, \dots, v_n \in T_p B$ with pairwise angles bigger than $\pi/2$. Then for all sufficiently small $R > 0$ the points $p_i = \exp_p(Rv_i)$ define an admissible map $f: B \rightarrow \mathbb{R}^n$ given by $y \mapsto (d(y, p_1), \dots, d(y, p_n))$. This map is obviously regular on $B(p, r)$ for $r \ll R$ and it gives a bi-Lipschitz open embedding (in fact a smooth one) $f: B(p, R) \rightarrow \mathbb{R}^n$. Let $F_i = \pi^{-1}(p_i)$. Let $\tilde{f}: X \rightarrow \mathbb{R}^n$ be given by $\tilde{f}(x) = (d(x, F_1), \dots, d(x, F_n))$.

Since π is a submetry we obviously have that $d(x, F_i) = d(\pi(x), p_i)$ for any i and any $x \in X$. Therefore $\tilde{f} \equiv f \circ \pi$. It is also obvious that \tilde{f} is regular at $x \in X$ iff f is regular at $\pi(x)$.

In particular, \tilde{f} is regular on the r -neighborhood of $\pi^{-1}(\pi(p))$. Thus, up to a bi-Lipschitz change of coordinates on the target, when restricted to $U_r(\pi^{-1}(\pi(p)))$, we can write π as a proper regular map \tilde{f} to \mathbb{R}^n . Let's cover B by finitely many coordinate neighborhoods $U_\alpha = B(p_\alpha, r_\alpha)$ as above and let $f_\alpha: U_\alpha \rightarrow \mathbb{R}^n$ be the corresponding coordinate projections. Since all f_α are bijections we obviously have that for any $x, y \in \pi^{-1}(U_\alpha \cap U_b)$, $\tilde{f}_\alpha(x) = \tilde{f}_\alpha(y)$ iff $\tilde{f}_\beta(x) = \tilde{f}_\beta(y)$.

Therefore Parameterized Stability Theorem 7.8 can be easily amended to include the case when a framing on X respects a submetry to a fixed manifold as all the arguments can be made local on B where instead of the submetry π one can work with regular maps \tilde{f}_α .

The case of variable base easily follows given the fact that by Cheeger-Gromov compactness the class of manifolds $\{B^n \mid \text{vol}(B) \geq V, |\text{sec}(B)| \leq \lambda, \text{diam}(B) \leq D\}$ is precompact in Lipschitz topology and its limit points are $C^{1,\alpha}$ -Riemannian manifolds (see e.g. [GW88]). \square

Remark 8.3. It's interesting to see whether Theorem 8.2 remains true if one removes the assumption about the uniform upper bound on the curvature of B .

9. STABILITY WITH EXTREMAL SUBSETS

The results proved in this section are new.

The notion of an extremal subset in an Alexandrov space was introduced in [PP93].

Definition 9.1. *A closed subset E in an Alexandrov space X is called extremal if for any $q \in X \setminus E$ and $f = d(\cdot, E)$ the following holds:*

If $p \in E$ is a point of local minimum of $f|_E$ then it's a critical point of maximum type of f on X , i.e.

$$df_p(\xi) \leq 0 \text{ for any } \xi \in \Sigma_p X$$

Alternatively, it was shown in [Pet06] that E is extremal iff it's invariant under gradient flows of all semiconcave functions on X .

An extremal subset is called *primitive* if it doesn't contain any proper extremal subsets with nonempty relative interiors.

We refer to [PP93, Pet06] for basic properties of extremal subsets. It is easy to see [PP93] that closures of topological strata in an Alexandrov space X are extremal. Therefore stratification into extremal subsets can be considered as a geometric refinement of the topological stratification of X .

It is of course obvious that a homeomorphism between two Alexandrov spaces has to preserve topological strata.

The goal of this section is to generalize the Stability Theorem by showing that the stability homeomorphisms can be chosen to preserve extremal subsets. Namely we will prove the following

Theorem 9.2 (Relative Stability Theorem). *Let $X_i \xrightarrow[i \rightarrow \infty]{G-H} X^n$ be a noncollapsing sequence of compact Alexandrov spaces in $\mathbf{Atr}^n(D, \kappa)$. Let $\theta_i: X \rightarrow X_i$ be a sequence of $o(i)$ -Hausdorff approximations. Let $E_i \subset X_i$ be a sequence of extremal subsets converging to an extremal subset E in X . Then for all large i there exist homeomorphisms $\theta'_i: (X, E) \rightarrow (X_i, E_i)$, $o(i)$ -close to θ_i .*

In order to prove this theorem we'll need to generalize all the machinery used in the proof of the regular Stability Theorem to its relative version respecting extremal subsets. This is fairly straightforward and only minor modifications of the proofs are required.

In particular we'll have to prove the relative version of Local Fibration Theorem 6.8. Along the way we'll obtain some new topological information about the way a general extremal subset is embedded into an ambient Alexandrov space.

It was shown in [PP93] that just as Alexandrov spaces, extremal subsets are naturally stratified in the sense of the following definition.

Definition 9.3. *A metrizable space X is called an \widetilde{MSC} -space of dimension $\leq n$ if every point $x \in X$ has a neighborhood pointed homeomorphic to an open cone over a compact \widetilde{MCS} -space of dimension $\leq n - 1$. As for \widetilde{MCS} -spaces we assume the empty set to be the unique \widetilde{MCS} -space of $\dim \leq -1$.*

We will also need a relative version of the above definition.

Definition 9.4. *Let X be an \widetilde{MSC} -space of $\dim \leq n$. A subset $E \subset X$ is called a stratified subspace of X of dimension $\leq k$ if every $x \in X$ has a pointed neighborhood (U, p) such that $(U, U \cap E, p)$ is homeomorphic to $(K\Sigma, K\Sigma', o)$ where Σ is a compact \widetilde{MSC} -space of $\dim \leq n - 1$ and $\Sigma' \subset \Sigma$ is a compact stratified subspace of Σ of dimension $\leq k - 1$. As usual, the only stratified subspace of $\dim \leq -1$ is the empty set.*

It is obvious from the definition that a stratified subspace in X of $\dim \leq k$ is an \widetilde{MCS} -space of dimension $\leq k$.

Remark 9.5. It is easy to see that a connected \widetilde{MSC} -space is an \widetilde{MCS} -space iff its local topological dimension is constant. It was shown in [PP93] that a primitive extremal subset is equal to the closure of its top dimensional strata and therefore is an \widetilde{MCS} -space by above.

In the process of proving the Relative Stability Theorem we'll obtain the following result which clarifies the relative topology of extremal subsets with respect to their ambient spaces.

Theorem 9.6 (Relative Stratification Theorem). *Let X be an Alexandrov space and let $E \subset X$ be an extremal subset. Then E is a stratified subspace of X .*

Just as in the non-relative case, this theorem is a Corollary of the following local fibration theorem applied to the natural map $X \rightarrow \mathbb{R}^0$.

Theorem 9.7 (Relative Local Fibration Theorem). *Let $f: X \rightarrow \mathbb{R}^k$ be regular at $p \in E$ where $E \subset X$ is an extremal subset. Then there exists an open neighborhood U of p , an \widetilde{MCS} -space A , a stratified subspace $B \subset A$ and a homeomorphism $\varphi: (U, E \cap U) \rightarrow (A, B) \times \mathbb{R}^k$ such that $\pi_2 \circ \varphi = f$.*

It was shown in [PP93] by Perelman and Petrunin that the intrinsic metric on an extremal subset of an Alexandrov space is locally bi-Lipschitz to the ambient metric. On closer examination their proof actually gives the following somewhat stronger statement which we'll need for the proof of the relative stability theorem:

Lemma 9.8. *There exists $\epsilon = \epsilon(n, D, \kappa, v) > 0$ such that if $X \in \mathbf{Alex}^n(D, \kappa, v)$ and $E \subset X$ is extremal, then for any $p, q \in E$ with $d(p, q) \leq \epsilon$ there exists a curve in E connecting p and q of length $\leq \epsilon^{-1}d(p, q)$.*

Proof. Because the argument is very easy we give it here for reader's convenience. It is well-known (see [PP93] or [GP88]) that for the class $\mathbf{Alex}^n(D, \kappa, v)$ there exists an $\epsilon > 0$ such that the following holds:

$$(9.1) \quad \text{If } X \in \mathbf{Alex}^n(D, \kappa, v) \text{ and } p, q \in X \text{ with } d(p, q) < \epsilon^2 \text{ then} \\ |\nabla_p d(\cdot, q)| > \epsilon \text{ or } |\nabla_q d(\cdot, p)| > \epsilon$$

Suppose the first alternative holds. Then there exists x near p such that $\angle xpq \geq \pi/2 + \epsilon$. Then $\nabla_p d(\cdot, x)$ is polar to \uparrow_p^x so that $\angle \nabla_p d(\cdot, x) \uparrow_p^q \leq \pi/2 - \epsilon$. This means that moving p along the gradient flow of $d(\cdot, x)$ decreases $d(p, q)$ in the first order (with the derivative at 0 at least ϵ). Since E is extremal, the flow through p remains in E .

Now a standard argument shows that we can construct a curve in E connecting p and q of length $\leq \epsilon^{-1}d(p, q)$. □

We will need the following generalization of Lemma 6.7 proved in [PP93]:

Lemma 9.9. [PP93, Pet06] *Suppose Σ^{n-1} has $\text{curv} \geq 1$ and let $f_0, \dots, f_k: \Sigma \rightarrow \mathbb{R}$ be functions of class DER such that $\epsilon = \min_{i \neq j} -\langle f_i, f_j \rangle > 0$. Let $E \subset \Sigma$ be an extremal subset. Then*

- (1) *There exists $w \in E$ such that $f_i(w) > \epsilon$ for all $i \neq 0$.*
- (2) *There exists $v \in E$ such that $f_0(v) > \epsilon, f_1(v) < -\epsilon$ and $f_i(v) = 0$ for $i = 2, \dots, k$.*

Just as in the case of regular functions on Alexandrov spaces, this lemma implies that if $f: X \rightarrow \mathbb{R}^k$ is regular at $p \in E$ where $E \subset X$ is extremal then $f|_E$ is co-lipschitz near p .

The main geometric ingredient in the proof of the Relative Local Fibration Theorem and the Relative Stratification Theorem is the following relative analogue of Lemma 6.9.

Lemma 9.10. *Let $p \in E$ be a regular point of $f: X \rightarrow \mathbb{R}^k$ where E is an extremal subset of X . Suppose f is incompressible at $p \in E$.*

Then there exists an admissible function $h: X \rightarrow \mathbb{R}$ with the following properties

- (i) $h(p) = 0$.
- (ii) h is strictly concave on $B(p, R)$ for some $R > 0$.
- (iii) There are $r > 0, A > 0$ such that $h < A$ on $f^{-1}(\bar{I}^k(f(p), r))$ and $f^{-1}(\bar{I}^k(f(p), r)) \cap \{h \geq -A\}$ is compact in $B(p, R)$.
- (iv) h has a unique maximum in $B(p, R) \cap f^{-1}(v)$ for all $v \in \bar{I}^k(f(p), r)$. Let S denote the set of such maximum points.
- (v) (f, h) is regular on $f^{-1}(\bar{I}^k(f(p), r)) \cap B(p, R) \setminus S$.
- (vi) $S \subset E$.

Proof. The proof is identical to the proof of Lemma 6.9 except for part (vi) which is new. Let x be a point of max of h on $E \cap f^{-1}(v)$. Let z be the unique point of maximum of h on $f^{-1}(v)$. If $x \neq z$ then (f, h) is regular at x by Lemma 6.9. However, by Lemma 9.9, a

point on E is regular for $F: X \rightarrow \mathbb{R}^m$ iff it's regular for $F|_E$. Thus, $(f, h)|_E$ is regular at x . Therefore, $(f, h)|_E$ is co-lipschitz near x and hence x is not a point of maximum of h on $E \cap f^{-1}(v)$. \square

This Lemma easily implies that if $f: X \rightarrow \mathbb{R}^k$ is regular at $p \in E$ where $E \subset X$ is extremal then the local dimension of E near p is $\geq k$ and the equality is only possible if $F|_E$ is locally bi-Lipschitz near p (cf. Corollary 6.10).

Lemma 9.10 also yields the relative local fibration theorem in exactly the same way as Lemma 6.9 yields the absolute local fibration theorem.

Proof of Theorem 9.7. Since the proof is almost identical to the proof of the fibration theorem in [Per93] we only give a sketch.

We argue by reverse induction in k . Since the base of induction is clear we only have to consider the induction step from $k + 1 \leq n$ to k . Let $f: X \rightarrow \mathbb{R}^k$ be regular at $p \in E$ where $E \subset X$ is extremal. If f is complementable at p the statement follows by induction assumption. Suppose f is incomplementable at p . Let h be the function provided by Lemma 9.10. Suppose for simplicity that h is identically zero on S . Let $U = f^{-1}(\bar{I}^k(f(p), r)) \cap \{-A < h < 0\} \cap B(p, R)$ and $W = f^{-1}(\bar{I}^k(f(p), r)) \cap \{-A < h \leq 0\} \cap B(p, R)$

Then $U = W \setminus S$ and (f, h) is regular on U . Therefore the relative local fibration theorem holds for (f, h) on U by induction assumption.

By Lemma 9.10, (f, h) is proper on U and hence, $(f, h): (U, U \cap E) \rightarrow \bar{I}^k(f(p), r) \times (-A, 0)$ is a relative bundle map by [Sie72, 6.10]. This means that $(U, U \cap E)$ is homeomorphic to $(F, B) \times \bar{I}^k(f(p), r) \times (-A, 0)$ respecting (f, h) where F is an MCS-space of $\dim = n - k - 1$ and B is a stratified subspace in F . By Lemma 9.10, we can extend this homeomorphism to a homeomorphism $(W, W \cap U) \rightarrow (KF, KB) \times \bar{I}^k(f(p), r)$ which proves the induction step.

The general case when h is not constant on S is handled in exactly the same way as in [Per93] and the proof of Key Lemma 7.9 by constructing an auxiliary function \tilde{h} obtained by shifting h by constants on each of the fiber of f to make it identically zero on S . \square

Before we can start with the proof of the Relative Stability Theorem we first need to observe that by [Sie72] the corresponding version of Theorem 4.4 and Strong Gluing Theorem 4.10 hold in relative category for pairs of MCS spaces and their stratified subspaces.

The relative version of Theorem 4.4 follows from [Sie72, Complement 6.10 to Union Lemma 6.9] by the same argument as in the proof of the absolute version of Theorem 4.4 given by [Sie72, Corollary 6.14].

The relative version of the Strong Gluing Theorem still follows from the same deformation of homeomorphism result [Sie72, Theorem 6.1] which also covers relative homeomorphisms. Here we'll need the following definition

Definition 9.11. *A pair of metric spaces (X, E) is called \varkappa -connected if both X and E (taken with the restricted ambient metric) are \varkappa -connected.*

Let us state the Relative Gluing Theorem. For simplicity we only state the unparameterized version.

Relative Gluing Theorem 9.12. *Let (X, E) be a stratified pair, $\{U_\alpha\}_{\alpha \in \mathfrak{A}}$ be a finite covering of X . Given a function \varkappa_0 , there exists $\varkappa = \varkappa((X, E), \{U_\alpha\}_{\alpha \in \mathfrak{A}}, \varkappa_0)$ such that the following holds:*

Given a \varkappa_0 -connected stratified pair (\tilde{X}, \tilde{E}) , an open cover of \tilde{X} $\{\tilde{U}_\alpha\}_{\alpha \in \mathfrak{A}}$, a δ -Hausdorff approximation $\varphi: (X, E) \rightarrow (\tilde{X}, \tilde{E})$ and a family of homeomorphisms $\varphi_\alpha: (U_\alpha, U_\alpha \cap E) \rightarrow (\tilde{U}_\alpha, \tilde{U}_\alpha \cap \tilde{E})$, δ -close to φ ,

then there exists a homeomorphism $\bar{\varphi}: (X, E) \rightarrow (\tilde{X}, \tilde{E})$, $\varkappa(\delta)$ -close to φ .

Observe that under the assumptions of the Relative Stability Theorem, all the elements of the sequence (X_i, E_i) and the limit (X, E) are \varkappa -connected by Lemma 9.8.

Furthermore, Lemma 6.15 and Remark 6.16 hold for regular level sets of admissible functions on extremal subsets. The proof is exactly the same as the proof of Lemma 6.15 modulo Lemma 9.8 and the fact that extremal subsets are invariant under all gradient flows.

We are now ready to prove the Relative Stability Theorem.

Proof of Theorem 9.2. The proof proceeds by reverse induction on the framing and is, in fact, *exactly* the same as the proof of the usual stability theorem except we make all the arguments relative. Everywhere in the proof substitute U (with various subindices) by $(U, U \cap E)$. In the proof of the relative version of Key Lemma 7.9, use Lemma 9.10 instead of Lemma 6.9 whenever necessary. Also, use the Relative Local Fibration Theorem instead of Local Fibration Theorem 4.4 and the Relative Gluing Theorem instead of the Gluing Theorem whenever called for.

□

Remark 9.13. The relative stability theorem trivially implies the following hitherto unobserved fact. Under the assumptions of the Relative Stability Theorem, $E_i \xrightarrow{i \rightarrow \infty} E$ without collapse. Then $\dim E = \dim E_i$ for all large i .

A fairly simple direct proof of this statement can be given without using the Relative Stability Theorem. However, we chose not to present it here because this indeed obviously follows from the Relative Stability Theorem.

REFERENCES

- [BG00] V. N. Berestovskii and L. Guijarro. A metric characterization of Riemannian submersions. *Ann. Global Anal. Geom.*, 18(6):577–588, 2000.
- [BGP92] Yu. Burago, M. Gromov, and G. Perel'man. A. D. Aleksandrov spaces with curvatures bounded below. *Uspekhi Mat. Nauk*, 47(2(284)):3–51, 222, 1992.
- [CF79] T. A. Chapman and S. Ferry. Approximating homotopy equivalences by homeomorphisms. *Amer. J. Math.*, 101(3):583–607, 1979.
- [Dav00] J. F. Davis. Manifold aspects of the Novikov conjecture. In *Surveys on surgery theory, Vol. 1*, volume 145 of *Ann. of Math. Stud.*, pages 195–224. Princeton Univ. Press, Princeton, NJ, 2000.
- [Fre82] M. H. Freedman. The topology of four-dimensional manifolds. *J. Differential Geom.*, 17(3):357–453, 1982.
- [GP88] K. Grove and P. Petersen. Bounding homotopy types by geometry. *Ann. of Math. (2)*, 128(1):195–206, 1988.

- [GPW91] K. Grove, P. Peterson, and J.-Y. Wu. Geometric finiteness theorems via controlled topology. *Invent. Math.*, 99:205–213, 1991.
- [GW88] R. E. Greene and H. Wu. Lipschitz convergence of Riemannian manifolds. *Pacific J. Math.*, 131(1):119–141, 1988.
- [GW97] K. Grove and F. Wilhelm. Metric constraints on exotic spheres via Alexandrov geometry. *J. Reine Angew. Math.*, 487:201–217, 1997.
- [Jak88] W. Jakobsche. Approximating homotopy equivalences of 3-manifolds by homeomorphisms. *Fund. Math.*, 130(3):157–168, 1988.
- [Kap02] V. Kapovitch. Regularity of limits of noncollapsing sequences of manifolds. *Geom. Funct. Anal.*, 12(1):121–137, 2002.
- [KPT05] V. Kapovitch, A. Petrunin, and W. Tuschmann. Nilpotency, almost nonnegative curvature and the gradient push. to appear in *Annals of Mathematics*, <http://arxiv.org/abs/math.DG/0506273>, 2005.
- [Kwu64] K.W. Kwun. Uniqueness of the open cone neighborhood. *Proc. Amer. Math. Soc.*, 15:476–479, 1964.
- [LN] A. Lytchak and K. Nagano. Topological regularity of spaces with curvature bounded above. in preparation.
- [Lyt02] A. Lytchak. *Allgemeine Theorie der Submetrien und verwandte mathematische Probleme*. Bonner Mathematische Schriften [Bonn Mathematical Publications], 347. Universität Bonn Mathematisches Institut, Bonn, 2002. Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, 2001.
- [Per91] G. Perelman. Alexandrov spaces with curvatures bounded from below II. preprint, 1991.
- [Per93] G. Perelman. Elements of Morse theory on Aleksandrov spaces. *St. Petersburg. Math. J.*, 5(1):205–213, 1993.
- [Per95] G. Perelman. DC structure on Alexandrov space with curvature bounded below. preprint, <http://www.math.psu.edu/petrunin/papers/papers.html>, 1995.
- [Pet97] A. Petrunin. Applications of quasigeodesics and gradient curves. In *Grove, Karsten (ed.) et al., Comparison geometry. Cambridge: Cambridge University. Math. Sci. Res. Inst. Publ. 30, 203-219*. 1997.
- [Pet06] A. Petrunin. Semiconcave functions in alexandrov geometry. Anton’s paper in the same volume, 2006.
- [PP93] G. Perelman and A. M. Petrunin. Extremal subsets in Aleksandrov spaces and the generalized Liberman theorem. *Algebra i Analiz*, 5(1):242–256, 1993.
- [PP96] G. Perelman and A. Petrunin. Quasigeodesics and gradient curves in alexandrov spaces. preprint, <http://www.math.psu.edu/petrunin/papers/papers.html>, 1996.
- [Qui82] F. Quinn. Ends of maps. III. Dimensions 4 and 5. *J. Differential Geom.*, 17(3):503–521, 1982.
- [Sie72] L. C. Siebenmann. Deformation of homeomorphisms on stratified sets. I, II. *Comment. Math. Helv.*, 47:123–136; *ibid.* 47 (1972), 137–163, 1972.
- [Sul79] D. Sullivan. Hyperbolic geometry and homeomorphisms. In *Geometric topology (Proc. Georgia Topology Conf., Athens, Ga., 1977)*, pages 543–555. Academic Press, New York, 1979.
- [SY00] T. Shioya and T. Yamaguchi. Collapsing three-manifolds under a lower curvature bound. *J. Differential Geom.*, 56(1):1–66, 2000.
- [SY05] T. Shioya and T. Yamaguchi. Volume collapsed three-manifolds with a lower curvature bound. *Math. Ann.*, 333(1):131–155, 2005.
- [Tap02] K. Tapp. Finiteness theorems for submersions and souls. *Proc. Amer. Math. Soc.*, 130(6):1809–1817 (electronic), 2002.
- [Wu96] J.-Y. Wu. A parametrized geometric finiteness theorem. *Indiana Univ. Math. J.*, 45(2):511–528, 1996.

VITALI KAPOVITCH, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA, M5S2E4

E-mail address: vtk@math.toronto.edu