

PUTNAM PROBLEMS

INEQUALITIES

2007-B-2. Suppose that $f : [0, 1] \rightarrow \mathbf{R}$ has a continuous derivative and that $\int_0^1 f(x)dx = 0$. Prove that for every $\alpha \in (0, 1)$,

$$\left| \int_0^\alpha f(x)dx \right| \leq \frac{1}{8} \max_{0 \leq x \leq 1} |f'(x)| .$$

2004-A-6. Suppose that $f(x, y)$ is a continuous real-valued function on the unit square $0 \leq x \leq 1$, $0 \leq y \leq 1$. Show that

$$\int_0^1 \left(\int_0^1 f(x, y)dx \right)^2 dy + \int_0^1 \left(\int_0^1 f(x, y)dy \right)^2 dx \leq \left(\int_0^1 \int_0^1 f(x, y)dxdy \right)^2 + \int_0^1 \int_0^1 [f(x, y)]^2 dxdy .$$

2003-A-2. Let a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n be nonnegative real numbers. Show that

$$(a_1 a_2 \cdots a_n)^{1/n} + (b_1 b_2 \cdots b_n)^{1/n} \leq ((a_1 + b_1)(a_2 + b_2) \cdots (a_n + b_n))^{1/n} .$$

2003-A-3. Find the minimum value of

$$|\sin x + \cos x + \tan x + \cot x + \sec x + \csc x|$$

for real numbers x .

2003-A-4. Suppose that a, b, c, A, B, C are real numbers, $a \neq 0$ and $A \neq 0$, such that

$$|ax^2 + bx + c| \leq |Ax^2 + Bx + c|$$

for all real numbers x . Show that

$$|b^2 - 4ac| \leq |B^2 - 4AC| .$$

2003-B-2. Let n be a positive integer. Starting with the sequence $1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}$, form a new sequence of $n - 1$ entries $3/4, 5/12, \dots, (2n - 1)/2n(n - 1)$, by taking the averages of two consecutive entries in the first sequence. Repeat the averaging of neighbours of the second sequence to obtain a third sequence of $n - 2$ entries and continue until the final sequence consists of a single number x_n . Show that $x_n < 2/n$.

2003-B-6. Let $f(x)$ be a continuous real-valued function defined on the interval $[0, 1]$. Show that

$$\int_0^1 \int_0^1 |f(x) + f(y)|dxdy \geq \int_0^1 |f(x)|dx .$$

2002-B-3. Show that, for all integers $n > 1$,

$$\frac{1}{2ne} < \frac{1}{e} - \left(1 - \frac{1}{n}\right)^n < \frac{1}{ne} .$$

1999-B-4. Let f be a real function with a continuous third derivative such that $f(x)$, $f'(x)$, $f''(x)$, $f'''(x)$ are positive for all x . Suppose that $f'''(x) \leq f(x)$ for all x . Show that $f'(x) < 2f(x)$ for all x .

1998-B-1. Find the minimum value of

$$\frac{(x + 1/x)^6 - (x^6 + 1/x^6) - 2}{(x + 1/x)^3 + (x^3 + 1/x^3)}$$

for $x > 0$.

1998-B-2. Given a point (a, b) with $0 < b < a$, determine the minimum perimeter of a triangle with one vertex at (a, b) , one on the x -axis, and one on the line $y = x$. You may assume that a triangle of minimum perimeter exists.

1996-B-2. Show that for every positive integer n ,

$$\left(\frac{2n-1}{e}\right)^{\frac{2n-1}{2}} < 1 \cdot 3 \cdot 5 \cdots (2n-1) < \left(\frac{2n+1}{e}\right)^{\frac{2n+1}{2}} .$$

1996-B-3. Given that $\{x_1, x_2, \dots, x_n\} = \{1, 2, \dots, n\}$, find, with proof, the largest possible value, as a function of n (with $n \geq 2$), of

$$x_1x_2 + x_2x_3 + \cdots + x_{n-1}x_n + x_nx_1 .$$

1988-B-2. Prove or disprove: If x and y are real numbers with $y \geq 0$ and $y(y+1) \leq (x+1)^2$, then $y(y-1) \leq x^2$.