

# Projectivization, Welded Knots and Alekseev-Torossian


"An Algebraic Structure"

## The Categorification Speculative Paradigm.

- Every object in math is the Euler characteristic of a complex.
- Every operation in math lifts to an operation between complexes.
- Every identity in mathematics is true up to homotopy.

## The Projectivization Tentative Speculative Paradigm. Projectivization?

- Every graded algebraic structure in mathematics is the projectivization of a plain ("global") one.
- Every equation written in a graded algebraic structure is an equation for a homomorphic expansion, or for an automorphism of such.



- Has kinds, objects, operations, and maybe constants.
- Perhaps subject to some axioms.
- We always allow formal linear combinations.

## Defining proj O. The augmentation "ideal":

$$I = I_{\mathcal{O}} := \left\{ \text{formal differences of objects "of the same kind"} \right\}.$$

Then  $I^n := \left\{ \text{all outputs of algebraic expressions at least } n \text{ of whose inputs are in } I \right\}$ , and

$$\text{proj } \mathcal{O} := \bigoplus_{n \geq 0} I^n / I^{n+1}.$$

- Has same kinds and operations, but different objects and axioms.

## Knot Theory Anchors.

- $(\mathcal{O}/I^{n+1})^*$  is "type  $n$  invariants".
- $(I^n/I^{n+1})^*$  is "weight systems".
- $\text{proj } \mathcal{O}$  is  $\mathcal{A}$ , "chord diagrams".



## Warmup Examples.

- The projectivization of a group is a graded associative algebra.
- A quandle: a set  $Q$  with a binary op  $\wedge$  s.t.


$$1 \wedge x = 1, \quad x \wedge 1 = x \wedge x = x, \quad (x \wedge y) \wedge z = (x \wedge z) \wedge (y \wedge z).$$

(appetizers) (main)

$L := \text{proj } Q$  is a graded Lie algebra: set  $\bar{v} := (v-1)$  (these generate  $I$ !), feed  $1+\bar{x}, 1+\bar{y}, 1+\bar{z}$  in (main), and collect the surviving terms of lowest degree:

$$(\bar{x} \wedge \bar{y}) \wedge \bar{z} = (\bar{x} \wedge \bar{z}) \wedge \bar{y} + \bar{x} \wedge (\bar{y} \wedge \bar{z}).$$

An Expansion is  $Z: \mathcal{O} \rightarrow \text{proj } \mathcal{O}$  s.t.  $Z(I^n) \subset (\text{proj } \mathcal{O})_{\geq n}$  and  $Z_{I^n/I^{n+1}} = \text{Id}_{I^n/I^{n+1}}$  (A "universal finite type invariant"). In practice, it is hard to determine  $\text{proj } \mathcal{O}$ , but easy to guess a surjection  $\rho: \mathcal{A} \rightarrow \text{proj } \mathcal{O}$ . So find  $Z': \mathcal{O} \rightarrow \mathcal{A}$  with  $Z'(I^n) \subset \mathcal{A}_{\geq n}$  and  $Z'_{I^n/I^{n+1}} \circ \rho_n = \text{Id}_{\mathcal{A}_n}$ :



Can you make this diagram less confusing?

- $e(x+y) = e(x)e(y)$  in  $\mathbb{Q}[[x, y]]$ .
- The pentagon and hexagons in  $\mathcal{A}(\uparrow_{3,4})$ .
- The equations defining a QUEA, the work of Etingof and Kazhdan.

## Graded Equations Examples

- The Alekseev-Torossian equations in  $\mathcal{U}(\text{sder}_n)$  and  $\mathcal{U}(\text{tder}_n)$ .

sder  $\leftrightarrow$  tree-level  $\mathcal{A}$   
tder  $\leftrightarrow$  more

$$F \in \mathcal{U}(\text{tder}_2); \quad F^{-1}e(x+y)F = e(x)e(y) \iff F \in \text{Sol}_0$$

$$\Phi = \Phi_F := (F^{12,3})^{-1}(F^{1,2})^{-1}F^{23}F^{1,23} \in \mathcal{U}(\text{sder}_3)$$

$$\Phi^{1,2,3}\Phi^{1,23,4}\Phi^{2,3,4} = \Phi^{12,3,4}\Phi^{1,2,34} \quad \text{"the pentagon"}$$

$t = \frac{1}{2}(y, x) \in \text{sder}_2$  satisfies  $4T$  and  $r = (y, 0) \in \text{tder}_2$  satisfies  $6T$

$$R := e(r) \text{ satisfies Yang-Baxter: } R^{12}R^{13}R^{23} = R^{23}R^{13}R^{12}$$

$$\text{also } R^{12,3} = R^{13}R^{23} \text{ and } F^{23}R^{1,23}(F^{23})^{-1} = R^{12}R^{13}$$

$$\tau(F) := RF^{21}e(-t) \text{ is an involution, } \Phi_{\tau(F)} = (\Phi_F^{321})^{-1}$$

$$\text{Sol}_0^r := \{F: \tau(F) = F\} \text{ is non-empty; for } F \in \text{Sol}_0^r,$$

$$e(t^{13} + t^{23}) = \Phi^{213}e(t^{13})(\Phi^{231})^{-1}e(t^{23})\Phi^{321}$$

$$\text{and } e(t^{12} + t^{13}) = (\Phi^{132})^{-1}e(t^{13})\Phi^{312}e(t^{12})\Phi$$



Alekseev




Torossian

This is just a part of the Alekseev-Torossian work!

- Related to the Kashiwara-Vergne Conjecture! So What?
- Will likely lead to an explicit tree-level associator, a linear equation away from a 1-loop equation, two linear equations away from a 2-loop associator, etc.!
- A baby version of the QUEA equations; we may be on the right tracks!


## Knotted Trivalent Graphs

$$\mathcal{O}(\Delta) = \left\{ \text{trivalent graphs}, \dots \right\}$$



**Theorem.** KTG is generated by the unknotted  $\Delta$  and the Möbius band, with identifiable relations between them.

**Theorem.**  $Z(\Delta)$  is equivalent to an associator  $\Phi$ .



Algebraic Knot Theory

**Theorem.**  $\{\text{ribbon knots}\} \sim \{u\gamma: \gamma \in \mathcal{O}(\circ\circ), d\gamma = \bigcirc\bigcirc\}$ .




Hence an expansion for KTG may tell us about ribbon knots, knots of genus 5, boundary links, etc.

Homomorphic Expansions are expansions that intertwine the algebraic structure on  $\mathcal{O}$  and  $\text{proj } \mathcal{O}$ . They provide finite / combinatorial handles on global problems.




X-S. Lin

# Projectivization, Welded Knots and Alekseev-Torossian



## Circuit Algebras

- \* Have "circuits" with "ends"
- \* Can be wired arbitrarily.
- \* May have "relations" - de-Morgan, etc.




add F1  
 $wK = \text{Welded knots + links}$

## "Welded trivalent (framed) tangles" are a circuit algebra:


"Welded Braids" are due to Fenn, Rimanyi and Rourke

$WT = \langle \text{diagrams} \rangle / R123, R4 \text{ (for vertices), F1.}$   
Further operations: delete, unzip.

## Partial Dictionary.





The "Chord Diagrams" -  $\mathcal{A}_n^{wt}$ . As we did for quandles, substitute



## The "Jacobi Diagrams" - $\mathcal{A}_n^{cc}$ .


Theorem. (95%)  $\mathcal{A}_n^{wt}$  is  $\mathcal{A}_n^{cc}$  is  $U(\text{tder}_n)$ . (pretend wheels dit.)



## The Map $\alpha: \mathcal{A}_n^{tree} \rightarrow \mathcal{A}_n^{cc}$ :

Theorem. (90%)  $\alpha$  is an injection on  $\mathcal{A}_n^{tree} \cong U(\text{sder}_n)$ .

Furthermore, there is a simple characterization of  $\text{im } \alpha$ , so we can tell "an arrowless element" when we see it.



The pentagon and The hexagons Follow, with a minor twist, from the fact that we have an unzip behaved invariant of KTG's.

The Main Theorem. (80%/0%) F's in  $\text{Sol}_0^7$  are in a bijective correspondance with tree-level associators for ordinary paranthesized tangles (or ordinary knotted trivalent graphs) / with homomorphic expansions for knotted welded trivalent tangles.

Disclaimer. Orientations, rotation numbers, framings, the vertical direction and the cyclic symmetry of the vertex may still make everything uglier. I hope not.

"God created the knots, all else in topology is the work of mortals"

Leopold Kronecker (paraphrased)

Visit! Edit!

The Knot Atlas - Anyone Can Edit

<http://katlas.org>