

$$\begin{aligned}
dm_{i,j \rightarrow k} &\rightarrow a_k (\alpha_i + \alpha_j) + b_k (\beta_i + \beta_j) + y_k \eta_i + \frac{y_k \eta_j}{\alpha_i} + \frac{x_k \xi_i}{\alpha_j} + \eta_j \xi_i - \\
& B_k \eta_j \xi_i + \frac{1}{4 \alpha_i \alpha_j} \in \left(2 y_k \eta_j (2 x_k \xi_i + \alpha_j (-2 \beta_i + (1 - 3 B_k) \eta_j \xi_i)) + \right. \\
& \quad \alpha_i \xi_i (x_k (-4 \beta_j + 2 (1 - 3 B_k) \eta_j \xi_i) + \\
& \quad \left. \alpha_j \eta_j (4 a_k B_k + (1 - 4 B_k + 3 B_k^2) \eta_j \xi_i) \right) + x_k \xi_j \\
d\Delta_{i \rightarrow j, k} &\rightarrow a_j \alpha_i + a_k \alpha_i + b_j \beta_i + b_k \beta_i + y_j \eta_i + B_j y_k \eta_i + \\
& x_j \xi_i + x_k \xi_i + \frac{1}{2} \in (B_j y_j y_k \eta_i^2 + x_k \xi_i (-2 a_j + x_j \xi_i)) \\
dS_i &\rightarrow -a_i \alpha_i - b_i \beta_i - \frac{\alpha_i (y_i \eta_i + (-\eta_i + B_i (x_i + \eta_i)) \xi_i)}{B_i} - \\
& \frac{1}{4 B_i^2} \in \alpha_i (\alpha_i \eta_i^2 (2 y_i^2 - 6 y_i \xi_i + 3 \xi_i^2) + B_i^2 \xi_i (4 a_i x_i + 2 x_i^2 \alpha_i \xi_i + \\
& \quad 2 x_i (2 \beta_i + \alpha_i \eta_i \xi_i) + \eta_i (-4 + 4 \beta_i + \alpha_i \eta_i \xi_i)) + \\
& \quad 2 B_i \eta_i (y_i (-2 + 2 \beta_i + 2 x_i \alpha_i \xi_i + \alpha_i \eta_i \xi_i) - \\
& \quad \xi_i (-2 + 2 a_i + 2 \beta_i + 3 x_i \alpha_i \xi_i + 2 \alpha_i \eta_i \xi_i)) \\
R_{i,j} &\rightarrow a_j b_i + x_j y_i - \frac{1}{4} \in x_j^2 y_i^2 \\
P_{i,j} &\rightarrow \alpha_j \beta_i + \eta_i \xi_j + \frac{1}{4} \in \eta_i^2 \xi_j^2
\end{aligned}$$

$$\begin{aligned}
E_{() \rightarrow (1)} &\left[\mathbf{0}, \mathbf{0}, \frac{B}{1 - B + B^2} + \right. \\
& \left. \frac{B (-B + 2 B^2 + 2 B^4 + a (-1 + B - B^3 + B^4) - 2 x y - B^3 (3 + 2 x y))}{(1 - B + B^2)^3} \in + \right. \\
& \left. \frac{1}{2 (1 - B + B^2)^5} \right. \\
& \left. B (4 B^8 + a^2 (1 - B + B^2)^2 (1 + B - 6 B^2 + B^3 + B^4) + 6 B^5 x^2 y^2 + \right. \\
& \quad 2 x y (-2 + 3 x y) - B^7 (11 + 4 x y) - 2 B^2 (1 + 6 x^2 y^2) - \\
& \quad 2 B^4 (1 - 2 x y + 6 x^2 y^2) + B (1 + 8 x y + 6 x^2 y^2) + \\
& \quad B^6 (6 + 8 x y + 6 x^2 y^2) + B^3 (4 + 4 x y + 30 x^2 y^2) + \\
& \quad 2 a (1 - B + B^2) (2 B^6 + 2 x y + 8 B^3 (1 + x y) - 5 B^2 (1 + 2 x y) - \\
& \quad \left. 2 B^5 (1 + 2 x y) - B^4 (7 + 2 x y) + B (2 + 4 x y)) \right) \in^2 + 0[\in]^3]
\end{aligned}$$

A Quantum Algebra Example.

Proto-Proposition^{†0} (with Jesse Frohlich and Roland van der Veen, near [Ma, Proposition 1.7.3]). Let H be a finite dimensional Hopf algebra and let $U = H^{*cop} \otimes H$ be its Drinfel'd double, with R -matrix $R \in H^* \otimes H \subset U \otimes U$. Write $R^{\dagger 1} = \sum \rho_a \otimes r_a$, and let $\langle \cdot | \cdot \rangle: H^* \otimes H \rightarrow \mathbb{F}$ be the duality pairing. Then the functional $\int \in U^*$ defined by

$$\int \phi \otimes x := \sum \langle \phi \rho_a^{\dagger 2} | x r_a^{\dagger 3} \rangle$$

is a right^{†4} integral in U^* . (Meaning $\Delta_{jk}^i // \int_j = \int_i // \epsilon_k$ in $\text{Hom}(U^{\otimes \{i\}} \rightarrow U^{\otimes \{k\}})$).

†0 A “proto-proposition” is something that will become a proposition once you figure out the correct statement. †1 Or did we want it to be $R // S_1^2$? Or $R // S_2^2$? †2 Or is it $\rho_a \phi$? †3 Or is it $r_a x$? †4 Or maybe “left”?

`inp = $\mathbb{E}_{() \rightarrow (1)}$ [3 a1 b1, 5 x1 y1, 1] // dmi,1→i;`


`Table[`

```

HL@TrueQ[
  (inp // (SYi→1,1,2,2 RR) // BM // AM // P1,2) dej =
  (inp //  $\Delta\Delta$  // (SYi→1,1,2,2 RR) // BM // AM // P1,2) ],
{ $\Delta\Delta$ , {d $\Delta_{i \rightarrow j, i}$ , d $\Delta_{i \rightarrow j, i}$ }}, {AM, {dm2,4→2, dm4,2→2}},
{BM, {dm1,3→1, dm3,1→1}},
{RR, {R3,4, R3,4 // dS3 // dS3, R3,4 // dS4 // dS4}}
] // MatrixForm
( (False False False) (False False True)
  (False False False) (False False False)
  (False False False) (False False False)
  (False False True) (False False False) )

```

A Knot Theory Example.



`$k = 2;`

`Simplify[`

```

R1,5 R6,2 R3,7 C4 Kink8 Kink9 Kink10 // dm1,2→1 // dm1,3→1 //
dm1,4→1 // dm1,5→1 // dm1,6→1 // dm1,7→1 // dm1,8→1 //
dm1,9→1 // dm1,10→1] / . v-1 → v

```

References.

- [BG] D. Bar-Natan and S. Garoufalidis, *On the Melvin-Morton-Rozansky conjecture*, Invent. Math. **125** (1996) 103–133.
- [BV] D. Bar-Natan and R. van der Veen, *A Polynomial Time Knot Polynomial*, arXiv:1708.04853.
- [Fa] L. Faddeev, *Modular Double of a Quantum Group*, arXiv:math/9912078.
- [GR] S. Garoufalidis and L. Rozansky, *The Loop Expansion of the Kontsevich Integral, the Null-Move, and S-Equivalence*, arXiv:math.GT/0003187.
- [LT] R. J. Lipton and R. E. Tarjan, *A Separator Theorem for Planar Graphs*, SIAM J. Appl. Math. **36-2** (1979) 177–189.
- [Ma] S. Majid, *Foundations of Quantum Group Theory*, Cambridge University Press, 1995.
- [MM] P. M. Melvin and H. R. Morton, *The coloured Jones function*, Commun. Math. Phys. **169** (1995) 501–520.
- [Ov] A. Overbay, *Perturbative Expansion of the Colored Jones Polynomial*, University of North Carolina PhD thesis, ωεβ/Ov.
- [Qu] C. Quesne, *Jackson’s q-Exponential as the Exponential of a Series*, arXiv:math-ph/0305003.
- [Ro1] L. Rozansky, *A contribution of the trivial flat connection to the Jones polynomial and Witten’s invariant of 3d manifolds, I*, Comm. Math. Phys. **175-2** (1996) 275–296, arXiv:hep-th/9401061.
- [Ro2] L. Rozansky, *The Universal R-Matrix, Burau Representation and the Melvin-Morton Expansion of the Colored Jones Polynomial*, Adv. Math. **134-1** (1998) 1–31, arXiv:q-alg/9604005.
- [Ro3] L. Rozansky, *A Universal U(1)-RCC Invariant of Links and Rationality Conjecture*, arXiv:math/0201139.
- [Za] D. Zagier, *The Dilogarithm Function*, in Cartier, Moussa, Julia, and Vanhove (eds) *Frontiers in Number Theory, Physics, and Geometry II*. Springer, Berlin, Heidelberg, and ωεβ/Za.