Simple 2-Knots.








Satoh

Dalvit
ωεβ/Dal



The Generators





The Double Inflation Procedure δ.

w-Knots.



A Big Open Problem. δ maps w-knots onto simple 2-knots. To what extent is it a bijection? What other relations are required? In other words, **find a simple description of simple 2-knots**.

The Full

2-Knot Story



Rewrites of IHX.



Riddles, in case you are bored.

- Can you find uncountably many distinct subsets $\{A_\alpha\}$ of $\mathbb{Z}$ such that whenever $\alpha \neq \beta$ either $A_\alpha \subset A_\beta$ or $A_\beta \subset A_\alpha$?
- Can you find uncountably many distinct subsets $\{B_\alpha\}$ of $\mathbb{Z}$ such that whenever $\alpha \neq \beta$ the intersection $B_\alpha \cap B_\beta$ is finite?

Question. Does it all extend to arbitrary 2-knots (not necessarily "simple")? To arbitrary codimension-2 knots?

BF Following [CR]. $A \in \Omega^1(M = \mathbb{R}^4, \mathfrak{g})$, $B \in \Omega^2(M, \mathfrak{g}^*)$,

$$S(A,B) := \int_M \langle B, F_A \rangle.$$


Rossi

With $\kappa\colon (S = \mathbb{R}^2) \to M$, $\beta \in \Omega^0(S,\mathfrak{g})$, $\alpha \in \Omega^1(S,\mathfrak{g}^*)$, set

$$\mathcal{O}(A,B,\kappa) := \int \mathcal{D}\beta\mathcal{D}\alpha \exp\left(\frac{i}{\hbar}\int_S \langle \beta, d_{\kappa^*A}\alpha + \kappa^* B\rangle\right).$$


Cattaneo

The BF Feynman Rules. For an edge e, let Φ_e be its direction, in S^3 or S^1. Let ω_3 and ω_1 be volume forms on S^3 and S_1. Then

$$Z_{BF} = \sum_{\substack{\text{diagrams}\\ D}} \frac{[D]}{|\mathrm{Aut}(D)|} \underbrace{\int_{\mathbb{R}^2}\cdots\int_{\mathbb{R}^2}}_{S\text{-vertices}} \underbrace{\int_{\mathbb{R}^4}\cdots\int_{\mathbb{R}^4}}_{M\text{-vertices}} \prod_{\substack{\text{red}\\ e\in D}} \Phi_e^*\omega_3 \prod_{\substack{\text{black}\\ e\in D}} \Phi_e^*\omega_1$$

(modulo some IHX-like relations). See also [Wa]



Issues. • Signs don't quite work out, and BF seems to reproduce only "half" of the wheels invariant on simple 2-knots.
• There are many more configuration space integrals than BF Feynman diagrams and than just trees and wheels.
• I don't know how to define / analyze "finite type" for general 2-knots.
• I don't know how to reduce Z_{BF} to combinatorics / algebra.

References.

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, ωεβ/KBH, arXiv:1308.1721.

[BND1] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I: W-Knots and the Alexander Polynomial*, ωεβ/WKO1, arXiv:1405.1956.

[BND2] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects II: Tangles and the Kashiwara-Vergne Problem*, ωεβ/WKO2, arXiv:1405.1955.

[BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.

[CS] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, Math. Surv. and Mono. **55**, Amer. Math. Soc., Providence 1998.

[CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, Commun. in Math. Phys. **256-3** (2005) 513–537, arXiv:math-ph/0210037.

[Fa] M. Farber, *Noncommutative Rational Functions and Boundary Links*, Math. Ann. **293** (1992) 543–568.

[Le] J. Levine, *A Factorization of the Conway Polynomial*, Comment. Math. Helv. **74** (1999) 27–53, arXiv:q-alg/9711007.

[Wa] T. Watanabe, *Configuration Space Integrals for Long n-Knots, the Alexander Polynomial and Knot Space Cohomology*, Alg. and Geom. Top. **7** (2007) 47–92, arXiv:math/0609742.

"God created the knots, all else in topology is the work of mortals."

Leopold Kronecker (modified)

www.katlas.org

Video and more at http://www.math.toronto.edu/~drorbn/Talks/Hamilton-1412/