



Warning. Conventions on this page change randomly from line to line.

$Z^{w/2}$. The GGA story is about $Z^{w/2}: \mathcal{K} \rightarrow \mathcal{A}^{w/2}$, defined on arrows a by $\pm a \mapsto \exp(\pm a)$:




Where the target space $\mathcal{A}^{w/2}$ is the space of unsigned arrow diagrams modulo




($Z^{w/2}$ is a reduction of the much-studied Z^w [BND, BN]).

The Euler Trick. How best do non-commutative algebra with exponentials? Logarithms are from hell as $e^f e^g = e^{\text{bch}(f,g)}$, but Euler's from heaven: Let E be the derivation $Ef := (\deg f)f (= xf')$, in $\mathbb{Q}[[x]]$ and let $\tilde{E}Z := Z^{-1}EZ (= x(\log Z)'$ in same). If $\deg x = 1$ then $\tilde{E}e^x = x$ and if $F = e^f$ and $G = e^g$, then $\tilde{E}(FG)$ is $(FG)^{-1}((EF)G + F(EG)) = G^{-1}(\tilde{E}F)G + \tilde{E}G = e^{-\text{ad } g}(\tilde{E}F) + \tilde{E}G$.

Scatter and Glow. Apply \tilde{E} to $Z(K)$. EZ is shown:




Tail scattering. The algebra $\mathbb{Q}[[b_i]]\langle a_{ij} \rangle$ modulo $[a_{ij}, a_{kl}] = 0$ (loc), $[a_{ij}, a_{ik}] = 0$ (TC), and $[a_{ik}, a_{jk}] = -[a_{ij}, a_{jk}] = b_j a_{ik} - b_i a_{jk}$ (CH and $\overrightarrow{4T}$), acts on $V = \mathbb{Q}[[b_i]]\langle x_i = a_{i\infty} \rangle$ by $[a_{ij}, x_i] = 0$, $[a_{ij}, x_j] = b_i x_j - b_j x_i$. Hence $e^{\text{ad } a_{ij}} x_i = x_i$, $e^{\text{ad } a_{ij}} x_j = e^{b_i} x_j + \frac{b_j}{b_i}(1 - e^{b_i})x_i$. Renaming $\bar{x}_i = x_i/b_i$, $T_i = e^{b_i}$, get $[e^{\text{ad } a_{ij}}]_{\bar{x}_i, \bar{x}_j} = \begin{pmatrix} 1 & 1 - T_i \\ 0 & T_i \end{pmatrix}$. Alternatively,



Linear Control Theory.

If $\begin{pmatrix} y \\ y_n \end{pmatrix} = \begin{pmatrix} \Xi & \phi \\ \theta & \alpha \end{pmatrix} \begin{pmatrix} x \\ x_n \end{pmatrix}$, and we further impose $x_n = y_n$, then $y = Bx$ where $B = \Xi + \frac{\phi\theta}{1 - \alpha}$. This fully explains the Gassner formulas and the GGA formula!



All that remains now is to replace TC by something more interesting: with $\epsilon^2 = 0$,

$$[a_{ij}, a_{ik}] = \epsilon(c_j a_{ik} - c_k a_{ij}).$$

Many further changes are also necessary, and the algebra is a lot more complicated and revolves around “quantization of Lie bialgebras” [EK, En]. But the spirit is right.

References.

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, œβ/KBH, arXiv:1308.1721.
 [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I, II, IV*, œβ/WKO1, œβ/WKO2, œβ/WKO4, arXiv:1405.1956, arXiv:1405.1955, arXiv:1511.05624.
 [BNG] D. Bar-Natan and S. Garoufalidis, *On the Melvin-Morton-Rozansky conjecture*, Invent. Math. **125** (1996) 103–133.
 [BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.
 [CT] D. Cimasoni and V. Turaev, *A Lagrangian Representation of Tangles*, Topology **44** (2005) 747–767, arXiv:math.GT/0406269.
 [En] B. Enriquez, *A Cohomological Construction of Quantization Functors of Lie Bialgebras*, Adv. in Math. **197-2** (2005) 430-479, arXiv:math/0212325.
 [EK] P. Etingof and D. Kazhdan, *Quantization of Lie Bialgebras, I*, Selecta Mathematica **2** (1996) 1–41, arXiv:q-alg/9506005.
 [GST] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjectures*, Geom. and Top. **14** (2010) 2305–2347, arXiv:1103.1601.
 [GPV] M. Goussarov, M. Polyak, and O. Viro, *Finite type invariants of classical and virtual knots*, Topology **39** (2000) 1045–1068, arXiv:math.GT/9810073.
 [KLW] P. Kirk, C. Livingston, and Z. Wang, *The Gassner Representation for String Links*, Comm. Cont. Math. **3** (2001) 87–136, arXiv:math/9806035.
 [LD] J. Y. Le Dimet, *Enlacements d’Intervalles et Représentation de Gassner*, Comment. Math. Helv. **67** (1992) 306–315.
 [MM] P. M. Melvin and H. R. Morton, *The coloured Jones function*, Commun. Math. Phys. **169** (1995) 501–520.
 [PV] M. Polyak and O. Viro, *Gauss Diagram Formulas for Vassiliev Invariants*, Inter. Math. Res. Notices **11** (1994) 445–453.
 [Ro] L. Rozansky, *A contribution of the trivial flat connection to the Jones polynomial and Witten’s invariant of 3d manifolds, I*, Comm. Math. Phys. **175-2** (1996) 275–296, arXiv:hep-th/9401061.