

The Main Course

$B^{(m)} = (\text{PaB}^{(m)}, \mathbf{S} : \text{PaB}^{(m)} \rightarrow \text{PaP}, d_i, s_i, \square, \sigma)$

same-skeleton linear combinations allowed

d_0 (crossing) = ... ; d_0 (cup) = ...

d_2 (cup) = ... ; d_2 (crossing) = ...

$a =$ (cup) = ... ; $\sigma =$ (crossing) = ...

\square (cup) = ... and \square (crossing) = ...

$C^{(m)} = (\text{PaCD}^{(m)}, \mathbf{S} : \text{PaCD}^{(m)} \rightarrow \text{PaP}, d_i, s_i, \square, \tilde{R})$

same-skeleton linear combinations allowed

d_2 (cup) = ... ; d_0 (cup) = ...


$a =$ (cup) = ... ; $x =$ (crossing) = ...

$\tilde{R} = X \exp \frac{H}{2}$

$\Phi \cdot \exp \left(\pm \frac{1}{2} t^{23} \right) \cdot (\Phi^{-1})^{132} \cdot \exp \left(\pm \frac{1}{2} t^{13} \right) \cdot \Phi^{312}$

$s_1 \Phi = s_2 \Phi = s_3 \Phi = 1$

$\square \Phi = \Phi \otimes \Phi$




$d_4 \Gamma \cdot d_2 \Gamma \cdot d_0 \Gamma = d_1 \Gamma \cdot d_3 \Gamma$

$1 = \Gamma \cdot (\Gamma^{-1})^{132} \cdot \Gamma^{312}$

$d_1 t^{12} = \Gamma \cdot (t^{23} \cdot (\Gamma^{-1})^{132} + (\Gamma^{-1})^{132} \cdot t^{13}) \cdot \Gamma^{312}$

$e^{\epsilon(t^{13} + t^{23})} = \Gamma \cdot e^{\epsilon t^{23}} \cdot (\Gamma^{-1})^{132} \cdot e^{\epsilon t^{13}} \cdot \Gamma^{312}$

 I have a nifty Free Lie calculator. Wanna play?

GT

GRT

Braids and the Grothendieck–Teichmüller Group, II