

THE KAROUBI ENVELOPE AND LEE'S DEGENERATION OF KHOVANOV HOMOLOGY

DROR BAR-NATAN AND SCOTT MORRISON

ABSTRACT. We give a simple proof of Lee's result from [5], that the dimension of the Lee variant of the Khovanov homology of an c -component link is 2^c , regardless of the number of crossings. Our method of proof is entirely local and hence we can state a Lee-type theorem for tangles as well as for knots and links. Our main tool is the "Karoubi envelope of the cobordism category", a certain enlargement of the cobordism category which is mild enough so that no information is lost yet strong enough to allow for some simplifications that are otherwise unavailable.

CONTENTS

1. Introduction	1
1.1. Is there anything left to do?	2
1.2. The plan	2
1.3. Acknowledgement	2
2. A quick review of the local Khovanov theory	3
3. A quick review of the Karoubi envelope	4
4. Red and green in Lee's theory	5
5. Alternately coloured smoothings and orienting components	6
References	8

1. INTRODUCTION

In a beautiful article [5], Eun Soo Lee introduced a second differential Φ on the Khovanov complex of a knot (or link) and showed that the resulting double complex has non-interesting homology. In a seemingly contradictory manner, this is a very interesting result — for this "degeneration" of the Lee theory is in itself an extra bit of information about the original Khovanov homology, masterfully used by Rasmussen [7] to define the aptly called "Rasmussen invariant" of a knot and to give a combinatorial proof of an old conjecture of Milnor.

Unfortunately Lee's proof of her degeneration result is a bit technical and inductive in nature.

The purpose of this note is to re-prove Lee's degeneration result in local terms, using tools in the spirit of [2]. Thus in addition to being a bit more conceptual, our methods work for tangles as well as for knots and links.

Another proof of Lee's result was found by S. Wehrli [8]. His proof is quick, short and elegant; the only small advantage of our proof over his is its locality and applicability to tangles.

Date: First edition: June 21, 2006. This edition: June 28, 2006.

1991 Mathematics Subject Classification. 57M25; 57M27; 18E05.

Key words and phrases. Categorification, Cobordism, Karoubi envelope, Jones Polynomial, Khovanov, Knot Invariants.

This work was partially supported by NSERC grant RGPIN 262178. Electronic versions: <http://www.math.toronto.edu/~drorbn/papers/Karoubi/> and <http://math.berkeley.edu/~scott/papers/karoubi.pdf> and arXiv:math.GT/0606542.

Let us sketch our results now; the relevant terminology (which closely follows [2]) will be quickly recalled in §2 below.

A “confluence” within a smoothing S of a tangle T is a pair of arc segments in S that correspond to a small neighbourhood of a crossing in T . Thus if T has n crossings, S will have n confluences; a crossing such as \times in T becomes a confluence such as \succ or \prec in S .

Definition 1.1. Let T be a tangle and let S be a smoothing of T . An alternate colouring of S is a colouring of the components of S with two colours (always “red” and “green” below), so that the two arc segments at every confluence of S are coloured with different colours.

Our main theorem is the following:

Theorem 1.2. *Within the appropriate category (see below), the Khovanov-Lee complex of a tangle T is homotopy equivalent to a complex with one generator for each alternately coloured smoothing of T and with vanishing differentials.*

Alternately coloured smoothings are easy to manage:

Proposition 1.3. *A c -component tangle T has exactly 2^c alternately coloured smoothings. These smoothings are in a bijective correspondence with the 2^c possible orientations of the c components of T .*

Together, Theorem 1.2 and Proposition 1.3 imply Lee’s result, that the Khovanov-Lee homology of an c -component link is 2^c dimensional.

What’s that “appropriate category” of Theorem 1.2? It is the category of complexes over the “Karoubi envelope” $\text{Kar}(\mathcal{C}ob_1)$ of the category $\mathcal{C}ob_1$ used to describe the Khovanov-Lee complex. It is perhaps the nicest gadget appearing in our note — it introduces more objects into $\mathcal{C}ob_1$ allowing for more opportunities to simplify complexes over $\mathcal{C}ob_1$, yet it does not introduce new morphisms between existing objects of $\mathcal{C}ob_1$, and hence no new homotopies or homotopy equivalences. Thus there is no loss of information in the passage from $\mathcal{C}ob_1$ to $\text{Kar}(\mathcal{C}ob_1)$.

1.1. Is there anything left to do? Plenty. Lee’s degeneration is critical to understanding Rasmussen [7]. We have a clean approach to Lee’s degeneration, and one may hope it will help improving our understanding of Rasmussen’s work. This still remains to be done. More specifically, the Khovanov-Lee homology is filtered and the definition of the Rasmussen invariant requires this filtration. Thus:


Problem 1.4. *Figure out how the filtration in the Khovanov-Lee homology interacts with everything down below.*

1.2. The plan. In §2 we quickly recall the relevant definitions of the Khovanov and Khovanov-Lee theories. Then in §3 we review the definition of the Karoubi envelope of a general category \mathcal{C} . §4 is the heart of the paper. In it we introduce the “red” and the “green” projections, the red-green splitting within $\text{Kar}(\mathcal{C}ob_1)$ of objects in $\mathcal{C}ob_1$ and prove Theorem 1.2. Finally, in §5 we prove Proposition 1.3.

1.3. Acknowledgement. We wish to thank S. Wehrli for drawing our attention to [8].

2. A QUICK REVIEW OF THE LOCAL KHOVANOV THEORY

Let us briefly recall the definition of the Khovanov complex for tangles, following [2]. Given an n -crossing tangle T with boundary ∂T , such as the 2-crossing tangle displayed on the right, one constructs an n -dimensional “cube” of smoothings and cobordisms between them (as illustrated on the right). This cube is then “flattened” to a “formal complex” $[T]_0$ in the additive category $Cob_0(\partial T)$ (denoted $Cob_{\bullet/l}^3(\partial T)$ in [1, 2]) whose objects are formally graded smoothings with boundary ∂T and whose morphisms are formal linear combinations of cobordisms whose tops and bottoms are smoothings and whose side boundaries are $I \times \partial T$, modulo some local relations. An overall height shift ensures that the ‘oriented smoothing’ appears in homological height 0.



The Khovanov complex $[T]_0$ is an object in the category $\text{Kom}(\text{Mat}(Cob_0(\partial T)))$ of complexes of formal direct sums of objects in $Cob_0(\partial T)$ and it is invariant up to homotopies.

For simplicity we are using as the basis to our story one of the simpler cobordism categories $Cob_0 := Cob_{\bullet/l}^3$ that appear in [2], rather than the most general one, $Cob_{/l}^3$. It is worthwhile to repeat here the local relations that appear in the definition of Cob_0 (see [2, §11.2]):

(2.1) $\begin{matrix} \text{circle with dashed line} = 0, & \text{circle with solid line and dot} = 1, & \text{rectangle with two dots} = 0, \\ \text{and } \text{cylinder with dashed line} = \text{cup with dot} & \text{cup} + \text{cup} = \text{cup with dot} \end{matrix}$

Also recall from [2, §5] that $[\cdot]_0$ is a planar algebra morphism. That is, if T_1 and T_2 are tangles and $D(T_1, T_2)$ denotes one of their side-by-side compositions (a side by side placement of T_1 and T_2 while joining some of their ends in a certain way prescribed by a planar arc diagram D), then $[D(T_1, T_2)]_0 = [T_1]_0 \otimes_D [T_2]_0$. Here, as in [2, §5], $[T_1]_0 \otimes_D [T_2]_0$ is the “tensor product” operation induced on formal complexes by the horizontal composition operation D .

The Lee variant of Khovanov homology, or the Khovanov-Lee complex $[T]_1$ of a tangle T , is constructed in exactly the same way as $[T]_0$; the only difference is that the zero in the third local relation in (2.1) is replaced with a one. Hence it is valued in the category of complexes over the category $Cob_1(\partial T)$ defined in exactly the same manner as $Cob_0(\partial T)$, except with the following collection of local relations (spot the one difference!):

(2.2) $\begin{matrix} \text{circle with dashed line} = 0, & \text{circle with solid line and dot} = 1, & \text{rectangle with two dots} = 1, \\ \text{and } \text{cylinder with dashed line} = \text{cup with dot} & \text{cup} + \text{cup} = \text{cup with dot} \end{matrix}$

While Cob_0 can be made into a graded category (see [2, §6]), the degree of $\text{rectangle with two dots}$ is non-zero, so setting it to 1 breaks the grading in Cob_1 . Otherwise the Khovanov theory and the Khovanov-Lee theories are completely parallel. In particular, the Khovanov-Lee complex is also an up-to-homotopy knot invariant and it is also a planar algebra morphism.

3. A QUICK REVIEW OF THE KAROUBI ENVELOPE

A *projection* is an endomorphism p satisfying $p^2 = p$. In many contexts in mathematics, if p is a projection then so is $1 - p$, and together these two projections decompose space as a direct sum of the image of p with the image of $1 - p$. Thus often a projection gives rise to a decomposition of space into pieces which are hopefully simpler.

The equation $p^2 = p$ makes sense for an endomorphism in an arbitrary category, so projections make sense in an arbitrary category. And if p is a projection then so is $1 - p$ in an arbitrary *additive* category. But in a general (additive or not) category, “the image of p ” (or of $1 - p$) may or may not make sense.

The *Karoubi envelope*¹ of a category \mathcal{C} is a way of adding objects and morphisms to \mathcal{C} so that every projection has an image and so that if $p : \mathcal{O} \rightarrow \mathcal{O}$ is a projection and \mathcal{C} is additive, then (with the proper interpretation) $\mathcal{O} \cong \text{im } p \oplus \text{im}(1 - p)$. Thus sometimes complicated objects can be simplified in the Karoubi envelope of \mathcal{C} , while in \mathcal{C} they may be indecomposable.

Let us turn to the formal definitions.

Definition 3.1. Let \mathcal{C} be a category. An endomorphism $p : \mathcal{O} \rightarrow \mathcal{O}$ of some object \mathcal{O} in \mathcal{C} is called a *projection* if $p \circ p = p$. The Karoubi envelope $\text{Kar}(\mathcal{C})$ of \mathcal{C} is the category whose objects are ordered pairs (\mathcal{O}, p) where \mathcal{O} is an object in \mathcal{C} and $p : \mathcal{O} \rightarrow \mathcal{O}$ is a projection. If (\mathcal{O}_1, p_1) and (\mathcal{O}_2, p_2) are two such pairs, the set of morphisms in $\text{Kar}(\mathcal{C})$ from (\mathcal{O}_1, p_1) to (\mathcal{O}_2, p_2) is the collection of all $f : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ in \mathcal{C} for which $f = f \circ p_1 = p_2 \circ f$. An object (\mathcal{O}, p) in $\text{Kar}(\mathcal{C})$ may also be denoted by $\text{im } p$.²

The composition of morphisms in $\text{Kar}(\mathcal{C})$ is defined in the obvious way (by composing the corresponding f 's). The identity automorphism of an object (\mathcal{O}, p) in $\text{Kar}(\mathcal{C})$ is p itself. It is routine to verify that $\text{Kar}(\mathcal{C})$ is indeed a category. There is an obvious embedding functor $\mathcal{I} : \mathcal{C} \mapsto (\mathcal{O}, I)$ of \mathcal{C} into $\text{Kar}(\mathcal{C})$ and quite clearly, $\text{mor}_{\text{Kar}(\mathcal{C})}(\mathcal{I}\mathcal{O}_1, \mathcal{I}\mathcal{O}_2) = \text{mor}_{\mathcal{C}}(\mathcal{O}_1, \mathcal{O}_2)$ for any pair of objects $\mathcal{O}_{1,2}$ in \mathcal{C} . Thus we will simply identify objects in \mathcal{C} with their image via \mathcal{I} in $\text{Kar}(\mathcal{C})$.

Below we will assume that \mathcal{C} is an additive category and that direct sums of objects make sense in \mathcal{C} . As in [2], there is no loss of generality in making these assumptions as formal sums of morphisms and formal direct sums of objects may always be introduced.

Proposition 3.2. *Let $p : \mathcal{O} \rightarrow \mathcal{O}$ be an endomorphism in \mathcal{C} .*

- (1) *If p is a projection then so is $1 - p$.*
- (2) *In this case, $\mathcal{O} \cong \text{im } p \oplus \text{im}(1 - p)$ in $\text{Kar}(\mathcal{C})$.*

Proof. (1) $(1 - p)^2 = 1 - 2p + p^2 = 1 - 2p + p = 1 - p$ (sorry for the damage to the rainforest).

- (2) The isomorphism $\mathcal{O} \rightarrow \text{im } p \oplus \text{im}(1 - p)$ is given by the 1×2 matrix $(p \quad 1 - p)$. Its inverse

is the 2×1 matrix $\begin{pmatrix} p \\ 1 - p \end{pmatrix}$. □

In this paper we are mainly interested in complexes whose “chain spaces” are objects in some category \mathcal{C} as above. The previous proposition tells us that there may be some gain by switching to working over $\text{Kar}(\mathcal{C})$ as we may have new decompositions of old objects. The proposition below tells us that there is no loss of information in doing so.

Proposition 3.3. *Let Ω_1 and Ω_2 be complexes in $\text{Kom}(\mathcal{C})$. If Ω_1 and Ω_2 are homotopy equivalent as complexes in $\text{Kom}(\text{Kar}(\mathcal{C}))$, they were already homotopy equivalent as complexes in $\text{Kom}(\mathcal{C})$.*

¹The Karoubi envelope construction [10] was first described in [3], a few years before Karoubi. It has previously been used in motivic cohomology [6] and in diagrammatic representation theory, e.g. [4].

²If you're worried about just introducing $\text{im } p$ as notation, when you already know a category-theoretic definition of ‘image’, e.g. [9], don't be; this *is* actually an image.

Proof. A homotopy equivalence Υ in $\text{Kom}(\text{Kar}(\mathcal{C}))$ between Ω_1 and Ω_2 is a certain slew of morphisms between objects appearing in Ω_1 and objects appearing in Ω_2 (a chain morphism going one way, another going the other way and a couple of homotopies). All those morphisms are between objects in \mathcal{C} and as noted above, $\text{Kar}(\mathcal{C})$ introduces no new morphisms between objects in \mathcal{C} . So Υ is really in \mathcal{C} . \square

4. RED AND GREEN IN LEE'S THEORY

Viewed from our perspective, the key to Lee's theorem is the presence in Cob_1 of two complementary projections, the red projection r and the green projection g , that can be composed both vertically (in the "category" direction of Cob_1) and horizontally (in the "planar algebra" direction; see [2, §8.2]). Let us start with some elementary school algebra that contains all the calculations we will need regarding r and g .

Lemma 4.1. *Let b (for `\bullet`) be a variable satisfying $b^2 = 1$ (find it in (2.2)!), let r (for red) be $(1 + b)/2$ and let g (for green) be $(1 - b)/2$. Then r and g are the eigenprojections associated to the involution b . In particular,*

- (1) r and g are projections: $r^2 = r$ and $g^2 = g$.
- (2) r and g are complementary: $r + g = 1$.
- (3) r and g are disjoint: $rg = 0$.
- (4) r and g are eigenprojections of b : $br = r$ and $bg = -g$. \square

In Cob_1 , one may place a b (i.e., a bullet) anywhere on any cobordism and thus, treating linear combinations in the obvious manner, one may place an r or a g anywhere on any cobordism. In particular, we may place them on any "vertical curtain", i.e., on any connected component of the identity morphism from a smoothing to itself. The resulting "identities labeled r and/or g " are projections by the above lemma and hence they represent objects in the Karoubi envelope $\text{Kar}(\text{Cob}_1)$ of Cob_1 . Thus in $\text{Kar}(\text{Cob}_1)$ we have, for example, the isomorphism

$$\left| \begin{array}{c} \cong \\ \hline \end{array} \right| \oplus \left| \begin{array}{c} \hline \end{array} \right|$$

r g

between a single arc smoothing and a direct sum of two single arc smoothings, one paired with the r projection and one with g . Likewise we also have the isomorphism

$$(4.1) \quad \left. \begin{array}{c} \text{) } \left(\right. \\ \hline \end{array} \right) \cong \left. \begin{array}{c} \text{) } \left(\right. \\ \hline \end{array} \right) \oplus \left. \begin{array}{c} \text{) } \left(\right. \\ \hline \end{array} \right) \oplus \left. \begin{array}{c} \text{) } \left(\right. \\ \hline \end{array} \right) \oplus \left. \begin{array}{c} \text{) } \left(\right. \\ \hline \end{array} \right)$$

r r r g g r g g

We can now observe that a dramatic simplification occurs at the very first step of calculating a knot invariant; the complex associated to a single crossing now has an up-to-homotopy representative with a vanishing differential. Indeed, $[[\times]]_1$ is the two-step complex $\succ \xrightarrow{\times} \prec$ in which \times denotes the saddle morphism. As in (4.1), each of the two objects \succ and \prec appearing in this complex becomes a direct sum of four objects in $\text{Kar}(\text{Cob}_1)$. The differential \times becomes a 4×4 matrix M all of whose entries are saddles, and each such saddle carries r and g insertions to match the colourings of its domain and target smoothings. But r and g are disjoint (see Lemma 4.1) and the saddle cobordism is connected, so a saddle bearing insertions of more than one colour vanishes

and hence only two of the 16 entries of M survive. Thus in $\text{Kom}(\text{Kar}(\mathcal{Cob}_1))$,

$$[\times]_1 \cong \begin{bmatrix} r \rangle \langle_r \\ r \rangle \langle_g \\ g \rangle \langle_r \\ g \rangle \langle_g \end{bmatrix} \xrightarrow{\begin{pmatrix} \rangle \langle_r & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \rangle \langle_g \end{pmatrix}} \begin{bmatrix} \smile_r \\ \smile_r \\ \smile_g \\ \smile_r \\ \smile_g \\ \smile_g \end{bmatrix}.$$

This last complex is a direct sum of four complexes. The first and the last of the four summands, $r \rangle \langle_r \xrightarrow{\rangle \langle_r} \smile_r$ and $g \rangle \langle_g \xrightarrow{\rangle \langle_g} \smile_g$, are contractible as $\rangle \langle_r$ and $\rangle \langle_g$ are invertible (with inverses $\frac{1}{2}\smile_r$ and $-\frac{1}{2}\smile_g$, respectively)³. Thus up to homotopy,

$$[\times]_1 \simeq \begin{bmatrix} r \rangle \langle_g \\ g \rangle \langle_r \end{bmatrix} \xrightarrow{0} \begin{bmatrix} \smile_r \\ \smile_g \\ \smile_r \end{bmatrix},$$

and as promised, we found a representative for $[\times]_1$ with a vanishing differential.

Proof of Theorem 1.2. Theorem 1.2 is now simply a matter of assembling the pieces. The discussion above shows that it holds for tangles consisting of a single crossing.

If we build a tangle T by combining crossings X_1 through X_n using a planar operation D , then $[T]_1$ is the tensor product $\bigotimes_D [X_i]_1$.

Since the complexes $[X_i]_1$ have an object for each of the four alternately coloured smoothings of X_i , and no non-zero differentials, the complex $[T]_1$ also has no differentials, but at first sight too many objects. While every alternately coloured smoothing of T appears as an object (because an alternately colouring smoothing can be divided into alternating coloured smoothings of the constituent crossings), we also have smoothings which are alternately coloured, but have different colours appearing on arcs which are connected. Now remember that Lemma 4.1 holds for horizontal compositions as well as vertical ones. A strand with both colours inserted is the same as a strand with the zero projection inserted, and so is equivalent in the category to the zero object; the extra objects all disappear. \square

5. ALTERNATELY COLOURED SMOOTHINGS AND ORIENTING COMPONENTS

In order to return a little closer to Lee's language in [5], we will now show that the alternately coloured smoothings of §4 are in a one-to-one correspondence with orientations of the original tangle.

Proof of Proposition 1.3. To begin, consider a tangle diagram T , and give its regions a red-green checkerboard colouring. (The region 'outside the tangle disk' doesn't receive a colour; further, let's agree that the outer region of a knot, or the 'marked boundary region' of a tangle, is green.)

We can canonically associate to an orientation of T the 'oriented smoothing', in which each arc is consistently oriented. We now need to produce colours for the resulting arcs. As the arcs remain oriented, we can simply take the colour appearing in the region on the right. That this is consistent follows from the observation that as you pass through a crossing, switching from one strand to the other, the checkerboard colours appearing to your left and right remain the same. Further, the colouring we've produced alternates near each confluence, because as two incoming strands enter a crossing they have opposite colours to their right.

Conversely, from an alternately coloured smoothing of a tangle, we need to define an orientation of the tangle. Each arc of the knot has been coloured either red or green, and it has a red and a

³This follows from the neck cutting relation (the last of (2.2)), and the final part of Lemma 4.1.

green region on either side. We orient the arc so that its own colour appears on its right. Moving from one arc of the tangle to another through a crossing, the colour of the arc changes, because the smoothing is alternately coloured, and at the same time the checkerboard colourings on either side switch. This ensures that two opposite arcs at a crossing receive consistent orientations.

These two constructions are clearly inverses, so we have the desired bijection. The four orientations of the Whitehead link and the corresponding alternately coloured smoothings are shown in Figure 1. □




FIGURE 1. Generators of the Khovanov-Lee homology of the Whitehead link. (In greyscale, replace ‘red’ and ‘green’ with ‘dark’ and ‘light’.) All four generators are in homological height zero, because the two components have linking number 0 (see Proposition 5.1).

Without any difficulty, we can also describe the homological height of each generator. For this we need to compare a chosen orientation of our tangle with the fixed ‘original’ orientation (recall that the orientation of a tangle is required in the definition of the Khovanov homology to fix an overall homological height shift). Split the tangle into two parts; T_+ containing those components where the chosen and fixed orientations agree, and T_- containing the components where the orientations disagree.

Proposition 5.1. *The homological height of the corresponding generator is $\text{lk}(T_+, T_-)$, where T_+ and T_- are considered to carry the original orientations.*

Proof. This holds for the two component tangle consisting of a single crossing. If both strands carry the original orientation, or both carry the opposite orientation, then one of T_+ and T_- is empty, so the linking number is zero. The oriented smoothing of this crossing is also the oriented smoothing with respect to the original orientations, and so sits in homological height zero. If exactly one of the strands has been reversed, $\text{lk}(T_+, T_-) = \pm 1$, agreeing with the sign of the crossing (in the original orientation). Happily, the oriented smoothing of the crossing is actually the unoriented smoothing with respect to the original orientations, and so sits in homological height ± 1 , again depending on the sign of the crossing.

To extend the result to arbitrary tangles, it suffices to note that every generator of $[T]_1$ is a planar composition of generators for crossings, and that both $\text{lk}(T_+, T_-)$ and homological height are additive under planar composition. □

REFERENCES

- [1] Dror Bar-Natan, *Fast Khovanov Homology Computations*, arXiv:math.GT/0606318.
- [2] Dror Bar-Natan, *Khovanov's homology for tangles and cobordisms*, *Geom. Topol.* **9** (2005), 1443–1499 (electronic), arXiv:math.GT/0410495.
- [3] Peter Freyd, *Abelian categories. An introduction to the theory of functors*, Harper's Series in Modern Mathematics, Harper & Row Publishers, New York, 1964.
- [4] Greg Kuperberg, *Spiders for rank 2 Lie algebras*, *Comm. Math. Phys.* **180** (1996), no. 1, 109–151, arXiv:q-alg/9712003.
- [5] Eun Soo Lee, *An endomorphism of the Khovanov invariant*, *Adv. Math.* **197** (2005), no. 2, 554–586, arXiv:math.GT/0210213.
- [6] Barry Mazur, *What is ... a motive?*, *Notices Amer. Math. Soc.* **51** (2004), no. 10, 1214–1216.
- [7] Jacob A. Rasmussen, *Khovanov homology and the slice genus*, arXiv:math.GT/0402131.
- [8] Stephan M. Wehrli, *A spanning tree model for Khovanov homology*, arXiv:math.GT/0409328.
- [9] Wikipedia, *Image (category theory) — wikipedia, the free encyclopedia*, 2006, [Online; accessed 20-June-2006].
- [10] ———, *Karoubi envelope — wikipedia, the free encyclopedia*, 2006, [Online; accessed 20-June-2006].

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO ONTARIO M5S 2E4, CANADA
E-mail address: drorbn@math.toronto.edu
URL: <http://www.math.toronto.edu/~drorbn>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY CA 94720, USA
E-mail address: scott@math.berkeley.edu
URL: <http://math.berkeley.edu/~scott>