

**MAT 347**  
**Factorization in polynomial rings**  
**January 21, 2019**

Let us fix an integral domain  $R$  for this worksheet. We want to find out when  $R[X]$  is a UFD. Our strategy is as follows. Let  $F$  be the field of fractions of  $R$ . We know that  $F[X]$  is a Euclidean domain, hence a UFD. Let  $f(X) \in R[X]$ . We can factor  $f(X)$  uniquely as a product of irreducibles in  $F[X]$ , but does this mean we get a unique factorization in  $R[X]$ ?

0. Show that  $R$  is a UFD iff every non-zero non-unit in  $R$  is a product of prime elements.
1. Let  $I \trianglelefteq R$ . Denote by  $(I) \trianglelefteq R[X]$  the ideal of  $R[X]$  generated by the subset  $I$ . Notice that  $(I)$  consists of the polynomials in  $R[X]$  all of whose coefficients are in  $I$ . Prove that

$$R[X]/(I) \cong (R/I)[X].$$

(Hint: remember the isomorphism theorems.)

2. Continue with the notation of Question 1. Prove that  $I \trianglelefteq R$  is a prime ideal iff  $(I) \trianglelefteq R[X]$  is a prime ideal.

*Hint:* Use the characterization of prime ideals in terms of the quotient they generate.

3. Let  $p \in R$ . Prove that  $p$  is prime in  $R$  iff  $p$  is prime in  $R[X]$ .

*Hint:* Use the characterization of prime element in terms of the ideal it generates.

4. Prove that if  $R[X]$  is a UFD, then  $R$  is a UFD.

*Hint:* Remember Question 0.

**For the rest of this worksheet, we will assume that  $R$  is a UFD.**

**Definitions.** Let  $R$  be a UFD. Let  $f(X) \in R[X]$  be non-zero. We define the *content* of  $f(X)$ , denoted  $C_f$ , as the GCD of all the coefficients of  $f(X)$ . We could also interpret  $C_f$  to be the “greatest” divisor of  $f(X)$  among the elements in  $R$ . Notice that content is only defined up to associates. We say that  $f$  is *primitive* if its content is 1. Notice that every non-zero polynomial can be written as the product of its content and a primitive polynomial, and that this decomposition is unique up to multiplication by units.

5. Prove that the product of two primitive polynomials is a primitive polynomial.

*Hint:* Assume that  $f(X), g(X)$  are primitive and that  $d = C_{fg}$ . Let  $p$  be an irreducible factor of  $d$  in  $R$ . Use Question 3.

6. Prove that  $C_{fg} = C_f C_g$  for every non-zero  $f(X), g(X) \in R[X]$ .

7. **Gauss’ Lemma:** Let  $f(X) \in R[X]$ . Prove that if  $f(X)$  is reducible in  $F[X]$ , then it is reducible in  $R[X]$ .

More specifically, assume that

$$f(X) = a(X)b(X)$$

with  $a(X), b(X) \in F[X]$ ,  $\deg a(X) \geq 1$ ,  $\deg b(X) \geq 1$ . Then show that we can find  $\lambda \in F^\times$  such that

$$f(X) = A(X)B(X)$$

with  $A(X) = \lambda a(X) \in R[X]$  and  $B(X) = \lambda^{-1}b(X) \in R[X]$ .

Give an example that shows that it is not possible to conclude that  $a(X), b(X) \in R[X]$ .

*Hint:* first find a non-zero  $d \in R$  such that  $df(X) = a_1(X)b_1(X)$ , where  $a_1(X), b_1(X)$  are in  $R[X]$  and are scalar multiples of  $a(X), b(X)$ . Then try to get rid of  $d$ ...

8. Let  $f(X) \in R[X]$  be primitive. Show that  $f(X)$  is irreducible in  $R[X]$  if and only if it is irreducible in  $F[X]$ .
9. Suppose  $f(X), g(X) \in R[X]$  are primitive. Show that  $f(X), g(X)$  are associates in  $R[X]$  if and only if they are associates in  $F[X]$ .
10. Prove that  $R[X]$  is a UFD. How can you describe the irreducible elements in terms of the irreducibles of  $R$  and  $F[X]$ ?