

**MAT 347**  
**Quotient groups**  
**October 3, 2018**

Let  $G$  be a group and let  $S \subseteq G$ . We want to define an equivalence relation in  $G$  that will identify all the elements in  $S$ , and we want to maintain the group operation. Given  $a, b \in G$ , we say that  $a \sim b$  if there exists  $x \in S$  such that  $b = ax$ . In general, this relation will not be an equivalence relation.

1. Find necessary and sufficient conditions for  $\sim$  to be an equivalence relation.

For the rest of this worksheet, let  $G$  be a group and let  $H \leq G$ . We will consider the equivalence relation defined above with  $S = H$ . Given  $a \in G$ , the *left coset* of  $a$  is the equivalence class of this relation, and we denote it  $aH$ . (Why do we use this notation?) The *quotient set*  $G/H$  is the set of all equivalence classes. The *index* of  $H$  on  $G$ , written  $|G : H|$  is the number of equivalence classes.

2. Prove that  $aH = bH$  iff [there exists  $x \in H$  such that  $b = ax$ ] iff  $a^{-1}b \in H$
3. In general, what is the cardinality of each coset  $aH$ ? What is the relation between  $|G|$ ,  $|H|$ , and  $|G : H|$ ?
4. Consider the group  $G = D_8$  and consider the two subgroups  $H_1 := \langle s \rangle$  and  $H_2 := \langle r \rangle$ . For each of them, write the complete list of cosets, and list which elements are in each coset.

Next, we want to try to use the operation on  $G$  to define an operation on the set  $G/H$ . Given  $aH, bH \in G/H$ , we can try to define their product by

$$(aH) \star (bH) = (ab)H$$

[*Note:* I am using  $\star$  to emphasize that I am defining a new operation. As soon as we make sure this operation works and there is no ambiguity, we will drop the  $\star$ .]

5. In general, the operation  $\star$  is not well-defined. Going back to the example in Question 4, show that with one of those subgroups, the operation is well-defined, but with the other subgroup, the operation is not well-defined.

**Definition:** We say that the subgroup  $H$  is a *normal subgroup* of  $G$  if the operation  $\star$  in  $G/H$  is well-defined. We write  $H \triangleleft G$  (the book writes  $H \trianglelefteq G$ ).

6. Assume  $H \triangleleft G$ . In this case we know the operation in  $G/H$  is well-defined. What other conditions do we need to impose so that  $G/H$  is a group with this operation?

## The big theorem about normal subgroups

**Notation:** Let  $G$  be a group. Given subsets  $A, B$  and elements  $x, y$  we will use the following notation:

$$\begin{aligned}xA &:= \{xa \mid a \in A\}, \\xAy &:= \{xay \mid a \in A\}, \\AB &:= \{ab \mid a \in A, b \in B\}, \text{ etc.}\end{aligned}$$

7. Let  $G$  be a group and let  $H \leq G$ . Explore the relation between the following statements (which ones imply which ones)?
- (a)  $H \triangleleft G$ .
  - (b)  $aH = Ha$  for all  $a \in G$ .  
(Notice that this does not mean that  $a$  commutes with the elements of  $H$ . It only means that the sets  $aH$  and  $Ha$  are the same set.)
  - (c)  $aHa^{-1} = H$  for all  $a \in G$ .
  - (d)  $aHa^{-1} \subseteq H$  for all  $a \in G$ .
  - (e) There exists some group  $L$  and some group homomorphism  $f : G \rightarrow L$  such that  $H = \ker f$ .

## Another look at quotient groups

8. Suppose that  $G$  is a group and  $\sim$  an equivalence relation on  $G$  such that the set  $G/\sim$  of equivalence classes becomes a group in the natural way. In other words,  $\bar{g} \cdot \bar{h} = \overline{gh}$  for all  $g, h \in G$ , where  $\bar{x}$  denotes the equivalence class of any  $x \in G$ . Show that  $\sim$  is one of the equivalence relations considered in Problem 1, where  $S = N$  is a normal subgroup of  $G$ , i.e.  $G/\sim$  is nothing but the quotient group  $G/N$ . (Hint: it may help to notice that the map  $G \rightarrow G/\sim$  sending  $x$  to  $\bar{x}$  is a homomorphism.)