

Tychonoff's Theorem Lecture

Micheal Pawliuk

November 24, 2011

1 Preliminaries

In point-set topology, there are a variety of properties that are studied. When given a property of a topological space there are always 4 basic questions that need answering:

1. Is there any space with this property?
2. Is this property preserved under continuous maps?
3. Is this property preserved under products?
4. Is this property preserved under subspaces?

For example, the T_2 property is very nicely behaved in that it is preserved under continuous maps, products and subspaces. Connectedness is preserved under continuous maps, but not subspaces.

With compactness we have already seen that it is preserved under continuous maps, *closed* subspaces and *finite* products. Tychonoff's Theorem asserts that an arbitrary (i.e. finite or infinite) product of compact spaces is again compact. In the proof that compactness is preserved under finite products, we use that there are only finitely many coordinates. We cannot easily extend this proof to infinitely many coordinates, so we have to do something clever.

Let us first look at some examples of infinite products of compact spaces:

- The simplest example is $\{0\}^\omega$ which is just a single infinite sequence of zeros. Of course this is compact because it is finite.
- The first non-trivial example is $\{0, 1\}^\omega$. Showing that this is compact requires some thought, but can be seen by noting that $\{0, 1\}^\omega$ is homeomorphic to the Cantor set, which is compact. Another strategy is to use König's Lemma (which you can find online).

- Think about how you can show that $[0, 1]^\omega$ is compact. It may help to observe that this space is a metric space.

Tychonoff's Theorem says something about the product of compact spaces *with the product topology*. It does not say anything about spaces with the box topology. For example, check that $\{0, 1\}^\infty$ and $[0, 1]^\infty$ (both with the box topology) are not compact.

2 Tools-Filters

In order to prove Tychonoff's Theorem we will use an equivalent notion of compactness. Since we do not know anything about the extra topological properties possessed by the compact spaces we have (like metrizability), we cannot use Theorem 44 from class.

Definition 1. Recall that a collection \mathcal{F} of subsets of some set X has the **finite intersection property (FIP)** if for every finite subcollection $\{F_1, F_2, \dots, F_n\} \subseteq \mathcal{F}$ we have $\bigcap_{i=1}^n F_i \neq \emptyset$.

Theorem 2 (Proposition 46 in class). A topological space X is compact iff for every collection \mathcal{F} of closed subsets of X with the FIP, we have $\bigcap \mathcal{F} \neq \emptyset$.

This has some hints that it will be helpful when looking at the product topology, because the basic open sets in the product topology are given by *finite intersections* of subbasic open sets and subbasic sets only give information about an individual coordinate. This will be how we exploit compactness of individual coordinates.

One of the main tools we use is that of a maximal collection of sets with the FIP. These maximal collections turn out to satisfy the properties of being maximal filters.

Definition 3. Let \mathcal{C} be a collection of subsets of a topological space X , with the FIP. We say that \mathcal{C} is **maximal** (with respect to the FIP) if whenever \mathcal{A} is a collection of subsets with the FIP such that $\mathcal{C} \subseteq \mathcal{A}$, then $\mathcal{C} = \mathcal{A}$.

Proposition 4. For every collection of closed subsets \mathcal{C} with the FIP, there is a maximal \mathcal{A} such that $\mathcal{C} \subseteq \mathcal{A}$.

To prove this we will need a theorem (which is called a lemma for historical reasons) from set theory:

Lemma 5 (Zorn's Lemma). Let (P, \leq) be a (non-empty) partially ordered set such that every linearly ordered subset has an upper bound. Then P contains a maximal element.

The partial order we will be using is the family \mathcal{T} of all collections of subsets of a space X with the FIP that contain \mathcal{C} , ordered by \subseteq . So then $\mathcal{T} \neq \emptyset$ as $\mathcal{C} \in \mathcal{T}$.

Proof of Proposition 4. Let \mathcal{C} be a collection of subsets of X with the FIP. Let A be a linearly ordered subset of \mathcal{T} , the family of all collections of subsets of a space X with the FIP and that contain \mathcal{C} . We need to show that A has an upper bound.

Claim: $\bigcup_{\mathcal{F} \in A} \mathcal{F}$ is an upper bound for A .

It is clear that $\mathcal{F} \subseteq \bigcup_{\mathcal{F} \in A} \mathcal{F}$ for each $\mathcal{F} \in A$. All that remains to show is that $\bigcup_{\mathcal{F} \in A} \mathcal{F}$ has the FIP. Let $F_1, F_2, \dots, F_n \in \bigcup_{\mathcal{F} \in A} \mathcal{F}$, then there are $\mathcal{F}_1, \dots, \mathcal{F}_n$ such that $F_i \in \mathcal{F}_i$ for $1 \leq i \leq n$. Since A is linearly ordered there is a $1 \leq N \leq n$ such that $\mathcal{F}_i \subseteq \mathcal{F}_N$ for all $1 \leq i \leq n$. So then $F_i \in \mathcal{F}_N$ for all $1 \leq i \leq n$ and because \mathcal{F}_N has the FIP,

$$\bigcap_{i=1}^n F_i \in \mathcal{F}_N \subseteq \bigcup_{\mathcal{F} \in A} \mathcal{F}$$

Thus Zorn's Lemma applies and \mathcal{T} has a maximal element as desired. \square

Such a maximal extension will actually satisfy the stronger properties of being a filter.

Definition 6. A collection of subsets \mathcal{F} of a space X is called a **filter** if:

1. Finite intersections are contained in \mathcal{F} . That is: $(\forall n \in \mathbb{Z}^+) (\forall F_1, \dots, F_n \in \mathcal{F}) [\bigcap_{i=1}^n F_i \in \mathcal{F}]$
2. $\emptyset \notin \mathcal{F}$ and $X \in \mathcal{F}$
3. \mathcal{F} is closed upwards. That is if $B \subseteq A \subseteq X$ with $B \in \mathcal{F}$ then $A \in \mathcal{F}$.

Remark 7. Any filter satisfies the FIP by property (1) and (2).

Definition 8. Let \mathcal{F} be a collection of subsets of a topological space X , that is a filter. We say that \mathcal{F} is a **maximal filter** if it is maximal with respect to FIP.

Traditionally maximal filters are meant as filters maximal with respect to filters. You may wish to do this exercise after Theorem 11. In that case, make sure that in the proof of Theorem 11 we only ever use maximality with respect to the FIP and not maximality with respect to filters.

Exercise 9. Let \mathcal{F} be a filter on a topological space X . TFAE:

- \mathcal{F} is maximal with respect to the FIP.
- \mathcal{F} is maximal with respect to filters.

Example 10. For $X = \mathbb{R}$, consider the following:

- $\mathcal{F}_1 := \{(a, b) \subseteq \mathbb{R} : 100 \in (a, b)\};$
- $\mathcal{F}_2 := \{A \subseteq \mathbb{R} : \exists(a, b) \ni 100, (a, b) \subseteq A\};$ and
- $\mathcal{F}_3 := \{A \subseteq \mathbb{R} : 100 \in A\}.$

Here $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \mathcal{F}_3$ and all have the FIP. However \mathcal{F}_1 is not a filter, \mathcal{F}_2 is a filter and \mathcal{F}_3 is a maximal filter.

Theorem 11. Suppose that \mathcal{F} is a collection of subsets of some set X that is maximal with respect to the FIP. Then \mathcal{F} is a (maximal) filter.

Proof. (1) holds: We need to show that finite intersections are in \mathcal{F} . Let $F_1, \dots, F_n \in \mathcal{F}$.

Then $\bigcap_{i=1}^n F_i \neq \emptyset$ by the FIP. Now it is clear that $\{\bigcap_{i=1}^n F_i\} \cup \mathcal{F}$ has the FIP. So by maximality

of \mathcal{F} we have $\bigcap_{i=1}^n F_i \in \mathcal{F} = \{\bigcap_{i=1}^n F_i\} \cup \mathcal{F}$.

(2) holds: Clearly $\emptyset \notin \mathcal{F}$ as \mathcal{F} has the FIP. Thus $X \cap F = F \neq \emptyset$ for each $F \in \mathcal{F}$. So $\mathcal{F} \cup \{X\}$ has the FIP. By maximality of \mathcal{F} , $X \in \mathcal{F} = \mathcal{F} \cup \{X\}$.

(3) holds: This is analogous to how we showed $X \in \mathcal{F}$. \square

The most important part of a filter is property (1). The only other fact about maximal filters we will need is the following way to identify whether a set belongs to the maximal filter.

Theorem 12. Suppose that \mathcal{F} is a maximal filter on a set X and $A \subseteq X$. TFAE:

1. $A \in \mathcal{F}$

2. $\forall F \in \mathcal{F}, A \cap F \neq \emptyset$

Proof. Suppose that $A \in \mathcal{F}$. Then since \mathcal{F} has the FIP, this is clear. Conversely, suppose that $\forall F \in \mathcal{F}, A \cap F \neq \emptyset$. We show that $\{A\} \cup \mathcal{F}$ has the FIP, in which case the result follows by maximality of \mathcal{F} . Let $\mathcal{F}_1, \dots, \mathcal{F}_n \in \mathcal{F}$. Since \mathcal{F} is a filter we have that $\emptyset \neq \bigcap_{i=1}^n F_i \in \mathcal{F}$.

So then $(\bigcap_{i=1}^n F_i) \cap A \neq \emptyset$, by assumption. \square

The thing to observe is that maximal filters have a strong form of the FIP (property (1) in the definition of a filter) and we have an alternate way of detecting when a set is in a maximal filter. It is also worth noting that Munkres proves Tychonoff's theorem without

making mention of filters. Instead, the notion of a maximal collection with the FIP is used. This results in a proof with less notation, although on the other hand, filters are a useful notion in mathematics. Filters turn out to be a useful generalization of sequences.

3 Tychonoff's Theorem

We are now in a position to prove Tychonoff's Theorem:

Theorem 13 (Tychonoff's Theorem). *Let $\{X_\lambda : \lambda \in I\}$ be a collection of compact spaces. Then $\prod_{\lambda \in I} X_\lambda$ is compact.*

Proof. Let \mathcal{C} be a collection of closed subsets of $\prod X_\lambda$ with the FIP. We wish to show that $\bigcap \mathcal{C} \neq \emptyset$. Extend \mathcal{C} to a maximal filter \mathcal{F} (by Proposition 4 and Theorem 9). Since $\mathcal{C} \subseteq \mathcal{F}$ we get $\bigcap \mathcal{F} \subseteq \bigcap \mathcal{C}$. In fact we get $\bigcap_{F \in \mathcal{F}} \overline{F} \subseteq \bigcap \mathcal{C}$, so it is enough to show that $\emptyset \neq \bigcap_{F \in \mathcal{F}} \overline{F}$.

We now transfer the filter down to the coordinates using the usual projection maps π_λ . Let $\mathcal{F}_\lambda := \{\pi_\lambda(F) : F \in \mathcal{F}\}$. This collection has the FIP because $\emptyset \neq \pi_\lambda(F_1 \cap F_2) \subseteq \pi_\lambda(F_1) \cap \pi_\lambda(F_2)$. Thus the collection of closed sets $\{\overline{\pi_\lambda(F)} : F \in \mathcal{F}\}$ has the FIP. Now since X_λ is compact, we can choose $x_\lambda \in \bigcap_{F \in \mathcal{F}} \overline{\pi_\lambda(F)}$ for each $\lambda \in I$. Thus $\underline{x} = (x_\lambda) \in \prod_{\lambda \in I} X_\lambda$.

Claim: $\underline{x} \in \bigcap_{F \in \mathcal{F}} \overline{F}$.

Here is where we do something clever. We show that every basic open set $B \ni \underline{x}$ has the property that $B \in \mathcal{F}$. This will tell us that for all basic open neighbourhoods B of \underline{x} that $\forall F \in \mathcal{F}, B \cap F \neq \emptyset$, equivalently $\forall F \in \mathcal{F}, \underline{x} \in \overline{F}$ (by Theorem 14 in class). Thus we have the claim and the theorem.

To show that every basic open set containing \underline{x} is in \mathcal{F} it is enough to show that every subbasic open set containing \underline{x} is in \mathcal{F} , because \mathcal{F} is closed under finite intersections (property (1) of filter), and every basic open set is a finite intersection of subbasic open sets.

Subclaim: All subbasic open sets containing \underline{x} are in \mathcal{F} .

Let $S = \pi_\lambda^{-1}(U_\lambda)$ be a subbasic open set containing \underline{x} , where U_λ is open in X_λ . Then $x_\lambda \in U_\lambda$. By our choice of x_λ , we have $x_\lambda \in \bigcap_{F \in \mathcal{F}} \overline{\pi_\lambda(F)}$. Thus $x_\lambda \in U_\lambda \cap \overline{\pi_\lambda(F)}$ for each $F \in \mathcal{F}$. So $\emptyset \neq U_\lambda \cap \pi_\lambda(F)$ and so $\emptyset \neq \pi_\lambda^{-1}(U_\lambda) \cap F = S \cap F$. And by Theorem 12, $S \in \mathcal{F}$. \square

4 Axiom of Choice

Our use of Zorn's Lemma deserves some comment. It might seem to you that this proof doesn't need to use some theorem that we pulled from set theory. You might ask if we can

accomplish this proof in a “purely topological way”. As it turns out, set theorists have shown that you do need to use Zorn’s Lemma, or something equivalent:

Theorem 14. *TFAE:*

1. *Zorn’s Lemma*
2. *The axiom of choice*
3. *If $\{A_\lambda : \lambda \in I\}$ is a collection of non-empty sets, then $\emptyset \neq \prod_{\lambda \in I} A_\lambda$.*
4. *Every set can be well-ordered.*
5. *Tychonoff’s Theorem*
6. *(The Compactness Theorem in Logic) Every finitely consistent model is consistent.*

It seems like (3) is obviously true, right? However, (1) seems pretty non-intuitive. Due to some highly non-intuitive consequences of Zorn’s Lemma (like the Banach-Tarski Paradox) some mathematicians are skeptical about the use of the axiom of choice and its equivalences. As a result whenever there is a proof using the axiom of choice it is interesting to know whether or not you actually need to use the axiom of choice.

For example, consider Proposition 4. Our proof used Zorn’s Lemma (to help streamline the proof), but it turns out that it is provable from a weaker principle than Zorn’s Lemma called the boolean prime ideal theorem (BPIT).

In our proof of Tychonoff’s theorem we used the axiom of choice somewhere. Can you find it? It turns out that we can remove this use of choice if we assume something about our spaces:

Exercise: Prove the following version of Tychonoff’s Theorem without using the axiom of choice (or any of its equivalences), but assuming BPIT (thus also Proposition 4).

Theorem 15. *Let $\{X_\lambda : \lambda \in I\}$ be a collection of compact T_2 spaces. Then $\prod_{\lambda \in I} X_\lambda$ is compact and T_2 .*