

A CHARACTER FORMULA FOR COMPACT ELEMENTS (THE RANK ONE CASE)

JONATHAN KORMAN

ABSTRACT. In their 1997 paper, [SS2], Schneider and Stuhler gave a formula relating the value of an admissible character of a p -adic group at an elliptic element to the fixed point set of this element on the Bruhat-Tits building. Here we give a similar formula which works for compact elements. Elliptic elements have finitely many fixed facets in the building but compact elements can have infinitely many. In order to deal with the compact case we truncate the building so that we only look at a bounded piece of it. We show that for compact elements the (finite) information contained in the truncated building is enough to recover all of the information about the character. This works since the fixed point set of a compact (non elliptic) element is periodic. The techniques used here are more geometric in nature than the algebraic ones used by Schneider and Stuhler. We recover part of their result as a special case.

CONTENTS

| | |
|---|----|
| 0. Introduction | 1 |
| 1. Notation | 5 |
| 2. Preliminaries | 6 |
| 3. Overview of the Schneider-Stuhler result | 14 |
| 4. Truncated buildings and truncation operators | 17 |
| 5. Modified truncation operators | 21 |
| 6. A character formula for functions | 24 |
| 7. Nice decompositions | 26 |
| 8. Recovering Schneider-Stuhler's result for semisimple rank 1 groups | 36 |
| 9. A character formula for compact elements | 39 |
| 10. Concluding remarks | 50 |
| References | 50 |

Date: Jan 11, 2004.

1991 Mathematics Subject Classification. AMS subject classification: primary: 22E50; secondary: 22E35 .

0. INTRODUCTION

In this paper we use the geometry of the (semisimple) Bruhat-Tits building to obtain a character formula for finitely generated admissible representations of a connected reductive p -adic group.

0.1. Given an admissible representation (π, V) of such a group, one would like to understand its character Θ_π , which is a complex valued function on the group. Ideally one would like a formula that expresses the value of the character at any given element of the group.

Associated with a reductive p -adic group G there is a geometric space X , called the (semisimple) Bruhat-Tits building, or affine building, of G . This building is a polysimplicial complex endowed with a G -action.

As a geometric space, the building only encodes information about its associated group, and not about a specific representation V of that group. In order for it to contain information about such a representation and its character, we have to consider a certain sheaf on X associated to V . The action of the group G on the building extends to this sheaf.

The formula for the value of characters (of finitely generated admissible representations) on certain elements of G is in terms of the action of these elements on X and on this sheaf.

In 1997 Schneider and Stuhler [SS2], using algebraic techniques, gave such a formula for elliptic elements – elements which have a finite number of fixed facets in the building. Using different techniques, yet relying on some of their basic results, we give a formula which works for compact elements – which can have an infinite number of fixed facets. In the elliptic case, this gives a new proof of the Schneider-Stuhler result (for groups of semisimple rank 1).

The main idea is the use of certain truncation operators which pick out finite subsets of the fixed-point set. These subsets contain enough information from which to recover the character.

0.2. Let k be a p -adic field of characteristic zero. Let \mathbf{G} be a connected reductive algebraic group defined over k and $G = \mathbf{G}(k)$ its group of k -rational points. Denote by $X = \mathcal{B}(\mathcal{D}\mathbf{G}, k)$ the semisimple Bruhat-Tits building; that is the Bruhat-Tits building associated to the derived group, $\mathcal{D}\mathbf{G}$, of G . Let (π, V) be a finitely generated admissible representation of G . Let $U_F^{(e)}$ be the open-compact subgroups associated by Schneider-Stuhler [SS2] to the facets F of X . These subgroups satisfy the relations $U_{F'}^{(e)} \subset U_F^{(e)}$ whenever $F' \subset \overline{F}$. Consequently $V^{U_F^{(e)}}$ are finite dimensional subspaces of V associated to the facets F . The relations between the groups translate into relations between

these vector spaces: $V^{U_F^{(e)}} \subset V^{U_{F'}^{(e)}}$.

In [SS2], Schneider and Stuhler prove:

Theorem. Let G be a connected reductive group, $\gamma \in G$ regular semi-simple elliptic, and (π, V) a finitely generated admissible representation of G . Then there exists an integer $e_0 = e_0(V)$ which depends on V , such that for all $e \geq e_0$ the character Θ_π can be expressed as:

$$\Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma)_q} (-1)^q \text{trace}(\gamma, V^{U_F^{(e)}})$$

Here d is the dimension of X , X^γ (a polysimplicial complex, but not necessarily a subcomplex of X) is the fixed point set of γ , $(X^\gamma)_q$ is the set of q -facets of X^γ and $F(\gamma) := F \cap X^\gamma$ (a polysimplex).

Remark. The summation in the formula above should be understood to mean the sum over all of the γ -stable facets in X . When the action of G on X preserves the types¹ of the vertices in X , a facet is γ -stable if and only if it is γ -fixed. Thus if this is the case then the summation is over all the γ -fixed facets and the above formula simplifies to:

$$\Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F \in X_q^\gamma} (-1)^q \text{trace}(\gamma, V^{U_F^{(e)}})$$

0.3. Let $C_q := C_c^{or}(X_{(q)}; \gamma_e(V))$ be the vector space of oriented q -chains with compact support (see [SS2, II]). This is a *smooth* representation of G . Write T_g for the action of $g \in G$ on the C_q 's. Consider the complex of (smooth) G -modules:

$$(*) \quad 0 \longrightarrow C_d \xrightarrow{\partial} \cdots \xrightarrow{\partial} C_0 \xrightarrow{\epsilon} V \longrightarrow 0$$

For V finitely generated admissible there exists an integer $e_0 = e_0(V)$ depending on V such that for all $e \geq e_0$ the complex $(*)$ is exact ([SS1] and [SS2, Theorem II.3.1]).

We would like to apply the *Hopf trace formula* to the operators T_g acting on $(*)$. The T_g 's commute with the boundary operators (∂ and ϵ), but they are not of finite rank. In an attempt to address this issue we introduce truncation operators Q_q^r , ($r \geq 0$, a real number) on $X_{(q)}$ which can be viewed as acting on C_q . Roughly, one can think of these truncation operators as intersecting the building with a ball of radius r about some fixed point $o \in X^\gamma$. An advantage of the Q_q^r 's is that they are of finite rank, but unfortunately they do not commute with ∂ . To fix this, modify the truncation operators as follows.

¹This happens for example if G is semisimple simply connected.

Fix a vector space direct sum decomposition of C_q (see section 5 for details):

$$(\oplus) \quad C_q = B_q \oplus H'_q \oplus B'_{q-1}$$

Using such a decomposition it is possible to define modified truncation operators, \overline{Q}_q^r on C_q . These modified operators have all of the desired properties: they commute with ∂ (and ϵ), they have finite rank and they tend to the identity operator on C_q , Id_{C_q} , as r tends to ∞ . Using these modified operators we obtain:

Theorem. Let G be connected reductive, V a finitely generated admissible representation of G , $f \in C_c^\infty(G)$, a locally constant, compactly supported function and $e \geq e_0$. There exists a radius of truncation $r_0 = r_0(f)$, which depends on f , such that for all $r \geq r_0$ the trace of the operator $\pi(f)$ can be expressed as:

$$\text{trace } \pi(f) = \sum_{q=0}^d (-1)^q \text{trace}(T_f \overline{Q}_q^r, C_q).$$

We would like to give a more geometric interpretation of this formula in terms of the original truncation operators. With this purpose in mind, assume that the direct sum decomposition is T_γ -equivariant ($\gamma \in G^{cpt}$) and ‘nice’. Then $\text{trace}(T_\gamma \overline{Q}_q^r, C_q) = \text{trace}(T_\gamma Q_q^r, C_q)$ and the above formula gives the main result (see Theorem 9.10 for a more precise statement):

$$\Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma \cap X^r)_q} (-1)^q \text{trace}(\gamma, V^{U_F^{(e)}}).$$

Here X^r is a finite subcomplex of X called the truncated building.

0.4. For the following technical reasons we can prove this formula in full generality only for groups of semisimple rank 1. The direct sum decomposition (\oplus) is controlled by the following truncated complex:

$$(\overline{*}) \quad 0 \longrightarrow C_c^{or}(X_d^r; V) \xrightarrow{\partial'} \dots \xrightarrow{\partial'} C_c^{or}(X_0^r; V) \xrightarrow{\epsilon'} V$$

The exactness of this complex guarantees the existence of a nice direct sum decomposition. Exactness of $(\overline{*})$ may or may not depend on a parameter e_r . To prove the strong version of the main result it is necessary for it not to depend on e_r . We have the following:

- For $V = \mathbb{C}$ the trivial representation, the exactness of $(\overline{*})$ follows from the contractibility of the truncated building X^r , and is independent of e_r .
- For G of semisimple rank 1, we prove exactness (independently of e_r) using the fact that X has non-positive curvature, which implies that the distance function on the building is strictly convex.

- For a general connected reductive group G , we prove exactness of $(\bar{*})$ but the technique used is not independent of e_r . Thus in this case we can only obtain a weaker version of the main result.

0.5. The geometric techniques used here might suggest how one could proceed with the case of non-compact elements. Such elements do not have fixed points on X but they do have fixed points on the spherical building at infinity, X^∞ , which is another building associated with G . Finding a character formula for non-compact elements is still an open problem.

Acknowledgments. I would like to express deep gratitude to my thesis advisor Tom Hales for guiding, teaching and transforming me.

Special thanks to Ju-Lee Kim for her advice and kindness, to Jeff Adler for his generosity and to my friends: Julia Gordon, Elliot Lawes and Joel Pitkin for sharing with me this long path.

1. NOTATION

- k a p -adic field of characteristic zero.
- k^\times the set of non zero elements of k .
- O the ring of integers of k .
- O^\times the set of units in k .
- ϖ a fixed generator for the maximal ideal in O .
- $\bar{k} = O/\varpi O$ the residue field of k .
- \bar{k}^\times the set of non zero elements of \bar{k} .
- $\omega : k^\times \rightarrow \mathbb{Z}$ the discrete valuation normalized by $\omega(\varpi) = 1$.
- $G = \mathbf{G}(k)$ a connected reductive group.
- G^{reg} the set of regular semisimple elements in G .
- G^{ell} the set of regular semisimple elliptic elements in G .
- G^{cpt} the set of regular semisimple compact elements in G .
- γ a compact (sometimes also elliptic) element in G .
- X the semisimple Bruhat–Tits building of G .
- A a basic apartment in X .
- F a facet of X .
- \bar{F} the closure of the facet F in X (a polysimplex).
- X^r a truncated building with truncation parameter r .
- X_q the q -dimensional facets of X .
- $X_{(q)}$ the oriented q -dimensional facets of X .
- X^g the fixed point set of $g \in G$.
- (π, V) a finitely generated admissible representation.
- Θ_π the character of (π, V) .
- $\mathcal{O}_\gamma(f)$ the orbital integral of a function f with respect to an element γ .

2. PRELIMINARIES

Let k denote a p -adic field of characteristic zero, that is, a finite extension of \mathbb{Q}_p for some prime p . We will denote by O the ring of integers of k and pick a generator ϖ for the maximal ideal in O . The residue field of k will be denoted by \bar{k} . Let \mathbf{G} be a connected reductive group defined over k and denote by G the group $\mathbf{G}(k)$ of k -rational points of \mathbf{G} , equipped with the natural locally compact topology induced from that on k .

2.1. The building. A building is a polysimplicial complex which can be expressed as the union of subcomplexes called apartments satisfying certain axioms (see [Bro, p.78]). There are two types of buildings: affine buildings and spherical buildings. The apartments of affine buildings are Euclidean spaces and those of spherical buildings are spheres. To a connected reductive group G one can associate at least three kinds buildings:

- The semisimple Bruhat-Tits building of G . This is a building of affine type.
- The Bruhat-Tits building of G . This building is also of affine type.
- The spherical building (at infinity) of G . This is a building of spherical type.

Remark 2.1. If G is semisimple then the semisimple Bruhat-Tits building and the Bruhat-Tits building are the same. For a reductive, non-semisimple group G , the Bruhat-Tits building is a product of the semisimple Bruhat-Tits building and an affine building associated to the center of G .

In this paper we will only use the semisimple Bruhat-Tits building. We follow the review in [SS2, I.1] of the construction of such a building.

2.2. Review of the Semisimple Bruhat-Tits building. We will use the following notation.

$G = \mathbf{G}(k)$ be a connected reductive group.

S a maximal k -split torus in G .

$X^*(S) := \text{Hom}_k(S, K^\times)$ the lattice of rational characters of S .

$X_*(S) := \text{Hom}_k(K^\times, S)$ the (dual) lattice of rational co-characters of S .

$C := Z(G)^\circ$ be the connected component of the center of G .

$X_*(C)$ the lattice of rational co-characters of C .

Z the centralizer of S in G .

N the normalizer of S in G .

$W := N/Z$ the Weyl group.

Definition. The underlying affine space of the real vector space

$$A := (X_*(S)/X_*(C)) \otimes \mathbb{R}$$

is called the *basic apartment*.

The Weyl group W acts by conjugation on S which induces a faithful linear action of W on A .

Let $\langle \cdot, \cdot \rangle: X_*(S) \times X^*(S) \rightarrow \mathbb{Z}$ be the natural pairing; its \mathbb{R} -linear extension is also denoted by $\langle \cdot, \cdot \rangle$.

There is a unique homomorphism

$$\nu: Z \rightarrow X_*(S) \otimes \mathbb{R}$$

characterized by

$$\langle \nu(g), \chi|_S \rangle = -\omega(\chi(g))$$

for all $g \in Z$, and all characters χ of Z (here ω is the discrete valuation).

Using this homomorphism $g \in Z$ acts on A by translations

$$gx := x + \text{image of } \nu(g) \text{ in } A \quad x \in A.$$

This action of Z on A can be extended to an action of N on A . The N action is compatible with the action of the Weyl group W . Recall that there exists [Tit, p.31] a system of affine roots Φ_{af} (which are certain affine functions on A), and a mapping $\alpha \mapsto U_\alpha$ from Φ_{af} onto a set of subgroups of G .

Definition. Two points in A are called equivalent if all affine roots have the same sign on these two points; the corresponding equivalence classes are called *facets*. The facets of maximal dimension d are called *chambers* (they are also the connected components of the complements in A of the union of walls; a *wall* is the zero set of an affine root). The 0-dimensional facets are the *vertices*. The closure \overline{F} of a facet F is a *polysimplex* (in the sense of algebraic topology). This gives the basic apartment a polysimplicial structure.

Consider the following equivalence relation on $G \times A$:

$$(g, x) \sim (g', x') \text{ if there is an } n \in N \text{ such that } nx = x' \text{ and } g^{-1}g'n \in U_x.$$

We define $X := G \times A / \sim$. It is easy to see that G acts on X . This action extends the action of N on A and the polysimplicial structure of A extends to X .

Definition. The *semisimple Bruhat-Tits building* of G is the polysimplicial G -complex X , also denoted by $\mathcal{B}(\mathcal{D}\mathbf{G}, k)$.

2.3. Main properties of the building. Let $X = \mathcal{B}(\mathcal{D}\mathbf{G}, k)$ be the semi-simple Bruhat-Tits building of G ; this is equivalent to saying that X is the Bruhat-Tits building of the derived group $\mathcal{D}\mathbf{G}$ of \mathbf{G}/k .

We list the properties of the building which will be used in this paper (see [Bro], [Moy, 1.1] and [SS2, I.1] for more details).

- The building associated to G is made up of *apartments* which are glued together. Each apartment is a Euclidean space equipped with a polysimplicial structure and is isomorphic to A (see [SS2, p.10]).

- The building is a d -dimensional locally finite polysimplicial complex, where $d = \dim(A)$ is the semisimple k -rank of G . The group G acts on X polysimplicially. (If $\mathcal{D}\mathbf{G}$ is simple, as opposed to semisimple, then the building and the action are simplicial, as opposed to polysimplicial)
- Topologically, the building X is a contractible space with a natural G -action.
- There is a natural metric $d(\cdot, \cdot)$ on X with respect to which the action of G is by isometries.
- Any two points (and even any two facets) in X are contained in a common apartment.
- Any two points $x, y \in X$ are connected by a unique geodesic line segment, denoted $\text{geod}(x, y)$.

Here X_q will denote the space of all q -facets of X and $X_{(q)}$ will denote the space of all oriented q -facets. Also write $d = \dim(X)$ for the dimension of X as a locally finite polysimplicial complex.




FIGURE 1. The (semisimple) Bruhat-Tits building of the group $SL_2(\mathbb{Q}_2)$.

Example 2.2. For $G = SL_2(\mathbb{Q}_p)$, the semisimple Bruhat-Tits building X is a *tree* with $p + 1$ edges meeting at every vertex. Since the group SL_2 is semisimple, its semisimple Bruhat-Tits building and Bruhat-Tits building are the same. The apartments of this building (tree) are one dimensional Euclidean spaces (lines). The 0-dimensional facets are the vertices and the 1-dimensional facets are the edges. See Figure 1.

Since GL_2 and SL_2 have the same derived group, their semisimple Bruhat-Tits buildings are the same. For the actions of SL_2 on this building there are two types of vertices; for the action of GL_2 there is only one type².

2.4. The character. We include Fiona Murnaghan's [Mur] explanation of the character of an admissible representation.

A representation (π, V) of G is called *smooth* if

$$Stab_G(v) := \{g \in G \mid \pi(g)v = v\}$$

is open for every $v \in V$.

Let $C_c^\infty(G)$ denote the space of complex valued, locally constant, compactly supported function on G . Given $v \in V$ and $f \in C_c^\infty(G)$, the function $g \mapsto f(g)\pi(g)v$ belongs to $C_c^\infty(G, V)$, compactly supported, locally constant functions with values in V . Therefore, for each $f \in C_c^\infty(G)$, we can define an operator $\pi(f)$ on V as follows:

$$\pi(f)v = \int_G f(g)\pi(g)v dg, \quad v \in V.$$

Here, dg denotes a fixed Haar measure on G .

To define the character of π , we want to take the trace of $\pi(f)$. Since V is infinite dimensional in general this trace is not defined. In order to make sense of the trace of $\pi(f)$ we restrict the class of representations from smooth to admissible.

Recall that a smooth representation (π, V) of G is said to be *admissible* if for every open compact subgroup K of G , the space

$$V^K = \{v \in V \mid \pi(k)v = v \forall k \in K\}$$

is finite dimensional. It can be shown that π is admissible if and only if $\pi(f)$ has finite rank for all f in $C_c^\infty(G)$. Thus for π admissible we can talk about the trace of the operator $\pi(f)$. Write

$$\Theta_\pi(f) = \text{trace } \pi(f), \quad f \in C_c^\infty(G).$$

The distribution (linear functional on $C_c^\infty(G)$) defined by $f \mapsto \Theta_\pi(f)$ is the character of π .

It is a theorem of Harish-Chandra (see [Mur, p.2]) that for π an admissible representation of finite length³, there exists a locally integrable function, also denoted by Θ_π , on G , which is locally constant on the set of regular elements in G , and satisfies

$$\Theta_\pi(f) = \int_G f(g)\Theta_\pi(g)v dg, \quad f \in C_c^\infty(G).$$

²For explanation of *type* (in a special case), see [Bro, p.30].

³Recall that a finitely generated admissible representation has finite length.

That is, the distribution Θ_π is given by integration against a function. The function Θ_π is also called the character of G .

2.5. Three types of elements. Here G^{reg} will denote the set of regular semisimple elements in G . Recall that for an element $\gamma \in G^{reg}$, the connected component of its centralizer, $T := C_G(\gamma)^\circ$, is a maximal torus in G .

We classify elements $\gamma \in G^{reg}$ into three types⁴ according to their set of fixed points on the semisimple Bruhat–Tits building X .

Definition 2.3. Let $\gamma \in G^{reg}$ be a regular semisimple element.

We call γ *compact* if its fixed point set, X^γ , is non-empty. Write G^{cpt} for the set of regular semisimple compact elements.

An equivalent characterization is:

- γ is contained in some subgroup of G which is open, and compact modulo the center of G .

We call a compact element γ *elliptic*, if its fixed point set, X^γ , is compact. Write G^{ell} for the set of regular semisimple elliptic elements.

Equivalent characterizations are:

- γ is not contained in any parabolic subgroup of G .
- $C_G(\gamma)^\circ$ is an elliptic torus (i.e. compact modulo the center of G).

We call γ *non-compact* if it has no fixed points on the building X . (Sometimes non-compact elements are also called *hyperbolic*).

Remark 2.4. Intuitively a non-compact element acts on X by translations, hence has no fixed points on X . Non-compact elements do have fixed points on the spherical building at infinity X^∞ (see [Bro, VI.9]). We will not deal with non-compact elements here. Also note the following.

Let G be a connected reductive group, and (π, V) an admissible representation of G . In [Cas], Casselman shows how to attach to a general element $g \in G^{reg}$, a parabolic subgroup $P = MN$, such that $g \in M$ (g is compact in M). If $(\bar{\pi}, V_N)$ denotes the representation of M on the Jacquet module V_N , then it is shown in [Cas] that $\Theta_\pi(g) = \Theta_{\bar{\pi}}(g)$. That is the calculation of the value of a character at a general element g , can be reduced to the case where g is compact.

Example 2.5. The following elements in $\mathbf{GL}_2(k)$ provide examples of the different types of regular semisimple elements. Here ϖ is a uniformizer.

- elliptic: $\begin{pmatrix} a & b\varpi \\ b & a \end{pmatrix}$, where $a \in k$, $b \in k^\times$ and $a^2 - b^2\varpi \in k^\times$.

The centralizer of this element is an elliptic torus.

⁴We could classify all elements in G , not just those in G^{reg} , into these three types.

- compact, non elliptic: $\begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}$, where $u \in O^\times$ is a unit and $u \neq 1$.

This element is contained in parabolic subgroups and it is easy to show ([Lan, Lemma 5.2]) that its fixed point set contains the basic apartment A .

- non-compact: $\begin{pmatrix} 1 & 0 \\ 0 & \varpi^n \end{pmatrix}$, $n \neq 0$ an integer.

This element acts as a translation and so has no fixed points in X .

Since compact non-elliptic elements play a central role in this paper, we take a closer look at such an element and its fixed point set (see [Lan, chapet 5]).

Let $\gamma = \begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}$, where u is a unit and $u \neq 1$.

Suppose u has the form: $u = 1 + \alpha_r \varpi^r + \dots \in 1 + \varpi^r O^\times \subset O^\times$ $0 \neq \alpha_r \in \bar{k}$. Then the points of X fixed by γ are precisely those at a distance less than or equal to r from the basic apartment A [Lan, Lemma 5.2]. For example, if $r = 1$, then the fixed point set looks like Figure 2.




FIGURE 2. The fixed point set (in bold) of the compact non-elliptic element γ .

2.6. Some subgroups. In [SS2, I.1], Schneider and Stuhler attach to each facet F of the building the following subgroups:

- The stabilizer of the facet: $P_F^\dagger := \{g \in G \mid gF = F\}$.
- The ‘fixer’ of the facet: $P_F := \{g \in G \mid gz = z \text{ for all } z \in F\}$.
 P_F^\dagger and P_F are open subgroups of G ; P_F is a subgroup of finite index in P_F^\dagger .
- A filtration of subgroups: $U_F^{(e)}$, parameterized by integers $e \geq 0$. These are open compact in G and normal in P_F^\dagger .

Sometimes it will be convenient to attach such groups to any point in the building. Let $x \in X$ be any point in X and let F be the unique facet containing x . Define $P_x^\dagger = P_x = \{g \in G \mid gx = x\}$ and $U_x^{(e)} = U_F^{(e)}$.

We summarize some facts about the groups $U_F^{(e)}$ which will be used later. See [SS2, Chapter I] and [Vig, Lemma 1.28]. Here $e \geq 0$ is an integer and F is any facet of the building X .

(U1): $U_F^{(e)}$ are open compact subgroups of G . See [SS2, p.13].

(U2): $U_F^{(e)} \triangleleft P_F^\dagger$. See [SS2, p.21].

(U3): $U_F^{(e)}$, for $e \geq 0$, form a fundamental system of neighborhoods of the identity element, id , in G . See [SS2, I.2 Corollary 9].

(U4): $U_{F'}^{(e)} \subset U_F^{(e)}$ for any two facets F', F in X such that $F' \subset \overline{F}$. See [SS2, I.2 Proposition 11].

(U5): For any two vertices $x, y \in \overline{F}$ and for any $e, e' \geq 0$, the subgroups $U_x^{(e)}$ normalize the subgroups $U_y^{(e')}$. See [SS2, end of section I.2].

(U6):

$$U_F^{(e)} = \prod_{x \text{ vertex in } \overline{F}} U_x^{(e)}$$

for any facet F in X and any ordering of the factors on the right hand side. See [SS2, I.2 Proposition 11].

(U7): Fix two different points x and y in X . Recall that $geod(x, y)$ denotes the (closed) geodesic joining x and y . In an apartment A containing both x and y this geodesic can be realized as

$$geod(x, y) = \{(1-t)x + ty \mid 0 \leq t \leq 1\}.$$

For any point $z \in geod(x, y)$ and for any $e \geq 0$ we have:

$$U_z^{(e)} \subset U_x^{(e)} U_y^{(e)}.$$

See [SS2, I.3 Proposition 1] for x a special vertex; remark after proof of this proposition and [Vig, Lemma 1.28] for x any vertex.

2.7. Representations as coefficient systems. Let $\text{Alg}(G)$ be the category of smooth representations of G . In [SS2, II.2], Schneider and Stuhler define the following objects.

Definition. A *coefficient system* (of complex vector spaces) \underline{V} on the Bruhat-Tits building X consists of:

- complex vector spaces V_F for each facet $F \subset X$, and

- linear maps $r_{F'}^F : V_F \rightarrow V_{F'}$ for each pair of facets $F' \subset \overline{F}$ such that $r_F^F = id$ and $r_{F''}^F = r_{F''}^{F'} \circ r_{F'}^F$, whenever $F'' \subset \overline{F'}$ and $F' \subset \overline{F}$.

Coefficient systems form a category denoted by $\text{Coeff}(X)$.

Fix an integer $e \geq 0$. For any representation V in $\text{Alg}(G)$ we have the coefficient system $\underline{V} := (V^{U_F^{(e)}})_F$. Write $\gamma_e : \text{Alg}(G) \rightarrow \text{Coeff}(X)$ for the functor: $V \rightarrow (V^{U_F^{(e)}})_F$.

Recall that $X_{(q)}$ denotes the space of all oriented q -facets of X . We will denote an oriented facet by (F, c) , where c is an orientation of F (see [SS2, II.1]). For any $F \in X_{q+1}$ and any $F' \subset \overline{F}$ the map $\partial_{F'}^F$ ([SS2, pp. 28–29]), takes an orientation c on F and returns (the induced) orientation $\partial_{F'}^F(c)$, on F' . Often we will abuse the notation and write F , instead of (F, c) , for an oriented facet.

Definition. For any $d \geq q \geq 0$ the space of *oriented q -chains of compact support with values in the coefficient system $\gamma_e(V)$* is:

$C_q := C_c^{or}(X_{(q)}; \gamma_e(V)) := \mathbb{C}$ -vector space of all maps $\omega : X_{(q)} \rightarrow V$ such that

- ω has finite support,
- $\omega((F, c)) \in V^{U_F^{(e)}}$, and
- if $q \geq 1$, $\omega((F, -c)) = -\omega((F, c))$ for all $(F, c) \in X_{(q)}$.

The group G acts *smoothly* on these spaces via

$$(g\omega)((F, c)) := g(\omega((g^{-1}F, g^{-1}c))).$$

There is a natural boundary map

$$\begin{aligned} \partial : C_c^{or}(X_{(q+1)}; \gamma_e(V)) &\longrightarrow C_c^{or}(X_{(q)}; \gamma_e(V)) \\ \omega &\mapsto ((F', c') \mapsto \sum_{\substack{(F, c) \in X_{(q+1)} \\ F' \subset \overline{F} \\ \partial_{F'}^F(c) = c'}} \omega((F, c))). \end{aligned}$$

which satisfies: $\partial \circ \partial = 0$.

We obtain the following augmented G -equivariant chain complex:

$$(*) \quad 0 \longrightarrow C_d \xrightarrow{\partial_d} \dots \xrightarrow{\partial_1} C_0 \xrightarrow{\epsilon} V \longrightarrow 0$$

where the augmentation map is given by

$$\begin{aligned} \epsilon : C_c^{or}(X_{(0)}; \gamma_e(V)) &\longrightarrow V \\ \omega &\mapsto \sum_{F \in X_{(0)}} \omega(F). \end{aligned}$$

Let $V \in \text{Alg}(G)$. In [SS2, II.2-3], Schneider and Stuhler prove the following:

- If V is admissible then $C_c^{or}(X_{(q)}; \gamma_e(V))$ are finitely generated smooth G -modules.
- If V is finitely generated then there exists $e_0 = e_0(V)$ such that the complex $(*)$ is exact for all $e \geq e_0$.

For the rest of this paper V will be a finitely generated admissible representation of G , and e_0 large enough so that the complex $(*)$ is exact.

3. OVERVIEW OF THE SCHNEIDER-STUHLER RESULT

The main result of [SS2] in which we are interested, can be formulated as follows⁵:

For G a connected reductive group, (π, V) a finitely generated admissible representation of G , $\gamma \in G^{ell}$ a regular semisimple elliptic element, and $e \geq e_0(V)$, we have:

$$(1) \quad \Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma)_q} (-1)^q \text{trace}(\gamma; V^{U_F^{(e)}})$$

Here Θ_π is the character (function) of (π, V) evaluated on the regular element γ , $F(\gamma) := F \cap X^\gamma$, and $(X^\gamma)_q$ are the q -facets of X^γ . As stated in [Kot, p.635], $F(\gamma)$ is a polysimplex and X^γ is a polysimplicial complex.

Remark 3.1. Since $\gamma F = F$ for a facet F if and only if $F \cap X^\gamma \neq \emptyset$, the sum over $F(\gamma) \in (X^\gamma)_q$, over all q , is the same as the sum over all the γ -stable facets $F \in X$.

We now loosely sketch (our interpretation) of the proof given by Schneider and Stuhler in [SS2] of this main result. We will sometimes use the phrase ‘for sufficiently large e ’ to mean ‘for all $e \geq e_0$ ’.

There are two main steps:

Step 1: Establish the form of the character formula in terms of the fixed point set X^γ for all *large* e , i.e. $e \geq e(f)$, where $e(f)$ is a constant which depends on f :

$$\int_G f(g) \Theta_\pi(g) dg = \int_G f(g) \left\{ \sum_{q=0}^d \sum_{F(g) \in (X^g)_q} (-1)^q \text{trace}(g; V^{U_F^{(e)}}) \right\} dg$$

Here $f \in C_c^\infty(G)$ has support in G^{ell} so that X^g is compact and the sum on the right hand side is finite.

⁵This result is [SS2, III.4 Lemma 10] combined with [SS2, III.4 Proposition 16]. A cohomological interpretations of this result is given by the Hopf-Lefschetz type trace formula of [SS2, IV.1 Proposition 5].

Step 2: Show that for each fixed $g \in G^{ell}$ the following expression is independent of e (for $e \geq e_0$):

$$\sum_{q=0}^d \sum_{F(g) \in (X^g)_q} (-1)^q \text{trace}(g; V_F^{U_F^{(e)}})$$

Once independence of e has been shown in **Step 2**, we go back to the formula of **Step 1**, whose validity now holds for all $e \geq e_0$. We choose f to have support on a small neighborhood of γ , small enough so that both the character and the alternating sum above are constant on it. Thus we can eliminate f from the equation and obtain formula (1) to hold at an element $\gamma \in G^{ell}$.

The proof of **Step 1** is essentially included in Lemmas 13 and 14 of [SS2, III.4]. We now give a short derivation of the same result using elementary facts.

For simplicity of exposition we assume now that all stable facets are actually fixed facets (see Remark 0.2 and the discussion after Example 4.8).

Fix $\gamma \in G^{cpt}$ and let $F \in X^\gamma$ be a γ -fixed facet. Since $\gamma \in P_F^\dagger$ it acts on $V_F^{U_F^{(e)}}$. Recall Harish-Chandra's formula (section 2.4):

$$\Theta_\pi(f) = \int_G f(g) \Theta_\pi(g) dg \quad f \in C_c^\infty(G)$$

Let $K := U_F^{(e)}$, an open compact neighborhood of the identity, and choose $e(\gamma)$ large enough so that for all $e \geq e(\gamma)$, (the locally constant function) Θ_π is constant on γK : $\Theta_\pi(\gamma K) = \Theta_\pi(\gamma)$. Use $f = \frac{1_{\gamma K}}{\text{vol}(K)}$ in the Harish-Chandra's formula to obtain:

$$\begin{aligned} \Theta_\pi(\gamma) &= \int_K \frac{1}{\text{vol}(K)} \Theta_\pi(\gamma k) dk \\ &= \text{trace}\left(\pi\left(\frac{1_{\gamma K}}{\text{vol}(K)}\right); V\right) \\ &= \text{trace}(\pi(\gamma)\pi(K); V) \\ &= \text{trace}(\pi(\gamma); V^K) \\ &= \text{trace}(\pi(\gamma); V_F^{U_F^{(e)}}) \end{aligned}$$

Remark 3.2. This says that for $\gamma \in G^{cpt}$ (so in particular for $\gamma \in G^{ell}$), all the information about the value of the character at γ is contained in (the vector space $V_F^{U_F^{(e)}}$ above F for) any fixed facet F of γ . But to extract this information we are forced to increase the parameter e . Unfortunately we have no clear control over the way in which e increases.

We now recover the formula of **Step 1**. Recall that for any $\gamma \in G^{cpt}$ its fixed point set, X^γ , is contractible: for two γ -fixed points x, y their geodesic $geod(x, y)$ is also γ -fixed, hence X^γ is (geodesically) contractible; and that for $\gamma \in G^{ell}$ this fixed point set is compact (in fact it is a finite polysimplicial complex).

Let $\gamma \in G^{ell}$. Choose $e = e(\gamma)$ large enough so that $\Theta_\pi(\gamma U_F^{(e)}) = \Theta_\pi(\gamma)$, for all (finitely many) facets $F \in X^\gamma$. Thus $\Theta_\pi(\gamma) = trace(\gamma; V^{U_F^{(e)}})$, for all facets $F \in X^\gamma$. Using the contractibility of X^γ , the alternating sum below collapses and we recover formula (1), which is essentially the formula of **Step 1** (here $Euler(X^\gamma)$ is the Euler characteristic of X^γ):

$$\begin{aligned} \sum_{q=0}^d \sum_{F \in X^\gamma} (-1)^q trace(\gamma; V^{U_F^{(e)}}) &= Euler(X^\gamma) trace(\gamma; V^{U_F^{(e)}}) \\ &= trace(\gamma; V^{U_F^{(e)}}) = \Theta_\pi(\gamma) \end{aligned}$$

Remark 3.3. The only facts that went into the derivation of this formula are the local constancy of the character Θ_π and the contractibility of fixed point sets. In this sense our derivation is elementary. This means that the form of the character formula is not too surprising and that it is in showing that this formula is independent of the parameter e (**Step 2**), where most of the effort is expended.

We now sketch the main ideas in **Step 2**.

Recall the exactness of the resolution $(*)$ for sufficiently large e (i.e. $e \geq e_0$), and the fact that the G -modules $C_q = C_c^{or}(X_q; \gamma_e(V))$ are finitely generated. Both facts are proved in [SS2].

Consider the expression:

$$\sum_{q=0}^d (-1)^q dim H^q(\text{Hom}_G(C_q; V'))$$

Here $V' = (\pi', V')$ is any irreducible admissible representation of G . Since C_q are finitely generated as G -modules, the vector spaces $\text{Hom}_G(C_q; V')$ are finite dimensional and so the expression above makes sense. Since the resolution is exact for all $e \geq e_0$, this expression is independent of e (for $e \geq e_0$).

Using the *Hopf trace formula* and a formal fact about representations of finite groups, the expression above is related to the *Euler-Poincare function*⁶

⁶For definition of the Euler-Poincare function see [SS2, p.45]

$f_{EP}^{V,e}$:

$$\begin{aligned} \sum_{q=0}^d (-1)^q \dim H^q(\mathrm{Hom}_G(C_q; V')) &= \sum_{q=0}^d (-1)^q \dim(\mathrm{Hom}_G(C_q; V')) \\ &= \mathrm{trace}(\pi'(f_{EP}^{V,e}); V') \end{aligned}$$

The left hand side being independent of e implies that the right hand side is also independent of e for all irreducible admissible representations V' .

Recall *Kazhdan's Density Theorem* [Kaz, p.29]:

Theorem. *Let G be a connected reductive p -adic group. Let $f \in C_c^\infty(G)$. Suppose that $\mathrm{trace}(\pi'(f); V') = 0$ for all irreducible admissible representations (π', V') of G . Then $\mathcal{O}_\gamma(f) = 0$ for all strongly regular⁷ semisimple elements $\gamma \in G$.*

Since $\mathrm{trace}(\pi'(f_{EP}^{V,e}); V')$ is independent of e we obtain for any other e' :

$$0 = \mathrm{trace}(\pi'(f_{EP}^{V,e}); V') - \mathrm{trace}(\pi'(f_{EP}^{V,e'}); V') = \mathrm{trace}(\pi'(f_{EP}^{V,e} - f_{EP}^{V,e'}); V')$$

So by applying Kazhdan's Density Theorem, we obtain:

$$0 = \mathcal{O}_\gamma(f_{EP}^{V,e} - f_{EP}^{V,e'}) = \mathcal{O}_\gamma(f_{EP}^{V,e}) - \mathcal{O}_\gamma(f_{EP}^{V,e'}).$$

That is, the orbital integral $\mathcal{O}_\gamma(f_{EP}^{V,e})$ is also independent of e .

A formal calculation, [SS2, III.4 Lemma 10], equates this orbital integral and the alternating sum in equation (1):

$$\mathcal{O}_\gamma(f_{EP}^{V,e}) = \sum_{q=0}^d \sum_{F \in X_q^\gamma} (-1)^q \mathrm{trace}(\gamma; V^{U_F^{(e)}})$$

showing that the alternating sum is independent of e .

4. TRUNCATED BUILDINGS AND TRUNCATION OPERATORS

Fix a point $o \in X$ and call it the *origin*.

In this section we define a family of subsets, X^r ($r \geq 0$, a real number), of the building X . Each of these subsets will be referred to as a *truncated building* (with center o). Using such a truncated building X^r , we will define *truncation operators* Q_q^r on the C_q 's.

Warning: Do not confuse the truncated building X^r with fixed point sets (such as X^g or X^γ) or with X_q for that matter.

Let $B(o, r) := \{x \in X \mid d(o, x) \leq r\}$ be the closed ball of radius r about the point o . Here we denote by $d(\cdot, \cdot)$ the distance function on the building.

⁷A semisimple element is regular if its centralizer has the lowest possible dimension. A semisimple regular element is strongly regular if its centralizer is connected. In general an element is regular/strongly regular if its semisimple part is regular/strongly regular.

Definition 4.1. Let S be any subset of the building. We define two operations on S :

- (1) $cnvx(S) := \text{convex hull of } S$.

We call a subset of the building *convex* if for any two points x, y in the subset, the geodesic $geod(x, y)$ is also in the subset. The convex hull of S is the (unique) smallest convex subset of the building containing S .

Note that if $o \in S \cap X^\gamma$ and S is γ -invariant, then $o \in cnvx(S)$ and $cnvx(S)$ is also γ -invariant.

- (2) $simp(S) := \text{smallest subcomplex of the building containing } S$.

Note that, since the action of γ is simplicial, if $o \in S \cap X^\gamma$ and S is γ -invariant, then $o \in simp(S)$ and $simp(S)$ is also γ -invariant.

Now start with the subset $S = B(o, r)$ and apply the operations (1) and (2) above consecutively to obtain the sequence of increasing subsets:

$$(\dagger) \quad S \subseteq cnvx(S) \subseteq simp(cnvx(S)) \subseteq \dots$$

Claim 4.2. *This sequence stabilizes after a finite number of terms.*

Proof. We assume the results of section 4.1 where it is shown (see Proposition 4.10) that there exist finite polysimplicial subcomplexes of the building which are convex and are arbitrary large (in the sense that they contain any given ball). Starting with $B(o, r)$, fix such a subset E , ($E = X^{\gamma m(r)}$ in the notation of 4.1), containing it. Note that if $S \subseteq E$ then $cnvx(S) \subseteq E$ and also $simp(S) \subseteq E$. Thus the terms in the sequence (\dagger) can never leave the finite subcomplex E and hence the sequence stabilizes. \square

We denote by X^r the stable terms in the sequence (\dagger) and call it the *truncated building (with parameter r)*.

The main properties of the truncated building X^r are:

- $o \in B(o, r) \subseteq X^r$.
- X^r is convex.
- X^r is a finite subcomplex. (In particular it is compact)

Remark 4.3. If $B(o, r)$ contains a chamber, (e.g. r not too small), then X^r is a union of maximal dimensional polysimplices. (See Lemma 4.5).

Example 4.4. For $G = SL_2(\mathbb{Q}_p)$, X is a *tree* with $p + 1$ edges meeting at every vertex. In this case the truncated building X^r , for r an integer, is the closed ball $B(o, r)$. See Figure 3.

To see this recall (see [Bro, IV.3]) that the distance function, $d(o, \cdot)$ on X , is a geodesically convex function⁸.

⁸A function f on X is geodesically convex if for any geodesic $geod(x(0), x(1)) := \{x(t) = tx + (1-t)y \mid 0 \leq t \leq 1\}$: $f(x(t)) \leq tf(x(0)) + (1-t)f(x(1))$ for all $0 \leq t \leq 1$.




FIGURE 3. The truncated building X^r (part of building inside the dotted circle) for SL_2 with $r = 1$ and $r = 2$.

Hence the ball $B(o, r)$ is a convex set, and so applying *convx* to it does nothing. Now for $G = SL_2$, the ball $B(o, r)$ is already simplicial, so applying *simp* to it also does nothing.

The truncated building X^r induces a decomposition of each oriented q -skeletons as a disjoint union:

$$X_{(q)} = X_{(q)}^{in(r)} \amalg X_{(q)}^{out(r)},$$

where $X_{(q)}^{in(r)} := \{F \in X_{(q)} \mid F \subset X^r\}$ and $X_{(q)}^{out(r)} := \{F \in X_{(q)} \mid F \not\subset X^r\}$.

Let $C_q^{in(r)} :=$ the oriented q -chains in C_q supported on $X_{(q)}^{in(r)}$ and

let $C_q^{out(r)} :=$ the oriented q -chains in C_q supported on $X_{(q)}^{out(r)}$.

Note that $C_q^{in(r)} = C_q(X^r)$.

We obtain a vector space direct sum decomposition:

$$C_q = C_q^{in(r)} \oplus C_q^{out(r)}$$

Define the *truncation operator* Q_q^r on C_q to be the projection of C_q onto $C_q^{in(r)}$. Thus the truncation operator Q_q^r takes a map $\omega \in C_q^{or}(X_{(q)}; \gamma_e(V))$ and returns a map $Q_q^r \omega \in C_q^{or}(X_{(q)}; \gamma_e(V))$ supported on X^r .

Note the following properties of the truncation operators Q_q^r :

- Q_q^r is a finite rank operator.

This follows from the fact that its range $C_q^{in(r)}$ is a finite dimensional vector space.

- The Q_q^r 's do not necessarily commute with the boundary maps ∂ .

Notation: Write Q^r for the sequence of operators (Q_d^r, \dots, Q_0^r) , where each operator Q_q^r acts on the corresponding vector space C_q .

We would like to modify the truncation operators so that they *do* commute with the boundary maps ∂ . This is done in section 5.

4.1. Existence of certain finite subcomplexes. We now prove the existence of finite subcomplexes of the building which are convex and are arbitrary large (in the sense that they contain any given ball). This section is independent of previous results.

Lemma 4.5. *Let $\gamma \in G = \mathbf{G}(k)$ be a compact element of G . Then its fixed point set, X^γ , is a convex subset of the building X . If γ is elliptic, then X^γ is also compact.*

Proof. Convexity follows from the fact that for any two points $x, y \in X$ fixed by γ , their geodesic $geod(x, y)$ is also fixed by γ . Compactness is proved in Lemma 1 of [Rog]. \square

Lemma 4.6. *For any compact element $\gamma \in G$, if its fixed point set X^γ contains a chamber C , then X^γ is the union of polysimplices of maximal dimension; equivalently X^γ is the union of the closure of its chambers.*

Proof. Pick any element $x \in X^\gamma$. Choose an apartment A containing both x and the chamber C . Recall that any apartment is a Euclidean space (over \mathbb{R}) with the usual topology. Let $cone(C, x)$ be the cone on C with vertex x . Since both C and $\{x\}$ are subsets of X^γ , so is $cone(C, x)$. Let $star^\circ(x)$ denote the union of all the chambers C' in A which contain x in their closure. Both $cone(C, x)$ and $star^\circ(x)$ are open subsets of the Euclidean space A and so is their non-empty intersection. Let $C' \subseteq star^\circ(x)$ be a chamber which intersects $cone(C, x)$. This intersection is fixed under γ and it being an open subset of C' , forces C' to also be fixed by γ . Thus $x \in \overline{C'}$. \square

Remark 4.7. For an arbitrary elliptic element γ , its fixed point set X^γ need not be a subcomplex of the building.

Example 4.8. In $G = PGL_2(\mathbb{Q}_2)$, which has the same Bruhat-Tits building as $SL_2(\mathbb{Q}_2)$, the fixed point set of the elliptic element

$$\gamma = \left[\left(\begin{array}{cc} 0 & 1 \\ \varpi & 0 \end{array} \right) \right]$$

is a single point: the center of a standard chamber. This is not a subcomplex of the building.

For this reason we will restrict ourselves⁹ to elliptic elements $\gamma \in \mathcal{D}G = \mathcal{D}\mathbf{G}(k)$ which are coming from the simply connected cover¹⁰ $\tilde{G} = \tilde{\mathbf{G}}(k)$ of G . That is, let $\tilde{\gamma}$ be an elliptic element in \tilde{G} and consider its image γ under the natural map from \tilde{G} to $\mathcal{D}G$. (Recall that the projection map $\pi : \tilde{\mathbf{G}} \rightarrow \mathcal{D}\mathbf{G}$, which is a map on the level of algebraic groups, induces a map $\pi : \tilde{\mathbf{G}}(k) \rightarrow \mathcal{D}\mathbf{G}(k)$ on the level of k -rational points and that the image $\pi(\tilde{\mathbf{G}}(k))$ is a subgroup of $\mathcal{D}\mathbf{G}(k)$ of finite index.) The action of the simply connected cover \tilde{G} preserves the type (See [Tit, 2.5]) of each vertex (do not confuse type of a vertex with type of an element). Since any facet contains at most one vertex of each type, any element $\gamma \in \tilde{G}$ which stabilizes a facet must fix this facet. That is, if an element $\gamma \in \tilde{G}$ fixes a point $x \in X$, then it fixes every point of the unique facet in which x lies. Thus the fixed point sets X^γ of such elements are subcomplexes.

Let $\tilde{\gamma}_m \in \tilde{G}$ be a sequence of regular elliptic elements going to $id \in \tilde{G}$. Denote their images in $\mathcal{D}G$ by γ_m .

Lemma 4.9. *For each $r \geq 0$, there is a positive integer $m = m(r)$ such that γ_m fixes all points of $B(o, r)$.*

Proof. Consider the open subgroup $U := \{g \in \mathcal{D}G \mid gx = x \text{ for all } x \in B(o, r)\}$. For large enough $m = m(r)$, the element γ_m is inside U and so fixes all the points of $B(o, r)$ \square

Combining the last two lemmas we obtain:

Proposition 4.10. *For any ball $B(o, r)$ the subset $X^{\gamma_{m(r)}}$ contains the ball. It is a convex, finite subcomplex of X . If $B(o, r)$ contains a chamber, $X^{\gamma_{m(r)}}$ is a union of maximal dimensional polysimplices.*

5. MODIFIED TRUNCATION OPERATORS

Some of the ideas in this section were inspired by [AB1] and [AB2].

Fix direct sum decompositions of the vector spaces C_q ($d \geq q \geq 0$):

$$(\oplus) \quad C_q = Z_q \oplus B'_{q-1} = B_q \oplus H'_q \oplus B'_{q-1}$$

Here Z_q are the q -cycles and B'_{q-1} is a complement to Z_q inside C_q . The spaces $B_q := \partial(C_{q+1})$ are the q -boundaries, and so $B'_{q-1} \cong B_{q-1}$. The space H'_q is a complement of B_q inside Z_q (so that $H'_q \cong H_q$). Note that such decompositions depend on the parameter e – this dependence is suppressed in the notation here. Given the sequence of truncation operators

⁹Recall that the groups G , \tilde{G} , and $\mathcal{D}G$ all have the same semisimple Bruhat-Tits building and their actions are compatible.

¹⁰By definition the simply connected cover \tilde{G} of a reductive group G is the simply connected cover of its derived group $\mathcal{D}G$.

$Q^r = (Q_d^r, \dots, Q_0^r)$ we use the decompositions (\oplus) to define *modified truncation operators* \widetilde{Q}_q^r on the C_q 's. In contrast to the truncation operators, the modified truncation operators are constructed in such a way as to commute with the boundary maps: $\partial_q \circ \widetilde{Q}_q^r = \widetilde{Q}_{q-1}^r \circ \partial_q$. Thus $\overline{Q}^r := (\overline{Q}_d^r, \dots, \overline{Q}_0^r)$ is an *endomorphism* of the complex

$$(C) \quad 0 \longrightarrow C_d \xrightarrow{\partial_d} \dots \xrightarrow{\partial_1} C_0 \longrightarrow 0$$

For this construction we first use the isomorphism $B'_{q-1} \xrightarrow[\sim]{\partial} B_{q-1}$ to define operators \widetilde{Q}_{q-1}^r on B'_{q-1} in such a way that the following diagram commutes:

$$\begin{array}{ccc} B'_{q-1} & \xrightarrow[\sim]{\partial} & B_{q-1} \\ \downarrow \widetilde{Q}_{q-1}^r & & \downarrow Q_{q-1}^r[B_{q-1}; B_{q-1}] \\ B'_{q-1} & \xrightarrow[\sim]{\partial} & B_{q-1} \end{array}$$

Now if we represent Q_q^r on $C_q = B_q \oplus H'_q \oplus B'_{q-1}$ by the matrix:

$$Q_q^r = \begin{pmatrix} Q_q^r[B_q; B_q] & Q_q^r[B_q; H'_q] & Q_q^r[B_q; B'_{q-1}] \\ Q_q^r[H'_q; B_q] & Q_q^r[H'_q; H'_q] & Q_q^r[H'_q; B'_{q-1}] \\ Q_q^r[B'_{q-1}; B_q] & Q_q^r[B'_{q-1}; H'_q] & Q_q^r[B'_{q-1}; B'_{q-1}] \end{pmatrix}$$

then we define \overline{Q}_q^r by the matrix:

$$\overline{Q}_q^r := \begin{pmatrix} Q_q^r[B_q; B_q] & Q_q^r[B_q; H'_q] & Q_q^r[B_q; B'_{q-1}] \\ 0 & Q_q^r[H'_q; H'_q] & Q_q^r[H'_q; B'_{q-1}] \\ 0 & 0 & \widetilde{Q}_{q-1}^r \end{pmatrix}$$

Claim 5.1. *The modified truncation operators commute with the boundary operators: $\partial_q \circ \overline{Q}_q^r = \overline{Q}_{q-1}^r \circ \partial_q$ for $d \geq q \geq 1$.*

Proof. Write $\omega \in C_q = Z_q \oplus B'_{q-1}$ as $\omega = z + b'$, where $z \in Z_q$ and $b' \in B'_{q-1}$. Since $\overline{Q}_q^r z \in Z_q$, applying ∂ to it gives $\partial \overline{Q}_q^r z = 0$. Hence we obtain:

$$\begin{aligned} \partial \overline{Q}_q^r \omega &= \partial \overline{Q}_q^r z + \partial \overline{Q}_q^r b' &= \partial \overline{Q}_q^r b' \\ & &= \partial \widetilde{Q}_{q-1}^r b' \\ & &= Q_{q-1}^r[B_{q-1}; B_{q-1}] \partial b' \\ & &= \overline{Q}_{q-1}^r \partial b' = \overline{Q}_{q-1}^r \partial \omega \end{aligned}$$

□

Claim 5.2. *The modified truncation operators \overline{Q}_q^r have finite rank.*

Proof. Since the original truncation operators Q_q^r have finite rank, the operators $Q_q^r[*; \star]$ all have finite rank (since they are all of the form $Q_q^r[*; \star] = P_\star Q_q^r P_*$, where P_* and P_\star are projection operators). Note that since \widetilde{Q}_{q-1}^r acts the same (under the above isomorphism) as $Q_{q-1}^r[B_{q-1}; B_{q-1}]$, it also

has finite rank. Now, since all the operator entries in the matrix defining \overline{Q}_q^r have finite rank, this modified truncation operator also has finite rank. \square

Claim 5.3. *The modified truncation operators \overline{Q}_q^r tend to Id_{C_q} , as r tends to ∞ , in the following sense:*

$$\forall \omega \in C_q \quad \exists r_\omega \in \mathbb{R} \quad \text{s.t.} \quad \overline{Q}_q^r(\omega) = \omega \quad \forall r \geq r_\omega.$$

Proof. Given $\omega \in C_q$, write its components with respect to $C_q = B_q \oplus H'_q \oplus B'_{q-1}$ as $\omega = \omega_1 + \omega_2 + \omega_3$. Let r_ω be large enough so that $\text{support}(\omega_i) \subset B(o, r_\omega)$, for $1 \leq i \leq 3$. We have

$$\begin{aligned} \omega_1 &= Q_q^r \omega_1 = Q_q^r[B_q; B_q] \omega_1 + Q_q^r[H'_q; B_q] \omega_1 + Q_q^r[B'_{q-1}; B_q] \omega_1, \\ \omega_2 &= Q_q^r \omega_2 = Q_q^r[B_q; H'_q] \omega_2 + Q_q^r[H'_q; H'_q] \omega_2 + Q_q^r[B'_{q-1}; H'_q] \omega_2, \\ \omega_3 &= Q_q^r \omega_3 = Q_q^r[B_q; B'_{q-1}] \omega_3 + Q_q^r[H'_q; B'_{q-1}] \omega_3 + Q_q^r[B'_{q-1}; B'_{q-1}] \omega_3. \end{aligned}$$

Since the decomposition above is a direct sum decomposition, we see that

$$\begin{aligned} \omega_1 &= Q_q^r \omega_1 = Q_q^r[B_q; B_q] \omega_1 + 0 + 0, \\ \omega_2 &= Q_q^r \omega_2 = 0 + Q_q^r[H'_q; H'_q] \omega_2 + 0, \\ \omega_3 &= Q_q^r \omega_3 = 0 + 0 + Q_q^r[B'_{q-1}; B'_{q-1}] \omega_3. \end{aligned}$$

Thus by the definition of \overline{Q}_q^r we have

$$\begin{aligned} \overline{Q}_q^r \omega_1 &= Q_q^r[B_q; B_q] \omega_1 = \omega_1, \\ \overline{Q}_q^r \omega_2 &= Q_q^r[H'_q; H'_q] \omega_2 = \omega_2, \\ \overline{Q}_q^r \omega_3 &= \overline{Q}_q^r \omega_3. \end{aligned}$$

Note that since $\text{support}(\omega_3) \subset B(o, r_\omega)$ also $\text{support}(\partial \omega_3) \subset B(o, r_\omega)$. Hence $\partial \overline{Q}_q^r \omega_3 = Q_{q-1}^r \partial \omega_3 = \partial \omega_3$, and since $B'_{q-1} \xrightarrow[\sim]{\partial} B_{q-1}$, we obtain $\overline{Q}_q^r \omega_3 = \overline{Q}_q^r \omega_3 = \omega_3$. We see that for $1 \leq i \leq 3$, $\overline{Q}_q^r \omega_i = \omega_i$. That is $\overline{Q}_q^r \omega = \omega$. \square

We summarize the main properties of the modified truncation operators \overline{Q}_q^r :

- \overline{Q}_q^r is a finite rank operator.
- The \overline{Q}_q^r 's commute with the boundary maps ∂ .
- \overline{Q}_q^r tends to Id_{C_q} , as r tends to ∞ .

Hence \overline{Q} is an endomorphism of finite rank of the complex (\mathcal{C}) .

Remark 5.4. Since the complex $(*)$ is exact, the complex (\mathcal{C}) is exact at C_q , $d \geq q \geq 1$, and so:

$$\begin{aligned} C_0 &= B_0 \oplus H'_0 \\ C_q &= B_q \oplus B'_{q-1}, \quad d \geq q \geq 1. \end{aligned}$$

6. A CHARACTER FORMULA FOR FUNCTIONS

Let $f \in C_c^\infty(G)$ be a locally constant function of compact support on G . Recall that the vector spaces C_q are *smooth* G -modules, and let T_g denote the action of g on the C_q 's. As is usual for smooth representations, define the operators $T_f = (T_f)_q$ on the C_q 's:

$$T_f(\omega) := \int_G f(g)T_g(\omega) dg, \quad \omega \in C_q$$

Note that $T_f = ((T_f)_d, \dots, (T_f)_0)$ is an endomorphism of the complex (\mathcal{C}) and that the operators $(T_f)_q$ are not necessarily of finite rank: the representation C_q is smooth but not necessarily admissible.

Since \overline{Q}^r is a finite rank endomorphism of the complex $\mathcal{C} = (C_d, \dots, C_0)$, the composition $T_f \overline{Q}^r = (T_f \overline{Q}_d^r, \dots, T_f \overline{Q}_0^r)$ is also finite rank endomorphism. Recall that any endomorphism of a complex induces operators on its homology modules. We use the same notation $T_f \overline{Q}_q^r$ for the induced operators on $H_q := H_q(\mathcal{C})$.

Theorem 6.1. *For G connected reductive, V finitely generated admissible, $e \geq e_0$, and $f \in C_c^\infty(G)$ there exists a radius $r_0(f)$ large enough¹¹ so that for all $r \geq r_0(f)$:*

$$\sum_{q=0}^d (-1)^q \text{trace}(T_f \overline{Q}_q^r; C_q) = \text{trace}(\pi(f); V)$$

Proof.

$$\begin{aligned} \sum_{q=0}^d (-1)^q \text{trace}(T_f \overline{Q}_q^r; C_q) &= \sum_{q=0}^d (-1)^q \text{trace}(T_f \overline{Q}_q^r; H_q) \\ &= \text{trace}(T_f \overline{Q}_0^r; H_0) \\ &= \text{trace}(T_f; H_0) \\ &= \text{trace}(\pi(f); V) \end{aligned}$$

Applying the Hopf trace formula for finite rank operators [AB1, Proposition 2.1] to the finite rank endomorphism $T_f \overline{Q}^r$ we obtain the first equality. Since the complex (\mathcal{C}) is exact at all places C_q , $d \geq q \geq 1$, the only (possible) non-zero homology is H_0 . This explains the second equality. Using the exact G -equivariant complex $(*)$ we have that $H_0 = C_0/B_0 \cong V$ and hence that $\text{trace}(T_f; H_0) = \text{trace}(\pi(f); V)$. Note that since V is admissible the operator $\pi(f)$ is of finite rank and so taking its trace makes sense. This gives the fourth equality. The following explains the third equality.

¹¹Choose $\omega_1, \dots, \omega_n \in H'_0 \subset C_0$ such that $\{\omega_1 B_0, \dots, \omega_n B_0\} \subset H_0 = C_0/B_0$ is a basis for $W := T_f(H_0)$. Taking ' r large enough' means that $\omega_i \in C_c^{or}(X_{(0)}^r; \gamma_e(V))$, for all $0 \leq i \leq n$.

Consider the diagram:

$$\begin{array}{ccc} H_0 & \xrightarrow{\epsilon} & V \\ \downarrow T_f & \sim & \downarrow \pi(f) \\ H_0 & \xrightarrow{\epsilon} & V \end{array}$$

Since the exact complex (*) respects the action of g , for all $g \in G$, it also respects the action of f ; therefore the diagram above commutes. Since V is admissible, $\pi(f)$ is a finite rank operator on V . Hence T_f is also a finite rank operator on H_0 . To show

$$\text{trace}(T_f \overline{Q}_0^r; H_0) = \text{trace}(T_f; H_0),$$

it is enough to show that for large enough r , the operators \overline{Q}_0^r act trivially on the finite dimensional image $T_f(H_0)$ of T_f .

Choose $\omega_1, \dots, \omega_n \in H_0' \subset C_0$ such that $\{\omega_1 B_0, \dots, \omega_n B_0\} \subset H_0 = C_0/B_0$ is a basis for $W := T_f(H_0)$, and let W' be a complement of W inside H_0 : $H_0 = W' \oplus W$. With respect to this direct sum decomposition the operator T_f is represented by a matrix of the form:

$$T_f = \begin{pmatrix} 0 & 0 \\ * & T_{W,W} \end{pmatrix}.$$

Let r be large enough so that $\omega_i \in C_c^{or}(X_{(0)}^r; \gamma_\epsilon(V))$, for all $0 \leq i \leq n$. Then (Claim 5.3) $\overline{Q}_0^r(\omega_i) = \omega_i$. Thus $\overline{Q}_0^r(\omega_i B_0) = \overline{Q}_0^r(\omega_i) B_0 = \omega_i B_0$, and so

$$\overline{Q}_0^r|_W \equiv Id|_W.$$

Note that r depends on W which depends on f .

So with respect to the above decomposition of H_0 the operator \overline{Q}_0^r is represented by a matrix of the form:

$$\overline{Q}_0^r = \begin{pmatrix} * & 0 \\ * & 1 \end{pmatrix}.$$

Multiply T_f and \overline{Q}_0^r to obtain:

$$T_f \overline{Q}_0^r = \begin{pmatrix} 0 & 0 \\ * & T_{W,W} \end{pmatrix} \begin{pmatrix} * & 0 \\ * & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ * & T_{W,W} \end{pmatrix}.$$

Equality of the traces follows:

$$\text{trace } T_f \overline{Q}_0^r = \text{trace } T_f$$

This completes the proof of the theorem. \square

How to proceed. Keeping in mind the formula of Theorem 6.1, we have the following two objectives in mind:

- (1) To give a more geometric interpretation of this formula in terms of the original truncation operators.
- (2) To replace the function f in the formula with a compact element γ .

In the rest of this paper we show how to complete step (1) for groups of semisimple rank 1; we make some progress in the direction of showing (1) for a general reductive group. Assuming step (1) we show how to complete step (2) for all compact elements. For groups of semisimple rank 1, we recover the Schneider-Stuhler result.

7. NICE DECOMPOSITIONS

Let $\gamma \in G^{cpt}$ and fix $o \in X^\gamma$. Recall the decomposition: $C_q = C_q^{in(r)} \oplus C_q^{out(r)}$ which is T_γ -equivariant (see paragraph after Example 4.4). Let $B_q^{in(r)} := B_q \cap C_q^{in(r)}$. It is a T_γ -stable subspace of B_q as T_γ acts on B_q and on $C_q^{in(r)}$. Let $B_q^{out(r)}$ be any T_γ -stable complement of $B_q^{in(r)}$ in B_q (we will show existence of such complements later in this section).

Definition 7.1. We say that the decomposition (\oplus) is ‘nice’ relative to $\gamma \in G^{cpt}$ and $r \in \mathbb{R}$, (here γ is such that $o \in X^\gamma$), if it has the following form:

- $C_0 = B_0 \oplus H'_0$
- $C_q = B_q \oplus B'_{q-1}$, $d \geq q \geq 1$.
Where $B'_{q-1} = (B_{q-1}^{in(r)})' \oplus (B_{q-1}^{out(r)})'$, for some subspaces $(B_{q-1}^{in(r)})'$ and $(B_{q-1}^{out(r)})'$ which satisfy:
 $(B_{q-1}^{in(r)})' \xrightarrow[\sim]{\partial} (B_{q-1}^{in(r)})'$, $(B_{q-1}^{out(r)})' \xrightarrow[\sim]{\partial} (B_{q-1}^{out(r)})'$, and
 $(B_{q-1}^{in(r)})' \subset C_q^{in(r)}$.
- All \oplus decompositions here are T_γ -equivariant.

In this definition the dependence on the parameter e is suppressed in the notation. If for each $r \in \mathbb{R}$ nice decompositions exist for all $e \geq e_0$, we will say that the nice decomposition is *uniform in e* . If for each r nice decompositions exist only for all $e \geq e_r$, where e_r is an integer depending in r , we will say that the nice decomposition *depends on e_r* .

In this section we show that for groups of semisimple rank 1, such nice decompositions exist uniformly in e . For a general connected reductive group we show that nice decompositions exist, but that they depend on e_r .

When the decomposition (\oplus) is nice, we will show, under appropriate conditions, that:

$$\text{trace}(T_\gamma Q_d^r; C_q) = \text{trace}(T_\gamma \overline{Q}_d^r; C_q) \quad d \geq q \geq 0$$

7.1. Truncated complexes. For a general group G the direct sum decomposition is controlled by the following *truncated complex*:

$$(*) \quad 0 \longrightarrow C_c^{or}(X_{(d)}^r; \gamma_e(V)) \xrightarrow{\partial} \cdots \xrightarrow{\partial} C_c^{or}(X_{(0)}^r; \gamma_e(V)) \xrightarrow{\epsilon} V$$

The exactness of this complex will guarantee the existence of a nice direct sum decomposition (\oplus). The exactness will follow from the properties of the truncated building X^r . Ideally we would like to show exactness of ($\bar{*}$) for all $e \geq e_0$, but the techniques used here will show (for a general connected reductive group) exactness only for $e \geq e_r$, for some e_r which depends on r .

We recall the following averaging process: Let $K \subset G$ be an open compact subgroup and denote by 1_K its characteristic function. The operator $\pi(K) := \pi(1_K)$ on V is well defined as $1_K \in C_c^\infty(G)$:

$$\pi(K)v = \int_{k \in K} \pi(k)v dk, \quad v \in V$$

where dk is a normalized Haar measure on K .

Note the following facts about $\pi(K)v$:

- For any $v \in V$, $\pi(K)v \in V^K$.
This follows from the definition of $\pi(K)$.
- If for some open compact subgroup U , $v \in V^U$ and K normalizes U , then $\pi(K)v \in V^U$:

$$\begin{aligned} \pi(u)\pi(K)v &= \pi(u) \int_{k \in K} \pi(k)v dk = \int_{k \in K} \pi(u)\pi(k)v dk \\ &= \int_{k \in K} \pi(k)\pi(k^{-1})\pi(u)\pi(k)v dk = \int_{k \in K} \pi(k)\pi(k^{-1}uk)v dk \\ &= \int_{k \in K} \pi(k)\pi(u)v dk = \int_{k \in K} \pi(k)v dk = \pi(K)v \end{aligned}$$

Recall that for any facet F , the groups $U_F^{(e)}$ are normal subgroups of P_F^\dagger . For vertices x this means that $U_x^{(e)} \triangleleft P_x = P_x^\dagger$.

Let X^r be the truncated building with center $o \in X$.

Claim 7.2. *It is possible to choose an integer e_r large enough so that for any $e \geq e_r$ the groups $U_x^{(e)}$, for $x \in X^r$, all normalize each other.*

Proof. Let $U(r) := \bigcap_{y \in X^r} U_y^{(0)}$, an open compact subgroup (since this intersection is a finite intersection of open compact subgroups). Since for any vertex x , the subgroups $U_x^{(e)}$ form a filtration of the identity element in G , we can choose e_r large enough so that $U_x^{(e)} \subset U(r) \subset P_y$, for all $x, y \in X^r$ and $e \geq e_r$. Since $U_x^{(e)} \subset P_y$, it normalizes $U_y^{(e)}$. As this holds for all $x, y \in X^r$ and $e \geq e_r$, the claim is proved. \square

Theorem 7.3. *For $e \geq e_r$, the sequence ($\bar{*}$) is exact.*

Proof. Take a non-zero cycle $\omega \in C_c^{or}(X_{(q)}^r; \gamma_e(V))$; we need to show that it is a boundary. The proof is by induction on the number of facets in the support of ω .

We can use any open compact subgroup, $K \subset U(r)$, to average ω :

$$(\pi(K)\omega)((F', c')) = \int_{k \in K} \pi(k)(\omega((k^{-1}F', k^{-1}c'))) dk$$

Since $K \subset P_{F'}$ for all $F' \subset X^r$, $k \in K$ acts trivially on $(F', c') \in X_q^r$. Hence

$$(\pi(K)\omega)((F', c')) = \int_{k \in K} \pi(k)(\omega((F', c'))) dk.$$

Note that $(\pi(K)\omega)((F', c')) \in V^K \cap V^{U_{F'}^{(e)}}$ and that $\pi(K)\omega \neq 0$. Also, since the action of $K \subset G$ commutes with the boundary operators, $\pi(K)\omega$ is also a cycle.

Let $F \in X_{(q)}^r$ be an oriented q -facet such that $F \subset \text{support}(\omega)$. Use $K = U_F^{(e)}$. Then $\omega = \pi(U_F^{(e)})\omega + (\omega - \pi(U_F^{(e)})\omega)$. Being a difference of two cycles, $(\omega - \pi(U_F^{(e)})\omega)$ is itself a cycle. Note that $\omega(F) \in V^{U_F^{(e)}}$ implies that $(\pi(U_F^{(e)})\omega)(F) = \omega(F)$, hence $(\omega - \pi(U_F^{(e)})\omega)(F) = 0$. This means that the support of $\omega - \pi(U_F^{(e)})\omega$ is strictly smaller than the support of ω . Thus by induction we conclude that $\omega - \pi(U_F^{(e)})\omega$ is a boundary. The non-zero cycle $\pi(U_F^{(e)})\omega$ has the same support as ω and is invariant (by the averaging process) under $U_F^{(e)}$. Since $e \geq e_r$, the groups $U_F^{(e)}$, $F \in X_{(q)}^r$, all normalize each other, and so $\pi(U_{F'}^{(e)})\pi(U_F^{(e)})\omega$ is still $U_F^{(e)}$ -invariant. Thus after applying this process a finite number of times, i.e. as K ranges over all $F \subset \text{support}(\omega)$, we can assume with out loss of generality that the cycle ω is invariant under all $U_F^{(e)}$, $F \subset \text{support}(\omega)$.

Denote by $X_0(\omega)$ the set of vertices in $\overline{\text{support}(\omega)}$ and let

$$V_\omega^{(e)} := \bigcap_{x \in X_0(\omega)} V^{U_x^{(e)}},$$

then $\omega((F', c')) \in V_\omega^{(e)}$, all $(F', c') \subset X^r$.

We show that this situation can be reduced to a constant coefficients case.

Let $S \subset X$ be any subset of the building. We recall the *simplicial* operation and define an algorithmic version of the *convex* operation:

(1) $geod(S) := \{z \in X \mid z \in geod(x, y) \text{ for some } x, y \in S\}$.

(2) $simp(S) := \text{smallest subcomplex of the building containing } S$.

Lemma 7.4. *If $V_\omega^{(e)} \subset V_z^{U_z^{(e)}}$, for all $z \in S$ then*

(1) $V_\omega^{(e)} \subset V_z^{U_z^{(e)}}$, for all $z \in geod(S)$.

(2) $V_\omega^{(e)} \subset V_z^{U_z^{(e)}}$, for all $z \in simp(S)$.

Proof. For $z \in geod(x, y)$ have $U_z^{(e)} \subset U_x^{(e)}U_y^{(e)}$ [Vig, Lemma 1.28]. Hence $V_z^{U_z^{(e)}} \supset V_x^{U_x^{(e)}} \cap V_y^{U_y^{(e)}} \supset V_\omega^{(e)}$, which proves (1). If $z \in simp(S)$, then $z \in \overline{F}$ for some facet F s.t $F \cap S \neq \emptyset$. So (by property (U4)) $U_z^{(e)} \subset U_F^{(e)} = U_x^{(e)}$ for any $x \in F \cap S$ and hence $V_\omega^{(e)} \subset V_x^{U_x^{(e)}} \subset V_z^{U_z^{(e)}}$, which proves (2). \square

We continue with the proof of the theorem, where X^r is the truncated building with parameter r and we choose e_r as in the claim above. Let $\omega \in C_c^{or}(X_{(q)}^r; \gamma_{e_r}(V))$ be a q -cycle. We want to show it is a boundary. Let $S_\omega := supp(\omega)$ and consider the sequence of increasing subsets:

$$(\ddagger) \quad S_\omega \subseteq geod(S_\omega) \subseteq simp(geod(S_\omega)) \subseteq \dots$$

Essentially the same argument as that showing that the sequence (\dagger) stabilizes, shows that this sequence (\ddagger) also stabilizes after a finite number of terms. Denote by \overline{S}_ω the stable terms in the sequence (\ddagger) .

It follows from the construction of \overline{S}_ω that $\overline{S}_\omega \subset X^r$, and that \overline{S}_ω is simplicial and convex.

Also since $V_\omega^{(e)} \subset V_x^{U_x^{(e)}}$, for all $x \in S_\omega$, we can apply the lemma several times to conclude that $V_\omega^{(e)} \subset V_x^{U_x^{(e)}}$, for all $x \in \overline{S}_\omega$. Now consider the following commutative diagram of chain complexes:

$$\begin{array}{ccccccc} \xrightarrow{\partial} C_c^{or}(X_{(q+1)}^r; \gamma_{e_r}(V)) & \xrightarrow{\partial} & C_c^{or}(X_{(q)}^r; \gamma_{e_r}(V)) & \xrightarrow{\partial} & C_c^{or}(X_{(q-1)}^r; \gamma_{e_r}(V)) & \xrightarrow{\partial} & \\ \cup & & \cup & & \cup & & \\ \xrightarrow{\partial} C_c^{or}((\overline{S}_\omega)_{(q+1)}^r; V_\omega^{(e)}) & \xrightarrow{\partial} & C_c^{or}((\overline{S}_\omega)_{(q)}^r; V_\omega^{(e)}) & \xrightarrow{\partial} & C_c^{or}((\overline{S}_\omega)_{(q-1)}^r; V_\omega^{(e)}) & \xrightarrow{\partial} & \end{array}$$

The bottom line is the homology chain complex of the simplicial complex \overline{S}_ω with constant coefficients $V_\omega^{(e)}$. Since \overline{S}_ω is convex (hence contractible) its constant coefficient chain complex is exact. Hence ω , considered as a cycle in $C_c^{or}((\overline{S}_\omega)_{(q)}^r; V_\omega^{(e)})$ must be a boundary of some chain $\delta \in C_c^{or}((\overline{S}_\omega)_{(q+1)}^r; V_\omega^{(e)}) \subset C_c^{or}(X_{(q+1)}^r; \gamma_{e_r}(V))$. We have found $\delta \in C_c^{or}(X_{(q+1)}^r; \gamma_{e_r}(V))$ such that $\partial\delta = \omega$, i.e., the cycle ω is a boundary. This concludes the proof of the theorem. \square

7.2. Existence of nice decompositions. Let γ be a compact element such that $o \in X^\gamma$. The compact element γ is contained in some compact subgroup K of G . Let $K_\gamma := \overline{\langle \gamma \rangle}$ be the closure in K of the subgroup generated by γ . K_γ is a compact subgroup containing γ . It will be used for ‘averaging’ purposes.

Remark 7.5. Schneider-Stuhler make the assumption that $Z^\circ(G)$ is anisotropic which implies that the stabilizer P_x , of any vertex x , is a compact group. In particular P_o being a compact group containing γ could also be used for ‘averaging’ purposes. We will not need it here, so we will not need to make the above assumption on G .

Recall that for $B_q^{out(r)}$ we needed to take any T_γ -stable complement of $B_q^{in(r)}$ in B_q .

To see such a complement exists, take any projection $\pi : B_q \rightarrow B_q^{in(r)}$ and average it over the compact group K_γ :

$$\int_{K_\gamma} T_k^{-1} \circ \pi \circ T_k$$

This makes sense since K_γ acts on B_q and on $B_q^{in(r)}$. The result is a T_γ -equivariant projection onto $B_q^{in(r)}$. Let $B_q^{out(r)}$ be the kernel of this projection.

Get a T_γ -equivariant direct sum decomposition of B_q for each $d \geq q \geq 0$:

$$B_q = B_q^{in(r)} \oplus B_q^{out(r)}$$

Note that $B_q^{in(r)} \subset C_q^{in(r)}$ but $B_q^{out(r)} \not\subset C_q^{out(r)}$.

Recall that since the chain complex \mathcal{C} is exact at C_q , $d \geq q \geq 1$, the direct sum decomposition (\oplus) has the form :

$$\begin{aligned} C_0 &= B_0 \oplus H'_0 \\ C_q &= B_q \oplus B'_{q-1}, \quad d \geq q \geq 1. \end{aligned}$$

Here H'_0 is any T_γ -equivariant complement of B_0 in C_0 . As above, such a complement can be realized as the kernel of an appropriate T_γ -equivariant projection from C_0 to B_0 which can be produced by averaging over K_γ . For B'_{q-1} we will chose a particularly ‘nice’ section of ∂ in the following sense.

For $q \geq 1$ we will construct a particular section α , i.e. $\partial \circ \alpha = id$, of the surjective map $\partial : C_q \rightarrow B_{q-1}$ and then use it to define B'_{q-1} as the image $\alpha(B_{q-1})$.

Example 7.6. Let $G = SL_2$ and $V = \mathbb{C}$ be the trivial representation. The building X is a tree. The chain complex (\mathcal{C}) for X is: $0 \rightarrow C_1 \rightarrow C_0 \rightarrow 0$. Let o be a vertex fixed by γ . A convenient basis for B_0 is $\{\delta_x - \delta_o\}$ where

x runs over all vertices $\neq o$.

Define the section α of ∂ to be $\alpha(\delta_x - \delta_o) = \text{sum of all the edges connecting } o \text{ to } x$.

Note that α commutes with the action of γ so that it is T_γ -equivariant, and also that $\alpha(\delta_x - \delta_o) \in C_q^{in(r)}$ if $x \in X^r$.

The point of the following lemma is to show that in general we can always find a section with the well behaved properties α has in this example.

Lemma 7.7. (*Key Lemma*)

semisimple rank 1 case: *Suppose G has semisimple rank 1. Suppose $\gamma \in G$ is such that $o \in X^\gamma$. For each integer $r \geq 0$ and for all $e \geq e_0$ the map $\partial : C_1 \rightarrow B_0$ has a T_γ -equivariant section α , such that $\alpha(B_0^{in(r)}) \subset C_1^{in(r)}$.*

general case: *Let G be a connected reductive group. Suppose $\gamma \in G$ is such that $o \in X^\gamma$. For each integer $r \geq 0$ there exists an integer e_r so that for all $e \geq e_r$ each map $\partial : C_q \rightarrow B_{q-1}$, $d \geq q \geq 1$, has a T_γ -equivariant section α , such that $\alpha(B_{q-1}^{in(r)}) \subset C_q^{in(r)}$.*

Remark 7.8. The main difference between the general version of the key lemma and the semisimple rank 1 version is the dependence of the parameter e on the radius of truncation r . This dependence is what makes the general version (significantly?) weaker and prevents us from achieving steps (1) and (2) in general.

Proof. $C_q^{in(r)} \subset C_q$ hence $\partial C_q^{in(r)} \subset \partial C_q = B_{q-1}$. Since $C_q^{in(r)}$ consists of maps supported on X^r their boundary is also supported on X^r , so $\partial C_q^{in(r)} \subset C_{q-1}^{in(r)}$.

Put together we see that $\partial C_q^{in(r)} \subset B_{q-1} \cap C_{q-1}^{in(r)} =: B_{q-1}^{in(r)}$. So the map $\partial : C_q \rightarrow B_{q-1}$ restricts to a map $\partial' : C_q^{in(r)} \rightarrow B_{q-1}^{in(r)}$, which is T_γ -equivariant. Recall that $C_q^{in(r)} = C_q(X^r)$.

Claim 7.9. (*semisimple rank 1 case*) *Suppose G has semisimple rank 1. For each $r \geq 0$ and for each $e \geq e_0$, the map ∂' is surjective.*

Proof. Recall the list of facts (section 2.6) regarding the groups $U_F^{(e)}$.

Since in this case the chain complex (\mathcal{C}) is: $0 \rightarrow C_1 \xrightarrow{\partial} C_0 \rightarrow 0$, we only have to show that $\partial' : C_1^{in(r)} \rightarrow B_0^{in(r)}$ is surjective. That is, given a non-vanishing 0-cycle $\omega \in B_0^{in(r)} = C_c^{or}(X_{(0)}^r; \gamma_e(V)) \cap B_0(X)$, we need to show that it is a boundary of a chain in $C_c^{or}(X_{(1)}^r; \gamma_e(V))$. The proof is by induction on the radius of the support of ω . Let $B(o, r_\omega)$ be the smallest ball containing the support of ω , we will show that by adding a boundary to ω , we obtain a 0-cycle ω^* with support contained in a smaller ball $B(o, r_\omega - 1)$. Repeating this process will show that ω is a boundary.

Definition 7.10. Given a finite collection of points in the building, $S \subset X$, we say that a point $x \in S$ is an *extreme point* (relative to a point o), if it is farthest away from o . That is if $d(o, x) \geq d(o, y)$ for all $y \in S$.

The support of ω is finite so it makes sense to talk about an extreme vertex in this support. Let x be such a vertex. Let y be another vertex in the support of ω and let $v = \omega(x) \in V^{U_x^{(e)}}$ and $w = \omega(y) \in V^{U_y^{(e)}}$. Consider the average of w , $\pi(U_x^{(e)})w$, over the compact group $U_x^{(e)}$.

Let F be the facet of the geodesic $geod(y, x)$ whose closure \overline{F} contains x . We next show that $\pi(U_x^{(e)})w \in V^{U_F^{(e)}}$. For any $u_z \in U_z^{(e)}$, where $z \in F$, we have:

$$\begin{aligned} \pi(u_z)\pi(U_x^{(e)})w &= \pi(u_z)\pi(U_x^{(e)})\pi(u_z^{-1})\pi(u_z)w \\ &= \pi(u_z U_x^{(e)} u_z^{-1})\pi(u_z)w \\ &= \pi(U_x^{(e)})\pi(u_z)w \\ &= \pi(U_x^{(e)})\pi(u_x)\pi(u_y)w \\ &= \pi(U_x^{(e)})w. \end{aligned}$$

The first and second equalities are clear. By fact (U5) (section 2.6) we see that u_z normalizes $U_x^{(e)}$. This explains the third equation. Using (U7) we see that u_z is of the form $u_x u_y$ for some $u_x \in U_x^{(e)}$ and some $u_y \in U_y^{(e)}$, which gives the fourth equality. By absorbing u_x into $U_x^{(e)}$ and since u_y acts trivially on w we obtain the fifth equality. Thus $\pi(U_x^{(e)})w \in V^{U_F^{(e)}}$.




FIGURE 4. Geodesics to x

Now label the vertices in the support of ω which are distinct from x by y_1, \dots, y_n , (see Figure 4), and let $w_i = \omega(y_i)$. Let F_i denote the facet of

the geodesic $geod(y_i, x)$ whose closure $\overline{F_i}$ contains x , oriented as to point towards x . Since ω is a 0-cycle, we have:

$$\epsilon(\omega) = v + w_1 + \cdots + w_n = 0,$$

where ϵ is the augmentation map of the complex $(*)$.

Applying $\pi(U_x^{(e)})$ to each term above we obtain:

$$v + \pi(U_x^{(e)})(w_1) + \cdots + \pi(U_x^{(e)})(w_n) = 0.$$

Let $\overline{w}_i := \pi(U_x^{(e)})(w_i)$. Since $\pi(U_x^{(e)})(w_i) \in V_{F_i}^{U_x^{(e)}}$, the 1-chain: $\omega_x := \delta_{F_1}^{\overline{w}_1} + \cdots + \delta_{F_n}^{\overline{w}_n}$ is in $C_1^{in(r)}$ (where δ_F^v is a ‘delta chain’ supported on F with value v). Now, $\omega = (\omega + \partial\omega_x) - \partial\omega_x$, where $\omega + \partial\omega_x \in B_0^{in(r)}$. Note that $(\omega + \partial\omega_x)(x) = 0$, so that $\omega + \partial\omega_x$ is not supported on x . Note also that as a consequence of this process, we introduced into the support of $\omega + \partial\omega_x$ the (possibly) new vertices x_i , where $\partial\delta_{F_i}^{\overline{w}_1} = \delta_x^{\overline{w}_1} - \delta_{x_i}^{\overline{w}_1}$. We next show that the vertices x_i are strictly closer to o than x and so $\omega + \partial\omega_x$ is an improvement on ω in the sense of trying to show it is a boundary. Now repeating the process above for all extreme vertices of distance $d(o, x)$ we obtain the 0-cycle ω^* which is supported on $B(o, r_\omega - 1)$. Thus by induction we have shown that ω is a boundary of a 1-chain in $C_1^{in(r)}$ and the claim is proved.

Remark 7.11. In the proof above, since the building is 1-dimensional, all the facets F_i are actually the same facet. Since parts of this proof generalize to higher dimensional buildings we leave it intact.

In order to show that this process terminates we need to show that the vertices x_i are strictly closer to o than x .

We use the *negative curvature inequality*, [Bro, p.153], which says that for all $t \in [0, 1]$:

$$d^2(o, p_t) \leq (1-t)d^2(o, y) + td^2(o, x) - t(t-1)d^2(y, x)$$

where $p_t := (1-t)y + tx \in geod(y, x)$, $t \in [0, 1]$. For $t \in (0, 1)$ we have:

$$d^2(o, p_t) \leq (1-t)d^2(o, y) + td^2(o, x)$$

as $t(t-1)d^2(y, x) \geq 0$. Now $d(o, y) \leq d(o, x) = r_\omega$, so:

$$d^2(o, p_t) \leq (1-t)r_\omega^2 + tr_\omega^2 = r_\omega^2$$

hence:

$$d(o, p_t) \leq r_\omega.$$

Since $x_i = p_t$ for some $t \in (0, 1)$ we see that the vertices x_i are indeed closer to o than is x :

$$d(o, x_i) \leq d(o, x).$$

□

Claim 7.12. (general case) Let G be a connected reductive group. For each $r \geq 0$ there exists an integer e_r , depending on r , such that for all $e \geq e_r$ the map $\partial' : C_q^{in(r)} \rightarrow B_{q-1}^{in(r)}$ is surjective.

Proof. We have the following commutative diagram:

$$\begin{array}{ccc} C_q(X) & \xrightarrow{\partial} & B_{q-1}(X) \\ \cup & & \cup \\ C_q(X^r) & \xrightarrow{\partial'} & B_{q-1}^{in(r)} \quad := \quad B_{q-1}(X) \cap C_{q-1}(X^r) \end{array}$$

The map $\partial : C_q(X) \rightarrow B_{q-1}(X)$ is surjective by definition. Using $B_{q-1}(X^r) \subset B_{q-1}^{in(r)} \subset B_{q-1}(X)$ and the exactness of $(\bar{*})$ we obtain:

$$B_{q-1}(X^r) = Z_{q-1}(X^r) = B_{q-1}(X) \cap Z_{q-1}(X^r) = B_{q-1}(X) \cap C_{q-1}(X^r) =: B_{q-1}^{in(r)}$$

Thus the map $\partial' : C_q^{in(r)} \rightarrow B_{q-1}^{in(r)}$ is surjective and the claim is proved. \square

Now, continuing with the proof of the key lemma, let $\alpha' : B_{q-1}^{in(r)} \rightarrow C_q^{in(r)}$ be a T_γ -equivariant section of ∂' and let $\alpha'' : B_{q-1}^{out(r)} \rightarrow C_q(X)$ be any section of $\partial : C_q(X) \rightarrow B_{q-1}(X)$. Again use K_γ to average α'' and make it T_γ -equivariant. Let $\alpha : B_{q-1}^{in(r)}(X) \oplus B_{q-1}^{out(r)}(X) \rightarrow C_q(X)$ be $\alpha' \oplus \alpha''$. α satisfies the properties in the statement of the lemma. \square

Corollary 7.13. *Recall Definition 7.1 of nice \oplus decompositions.*

semisimple rank 1 case: *For G of semisimple rank 1, nice \oplus decompositions exist and are uniform in e .*

general case: *For G a connected reductive group, nice \oplus decompositions exist but they depend on e_r .*

Proof. Let H_0 be any T_γ -equivariant complement of B_0 inside C_0 . Recall the T_γ -equivariant decomposition $B_{q-1} = B_{q-1}^{in(r)} \oplus B_{q-1}^{out(r)}$. Use the T_γ -equivariant section α of Lemma 7.7 to define:

$$\begin{aligned} (B_{q-1}^{in(r)})' &:= \alpha(B_{q-1}^{in(r)}) \\ (B_{q-1}^{out(r)})' &:= \alpha(B_{q-1}^{out(r)}). \end{aligned}$$

The properties of α guarantee that this construction gives a nice \oplus decomposition. \square

7.3. Traces with respect to nice decompositions. Given direct sum decompositions of the vector spaces C_q ($d \geq q \geq 0$):

$$(\oplus) \quad C_q = B_q \oplus H'_q \oplus B'_{q-1}$$

we now show that if such decompositions are nice (with respect to γ and r) then the finite rank operators $T_\gamma Q_q^r$ and $T_\gamma \overline{Q}_q^r$ all have the same trace.

Proposition 7.14. *Let $\gamma \in G$ be an element such that $o \in X^\gamma$. If the decomposition (\oplus) is nice relative to γ and r , then*

$$\text{trace}(T_\gamma Q_q^r; C_q) = \text{trace}(T_\gamma \overline{Q}_q^r; C_q) \quad d \geq q \geq 0$$

Proof. For $q = 0$ have $C_0 = B_0 \oplus H'_0$. Since B_0 and H'_0 are T_γ -stable, the linear operator T_γ on $C_0 = B_0 \oplus H'_0$ is block diagonal:

$$T_\gamma = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$$

As \overline{Q}_0^r differs from Q_0^r only on the bottom left block (with respect to $C_0 = B_0 \oplus H'_0$):

$$Q_0^r = \begin{pmatrix} Q_0^r[B_0; B_0] & Q_0^r[B_0; H'_0] \\ Q_0^r[H'_0; B_0] & Q_0^r[H'_0; H'_0] \end{pmatrix} \quad \overline{Q}_0^r = \begin{pmatrix} Q_0^r[B_0; B_0] & Q_0^r[B_0; H'_0] \\ 0 & Q_0^r[H'_0; H'_0] \end{pmatrix},$$

they agree on the diagonal blocks and so the diagonal blocks of $T_\gamma Q_0^r$ and those of $T_\gamma \overline{Q}_0^r$ are the same. Hence these finite rank operators have the same trace.

For $d \geq q \geq 1$ the complex (\mathcal{C}) is exact at the q^{th} place, hence $C_q = B_q \oplus B'_{q-1}$ where $B'_{q-1} \xrightarrow[\sim]{\partial} B_{q-1}$ and \oplus is T_γ -stable.

Since for $d \geq q \geq 1$, Q_q^r and \overline{Q}_q^r differ also on the bottom (right) diagonal block:

$$Q_q^r = \begin{pmatrix} Q_q^r[B_q; B_q] & Q_q^r[B_q; B'_{q-1}] \\ Q_q^r[B'_{q-1}; B_q] & Q_q^r[B'_{q-1}; B'_{q-1}] \end{pmatrix} \quad \overline{Q}_q^r = \begin{pmatrix} Q_q^r[B_q; B_q] & Q_q^r[B_q; B'_{q-1}] \\ 0 & \widetilde{Q}_{q-1}^r \end{pmatrix}$$

we cannot conclude yet that $T_\gamma Q_q^r$ and $T_\gamma \overline{Q}_q^r$ have the same diagonal blocks with respect to $C_q = B_q \oplus B'_{q-1}$. We now use the (refined) nice \oplus decomposition of C_q . Recall that $(B_{q-1}^{\text{in}(r)})' = \alpha(B_{q-1}^{\text{in}(r)})$ and that $(B_{q-1}^{\text{out}(r)})' = \alpha(B_{q-1}^{\text{out}(r)})$, where α is the section of ∂ of Lemma 7.7.

With respect to the decomposition $C_q = B_q \oplus (B_{q-1}^{\text{in}(r)})' \oplus (B_{q-1}^{\text{out}(r)})'$ we have:

$$T_\gamma = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix}.$$

Since Q_q^r and \overline{Q}_q^r agree on B_q we show that they have the same diagonal blocks with respect to $(B_{q-1}^{\text{in}(r)})' \oplus (B_{q-1}^{\text{out}(r)})'$.

Since $\overline{Q}_q^r[B'_{q-1}; B'_{q-1}] = \widetilde{Q}_{q-1}^r = Q_{q-1}^r[B_{q-1}; B_{q-1}]$ we look at Q_{q-1}^r on $B_{q-1} = B_{q-1}^{\text{in}(r)} \oplus B_{q-1}^{\text{out}(r)}$.

Note that:

$$\begin{aligned} B_{q-1}^{\text{out}(r)} \cap C_{q-1}^{\text{in}(r)} &= (B_{q-1}^{\text{out}(r)} \cap B_{q-1}) \cap C_{q-1}^{\text{in}(r)} \\ &= B_{q-1}^{\text{out}(r)} \cap (B_{q-1} \cap C_{q-1}^{\text{in}(r)}) \\ &= B_{q-1}^{\text{out}(r)} \cap B_{q-1}^{\text{in}(r)} = \{0\}. \end{aligned}$$

Now for $b \in B_{q-1}^{out(r)}$, have $Q_{q-1}^r[B_{q-1}^{out(r)}; B_{q-1}^{out(r)}](b) \in B_{q-1}^{out(r)} \cap C_{q-1}^{in(r)} = \{0\}$, hence Q_{q-1}^r acts as the zero operator on $B_{q-1}^{out(r)}$.

On $B_{q-1}^{in(r)}$ we have Q_{q-1}^r acting as the identity operator since $B_{q-1}^{in(r)} \subset C_{q-1}^{in(r)}$. So \overline{Q}_q^r with respect to $C_q = B'_q \oplus (B_{q-1}^{in(r)})' \oplus (B_{q-1}^{out(r)})'$ has the form:

$$\overline{Q}_q^r = \begin{pmatrix} Q_q^r & * & * \\ 0 & 1 & * \\ 0 & 0 & 0 \end{pmatrix}.$$

We now find the form of Q_q^r with respect to $C_q = B'_q \oplus (B_{q-1}^{in(r)})' \oplus (B_{q-1}^{out(r)})'$.

Let $b \in (B_{q-1}^{out(r)})'$, then $\partial b \in B_{q-1}^{out(r)}$. $Q_q^r b = b_q + b^{in} + b^{out} \in C_q^{in(r)}$ so $\partial Q_q^r b = 0 + \partial b^{in} + \partial b^{out} \in B_{q-1}^{in(r)}$. Hence $\partial b^{out} = 0$ and so $b^{out} = 0$ which means that $Q_q^r[(B_{q-1}^{out(r)})'; (B_{q-1}^{out(r)})'] = 0$. Now, $(B_{q-1}^{in(r)})' \subset C_{q-1}^{in(r)}$ so $Q_q^r = 1$ on $(B_{q-1}^{in(r)})'$. Hence we obtain that Q_q^r has the form:

$$Q_q^r = \begin{pmatrix} Q_q^r & * & * \\ * & 1 & * \\ * & 0 & 0 \end{pmatrix}$$

Thus Q_q^r and \overline{Q}_q^r have the same diagonal blocks and so $trace(T_\gamma Q_q^r) = trace(T_\gamma \overline{Q}_q^r)$. \square

8. RECOVERING SCHNEIDER-STUHLER'S RESULT FOR SEMISIMPLE RANK 1 GROUPS

The following lemma is part of the proof of lemma 12 in [S-S III.4]. It holds for any connected reductive group G with building X . For the sake of completeness we repeat the proof here.

Lemma 8.1. *Let $\gamma \in G^{ell}$. There exists an open subgroup $U \subset G$ such that*

$$X^\gamma = X^{\gamma'} \text{ for all } \gamma' \in \gamma U.$$

Proof. Since γ is elliptic, its fixed point set is non-empty and compact. Fix a point $o \in X^\gamma$ and choose $r \geq 0$ large enough so that X^γ is contained in the interior of the closed ball $B(o, r)$. By construction $B(o, r)$ has no γ -fixed points on its boundary. Let U denote the open subgroup: $U := \{g \in G \mid gx = x \text{ for all } x \in B(o, r)\}$. For $\gamma' \in \gamma U$ we show that $X^\gamma = X^{\gamma'}$.

It is clear that the actions of γ and γ' agree on all the points of the closed ball $B(o, r)$, so in particular they have the same fixed points inside $B(o, r)$: $X^\gamma \cap B(o, r) = X^{\gamma'} \cap B(o, r)$. Suppose γ' has a fixed point x outside the ball. Then the whole geodesic $geod(o, x)$ must be fixed by γ' and so there is a γ' -fixed point $x_0 := \partial B(o, r) \cap geod(o, x)$ on the boundary of the ball. But any such γ' -fixed point is also a γ -fixed point, contradicting γ having no fixed points on the boundary of $B(o, r)$. \square

Let (π, V) be an admissible representation of G with character function $\Theta_\pi(g)$, defined on the regular elements in G . Let γ be a regular semisimple elliptic element: $\gamma \in G^{ell}$, with $o \in X^\gamma$, and fix $e \geq e_0$. We recover the character formula (1) of Schneider–Stuhler.

Claim 8.2. *We can fix an open compact subgroup K of G with the following properties:*

- (1) *The character is locally constant on the neighborhood γK of γ , i.e. $\Theta_\pi(\gamma) = \Theta_\pi(\gamma K)$*
- (2) *All elements $\gamma k, k \in K$, have the same fixed point set: $X^\gamma = X^{\gamma k}$*
- (3) *$K \subset \bigcap_{x \in X^\gamma} U_x^{(e)}$*
- (4) *γ normalizes K .*

Proof. To see that such a group K exists, consider the following. If U denotes the open subgroup of the lemma above, and $U_x^{(e)}$ the usual open (compact) subgroups attached to a point $x \in X$, then $U \bigcap_{x \in X^\gamma} U_x^{(e)}$ is an open neighborhood of the identity in G . Since γ fixes o , it is contained in the group P_o^\dagger , and so it normalizes all the open compact groups $U_o^{(e)}$, (property (U2) above). Thus if we let $K := U_o^{(e)}$ and choose e large enough, we can make sure that $\Theta_\pi(\gamma K) = \Theta_\pi(\gamma)$ and $K \subset U \bigcap_{x \in X^\gamma} U_x^{(e)}$. Such K satisfies all the properties above. \square

Set $f := \frac{1_{\gamma K}}{\text{vol}(K)} \in C_c^\infty(G)$ to be the characteristic function of the set γK normalized by its volume. We have:

$$\begin{aligned}
 \text{trace}(\pi(f); V) &= \int_K f(\gamma k) \text{trace}(\pi(\gamma k); V) dk \\
 &= \int_K \frac{1}{\text{vol}(K)} \Theta_\pi(\gamma k) dk \\
 &= \Theta_\pi(\gamma) \int_K \frac{1}{\text{vol}(K)} dk \\
 &= \Theta_\pi(\gamma)
 \end{aligned}$$

Choose $r = r(f, \gamma)$ large enough so that it satisfies the requirement of Theorem 6.1 and so that $X^\gamma \subset B(o, r)$.

For the rest of this section G will denote a group of semisimple rank 1. Recall that for such groups we have shown (Corollary 7.13) that nice \oplus decompositions exist uniformly in e (with respect to $\gamma \in G^{cpt}$ such that $o \in X^\gamma$).

All elements $\gamma k \in \gamma K$ share the same fixed point o . We claim that it is possible to choose a nice decomposition (\oplus) which is common to all such elements: the issue being that we want the decomposition to be $T_{\gamma k}$ -equivariant simultaneously for all $k \in K$.

In the construction of a nice decomposition (\oplus) we averaged over the group K_γ so as to make sure the construction was γ -equivariant. Now, we will average over K_γ and then average again over K . Property (4) of the group K implies that the construction is both γ -equivariant and K -equivariant. We demonstrate this process with the section α . Start with α and average over K_γ to obtain:

$$\alpha' := \int_{K_\gamma} T_k^{-1} \circ \alpha \circ T_k$$

which is γ -equivariant. Now average α' over K the obtain:

$$\alpha'' := \int_K T_k^{-1} \circ \alpha' \circ T_k$$

By construction, α'' is K -equivariant. We check that it is still γ -equivariant:

$$\begin{aligned} T_\gamma^{-1} \circ \alpha'' \circ T_\gamma &= \int_K T_\gamma^{-1} \circ T_k^{-1} \circ \alpha' \circ T_k \circ T_\gamma \\ &= \int_K (T_\gamma^{-1} \circ T_k^{-1} \circ T_\gamma) \circ (T_\gamma^{-1} \circ \alpha' \circ T_\gamma) \circ (T_\gamma^{-1} \circ T_k \circ T_\gamma) \\ &= \int_K T_{k'}^{-1} \circ \alpha' \circ T_{k'} = \alpha'' \end{aligned}$$

Thus the nice decomposition works uniformly on γK and so by Proposition 7.14:

$$\text{trace}(T_{\gamma k} Q_q^r; C_q) = \text{trace}(T_{\gamma k} \overline{Q}_q^r; C_q) \quad d \geq q \geq 0 \quad \text{for all } k \in K.$$

For γ regular elliptic $\text{trace}(T_\gamma; C_q)$ is essentially counting stable q -facets (with multiplicity) and hence we see that:

$$\text{trace}(T_\gamma; C_q) = \sum_{\substack{\gamma\text{-stable} \\ F \in X_q}} (-1)^{q - \dim F(\gamma)} \text{trace}(\gamma; V_F^{U_F^{(e)}}).$$

Here the quantity $q - \dim F(\gamma)$ is ± 1 , depending on whether γ preserves or reverses the orientation of F (see [SS1, p.45 and p.51] for details).

In the notation of [SS2] we have:

$$\begin{aligned} \sum_{q=0}^d (-1)^q \text{trace}(T_\gamma; C_q) &= \sum_{q=0}^d \sum_{\substack{\gamma\text{-stable} \\ F \in X_q}} (-1)^{\dim F(\gamma)} \text{trace}(\gamma; V^{U_F^{(e)}}) \\ &= \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma)_q} (-1)^q \text{trace}(\gamma; V^{U_F^{(e)}}) \end{aligned}$$

Among other things this says that the left-hand-side equation is constant on γK (since the right-hand-side is the same for all γk , $k \in K$, by the choice of K).

Using this equality and Theorem 6.1 (with $f := \frac{1_{\gamma K}}{\text{vol}(K)}$ and the r above) we obtain:

$$\begin{aligned} \sum_{q=0}^d (-1)^q \text{trace}(T_\gamma; C_q) \int_K dk &= \int_K \left\{ \sum_{q=0}^d (-1)^q \text{trace}(T_{\gamma k}; C_q) \right\} dk \\ &= \int_K \left\{ \sum_{q=0}^d (-1)^q \text{trace}(T_{\gamma k} Q_q^r; C_q) \right\} dk \\ &= \int_K \left\{ \sum_{q=0}^d (-1)^q \text{trace}(T_{\gamma k} \bar{Q}_q^r; C_q) \right\} dk \\ &= \sum_{q=0}^d (-1)^q \text{trace}(T_f \bar{Q}_q^r; C_q) \\ &= \text{trace}(\pi(f); V) = \Theta_\pi(\gamma). \end{aligned}$$

Putting all of the above together we recover the Schneider-Stuhler formula (1), for groups of semisimple rank 1:

Theorem 8.3. *For G a semisimple rank 1 group, V a finitely generated admissible representation of G , $\gamma \in G$ regular elliptic and $e \geq e_0(V)$:*

$$\Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma)_q} (-1)^q \text{trace}(\gamma; V^{U_F^{(e)}})$$

Note: unlike the Schneider-Stuhler proof of this theorem, our proof does not rely on Kazhdan's density theorem.

9. A CHARACTER FORMULA FOR COMPACT ELEMENTS

In this section we prove the main results of this paper: Theorem 9.10 and Corollary 9.11. The key fact used in proving the main result, which will be developed here, is that a regular semisimple compact element γ has an open

neighborhood in G such that, as we vary γ' inside such neighborhood, the following expression stays constant:

$$(2) \quad \sum_{q=0}^d \sum_{F(\gamma') \in (X^{\gamma'} \cap X^r)_q} (-1)^q \text{trace}(\gamma'; V^{U_F^{(e)}})$$

9.1. The periodic nature of X^γ . When γ is elliptic, we know that its fixed point set, X^γ , is finite (as a simplicial complex). When γ is compact, the fixed point set, X^γ , can be an infinite simplicial complex. Yet, as we now show, this set has periodic nature. [See Figures 5 and 2]. Recall that the centralizer of γ , $C_G(\gamma)$, acts on X^γ .




FIGURE 5. Schematic description of the periodic nature of X^γ

Let $K = K_o$ denote the stabilizer of the vertex $o \in X^\gamma$. K is an open subgroup of G , which is compact modulo the center of G . In fact it is a maximal such subgroup. Denote its characteristic function by ch_K .

We now assume for simplicity that all of the vertices of the building X have the same type. This means that we can identify the set of cosets G/K with the vertices of the building, X_0 , via $gK \rightarrow go$. (Without this assumption the vertices can have a finite number of types and a little more care is needed in keeping track of the different types.)

Fix a Haar measures dg on G and dt on T . Let $\frac{dg}{dt}$ be the invariant measure on $T \backslash G$ with respect to dg and dt . The orbital integral

$$\mathcal{O}_\gamma(ch_K) := \int_{T \backslash G} ch_K(g^{-1}\gamma g) \frac{dg}{dt}$$

where $T := C_G(\gamma)^\circ$ is known to converge, say to the (finite) number N .

Remark 9.1. An orbital integral such as $\int_G ch_K(g^{-1}\gamma g) dg$ does not converge in general. The issue is that the split part of T , might not be compact and may make this integral diverge. In our case we integrate over $T \backslash G$ so we are avoiding this type of problem. Also note that since T is generally of smaller dimension than G , we need two normalization factors, one for dg and one for dt .

Since a vertex $gK \in X$ is fixed by γ ($\gamma gK = gK$) if and only if $g^{-1}\gamma g \in K$, we have the following:

$$\begin{aligned} N = \int_{T \backslash G} ch_K(g^{-1}\gamma g) \frac{dg}{dt} &= \sum_{g \in T \backslash G/K} \frac{vol_G(K)}{vol_T(T \cap K)} ch_K(g^{-1}\gamma g) \\ &= \sum_{\substack{g \in T \backslash G/K \\ \gamma gK = gK}} \frac{vol_G(K)}{vol_T(T \cap K)} \\ &= \frac{vol_G(K)}{vol_T(T \cap K)} (\text{number of } T\text{-orbits in } (G/K)^\gamma). \end{aligned}$$

Thus the set of γ -fixed vertices, $X_0^\gamma = (G/K)^\gamma$, is the union of a finite number of T -orbits. Similarly, if we allow vertices of different types and consider also higher dimensional facets, we reach the same conclusion: that the fixed point set X^γ is the union of a finite number of T -orbits (Figure 5 shows one such T -orbit). That is, there is a *fundamental domain* D in X , which is a *finite* union of facets, such that any facet in X^γ is of the form tF , for some facet $F \in D$ and some $t \in T$ (see Figure 7). This is what we mean by saying that ‘the set X^γ has periodic nature’.

Remark 9.2. In the case that γ is elliptic, so that T is compact, each T -orbit on X_0^γ contains finitely many points. But in the case that γ is compact but not elliptic, each T -orbit is infinite. (Essentially the maximal k -split torus contained in T , acts as a translation.)

9.2. An open neighborhood of γ whose elements share the same number of fixed points. In the elliptic case, Lemma 8.1 guaranteed the existence of an open neighborhood γU of γ such that $X^\gamma = X^{\gamma'}$ for all $\gamma' \in \gamma U$. We used the existence of such a neighborhood in showing that the elements $\gamma' = \gamma u$, $u \in U$, all had the same (number of) fixed facets inside

any ball $B(o, r)$ of radius $r \geq 0$ about $o \in X^\gamma$. In general, for a compact element it is not possible to find an open neighborhood γU whose elements share the same fixed point set. In this section we show that it is possible to find an open neighborhood γU whose elements all share the same *number* of fixed points inside any such ball $B(o, r)$.

To understand the behavior of elements near γ it is convenient to assume γ to be regular semisimple (in fact until now it was not necessary to assume γ to be semisimple).

For any maximal torus T , let T' denote the set of regular elements of T . Recall Harish-Chandra's *submersion principle*, [H-C, Lemma 20, p.55], which says that the following map is submersive:

$$\begin{aligned} \psi : G/T \times T' &\longrightarrow G \\ (gT, t) &\longrightarrow gtg^{-1} \end{aligned}$$

This implies that the image of this map

$$\psi(G/T \times T') = \{gtg^{-1} \mid g \in G, t \in T'\} = \mathcal{O}_G(T'),$$

is an open subset in G . Given $\gamma \in G^{reg}$, let $T = C_G(\gamma)^\circ$. In this case $\psi(G/T \times T')$ is an open neighborhood of γ in G^{reg} . So in a small neighborhood of γ , the torus T and the conjugacy class $\mathcal{O}_G(\gamma) := \{g\gamma g^{-1} \mid g \in G\}$ are transversal and $\mathcal{O}_G(T')$ fills out an open neighborhood of γ .

This means that to understand an open neighborhood of (the regular semisimple) γ in G , it is enough to understand an open neighborhood of γ inside $T := C_G(\gamma)^\circ$ and an open neighborhood of γ inside the conjugacy class $\mathcal{O}_G(\gamma)$ [See Figures 6 and 8].



We first study the behavior of elements in an open neighborhood of γ in T .

Let D be a fundamental domain for the action of T on X^γ . Fix a point $o \in D$ and choose $r \geq 0$ large enough so that D is contained in the interior B° of the closed ball $B = B(o, r)$. Let U denote the open subgroup: $U := \{g \in G \mid gx = x \text{ for all } x \in B(o, r)\}$, the pointwise stabilizer of $B(o, r)$. We will show that any element $\gamma' \in T^\circ := T \cap \gamma U$ has the same fixed set as γ : $X^\gamma = X^{\gamma'}$.

Any facet $F \in X^\gamma$ belongs to a T -orbit: say $F \in TF_0$ for some $F_0 \in D$; so there exists a $t \in T$ such that $tF = F_0$. Since any element $\gamma' \in T^\circ$ fixes F_0 and commutes with all elements of T , we have:

$$\gamma'F = \gamma't^{-1}tF = t^{-1}\gamma'(tF) = t^{-1}\gamma'F_0 = t^{-1}F_0 = F.$$

We see that γ' fixes all the facets in X^γ ; that is for all $\gamma' \in T^\circ$: $X^\gamma \subset X^{\gamma'}$.

FIGURE 6. Local picture around γ FIGURE 7. A fundamental domain (in bold) D in the fixed point set of the compact non-elliptic element $\gamma \in GL_2$ of Example 2.5.

Example 9.3. We show that a compact element $\gamma \in G = GL_2$ has an open neighborhood T° in $T = C_G(\gamma)^\circ$, such that all elements in T° have the same

fix point set.

Let $G = GL_2$, and take a compact element $\gamma = \begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}$, where u has the form: $u = 1 + \alpha_r \varpi^r + \dots$ ($0 \neq \alpha_r \in \bar{k}$). Recall (Example 2.5) that the points of X fixed by γ are precisely those at a distance less than or equal to r from the basic apartment A .

Consider the following open neighborhood of u : $u + \varpi^{r+1}O = 1 + \alpha_r \varpi^r + \varpi^{r+1}O$. All elements in this neighborhood have the same valuation. Elements in the corresponding open neighborhood of γ in T :

$$T^\circ = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & u' \end{pmatrix} \mid u' \in u + \varpi^{r+1}O \right\}$$

all have the same fixed point set.

We show this is the case in general.

Lemma 9.4. *All elements $\gamma' \in T^\circ := T \cap \gamma U$ have the same fixed point set: $X^{\gamma'} = X^\gamma$.*

Proof. We imitate the proof of Lemma 8.1 which worked in the elliptic case. Using the notation above, let the open ball B° be the interior of the closed ball B . Let $\mathcal{T}^\circ := TB^\circ$ be the T -orbit B° . Since the subsets tB° are open and since $TB^\circ = \bigcup_{t \in T} tB^\circ$, we see that \mathcal{T}° is an open subset of X . We will call \mathcal{T}° , the *open tube* around X^γ , and its closure, $\mathcal{T} := \overline{\mathcal{T}^\circ}$, the *closed tube* around X^γ . Figure 7 shows these sets for the compact non-elliptic element γ (with $r = 1$) of Example 2.5. Note that $B^\circ \supset D$ and hence that $\mathcal{T}^\circ = TB^\circ \supset TD = X^\gamma$.


It is easy to see that $\mathcal{T} = TB$.

Now let $\gamma' \in \gamma U$. Since the actions of γ and γ' agree on the closed ball B , they also agree on the closed tube $\mathcal{T} = TB$:

$$\gamma tb = t\gamma b = t\gamma' b = \gamma' tb \quad t \in T \quad b \in B.$$

Since $X^\gamma \subset \mathcal{T}^\circ$, γ has no fixed points on the boundary, $\partial\mathcal{T}$, of \mathcal{T} . Hence also γ' has no fixed points on $\partial\mathcal{T}$.

Now suppose γ' has a fixed point x outside the closed tube \mathcal{T} . Then the whole geodesic $geod(o, x)$ must be fixed by γ' and so there is a γ' -fixed point $x_0 := \mathcal{T} \cap geod(o, x)$ on the boundary of the closed tube \mathcal{T} , contradicting γ' having no fixed points on the boundary of \mathcal{T} . Thus $X^{\gamma'} \subset \mathcal{T}$, and hence $X^{\gamma'} = X^\gamma$. \square

FIGURE 8. Tubular neighborhood of γ

Elements in the tubular neighborhood $\mathcal{O}_G(T^\circ)$ of γ all look like $g\gamma'g^{-1}$, $\gamma' \in T^\circ, g \in G$. [See Figure 8]. For such elements and $x \in X^\gamma$, we have: $g\gamma'g^{-1}(gx) = g\gamma'x = gx$. That is $X^{g\gamma'g^{-1}} = gX^{\gamma'} = gX^\gamma$. If we choose g in a small enough neighborhood U° of the identity in G (e.g. $g \in U_o^{(e)}$) so that it fixes o , then $X^{g\gamma'g^{-1}}$ looks like a rotated version of X^γ . [See Figure 9]. Even though the fixed point set $X^{g\gamma'g^{-1}}$ is not the same as X^γ , inside any ball $B(o, r)$, $X^{g\gamma'g^{-1}}$ and X^γ have the same number of fixed facets. Since the construction of X^r is g -equivariant, the same is true inside any truncated building X^r .

That is, since elements in a small enough open neighborhood of γ are all of the form $g\gamma'g^{-1}$, $\gamma' \in T^\circ, g \in U^\circ$, we conclude that there exists an open neighborhood of γ in G whose elements all have the same number of fixed points inside any ball $B(o, r)$, hence inside any truncated building X^r about o .

9.3. Traces of elements in T close to γ . We now study the traces (on the fibers over their fixed points) of elements $\gamma' \in T$ close to γ . We already saw that for elements $\gamma' \in T^\circ$: $X^\gamma = X^{\gamma'}$. Now we show that for $\gamma' \in T^\circ$ and $x \in X^{\gamma'} = X^\gamma$, $\text{trace}(\gamma', V_x^{U_x^{(e)}})$ is constant on the T -orbit of x .

Lemma 9.5. *Given $\gamma' \in T^\circ$, $e \geq 0$, and $x \in X^{\gamma'} = X^\gamma$. For any $s \in T$ we have:*

$$\text{trace}(\gamma', V_x^{U_x^{(e)}}) = \text{trace}(\gamma', V_{sx}^{U_{sx}^{(e)}})$$




FIGURE 9. The fixed point sets X^γ and $X^{g\gamma'g^{-1}}$

Proof. Since $U_{sx}^{(e)} = sU_x^{(e)}s^{-1}$ we have the isomorphism:

$$\begin{aligned} \phi : V^{U_x^{(e)}} &\longrightarrow V^{U_{sx}^{(e)}} \\ v &\mapsto sv \end{aligned}$$

and using the fact that T is abelian we get that for any $\gamma' \in T$ the following diagram commutes:

$$\begin{array}{ccc} v & \xrightarrow{\phi} & sv \\ \gamma' \downarrow & & \downarrow \gamma' \\ \gamma'v & \xrightarrow{\phi} & \gamma'sv = s\gamma'v \end{array}$$

Hence the *trace* of γ' on $V^{U_x^{(e)}}$ is the same as its *trace* on $V^{U_{sx}^{(e)}}$. □

Corollary 9.6. *For γ a (regular semisimple) compact element there is an open neighborhood $T^{\circ\circ} \subset T^\circ$ of γ in $T = C_G(\gamma)^\circ$ (which depends on γ and e) such that all elements in this neighborhood have the same fixed point set, X^γ , and they all act the same way on the fibers above each fixed point $x \in X^\gamma$:*

$$\text{trace}(\gamma', V^{U_x^{(e)}}) = \text{trace}(\gamma, V^{U_x^{(e)}}) \quad \text{for all } \gamma' \in T^{\circ\circ}$$

Proof. By the last lemma, for $\gamma' \in T^\circ$, the traces $\text{trace}(\gamma', V^{U_x^{(e)}})$ and $\text{trace}(\gamma, V^{U_x^{(e)}})$, are constant on the T -orbit of x . Hence it is enough to show this equality for representatives of the orbits. Let $x \in D$ and let $\bigcap_{x \in D} U_x^{(e)}$ be an open neighborhood of the identity in G of elements which act trivially

on all the fibers $V^{U_x^{(e)}}$, $x \in D$. Then all elements $\gamma' \in \gamma(\bigcap_{F \in D} U_F^{(e)})$ act on fibers the same way as γ , so that for $\gamma' \in T^{\circ\circ} := \gamma U \cap \gamma(\bigcap_{F \in D} U_F^{(e)})$, we have: $\text{trace}(\gamma', V^{U_x^{(e)}}) = \text{trace}(\gamma, V^{U_x^{(e)}})$. \square

Corollary 9.7. *The expression:*

$$\sum_{q=0}^d \sum_{F(\gamma') \in (X^{\gamma'} \cap X^r)_q} (-1)^q \text{trace}(\gamma'; V^{U_F^{(e)}})$$

is constant for all $\gamma' \in T^{\circ\circ}$.

Proof. By the last corollary $\text{trace}(\gamma'; V^{U_F^{(e)}})$ is constant for all $\gamma' \in T^{\circ\circ}$. Since $X^{\gamma'}$ is constant for all $\gamma' \in T^{\circ\circ}$ by Lemma 9.4, the expression $(X^{\gamma'} \cap X^r)_q$ is also constant for such γ' and so the summation is over the same facets F as γ' varies. Hence the whole expression above is constant for $\gamma' \in T^{\circ\circ}$. \square

9.4. Traces of elements in $\mathcal{O}_G(T^{\circ\circ})$ close to γ . In this subsection we show that the constancy of the expression above (same as expression (2)) on the open neighborhood $T^{\circ\circ}$ of γ inside T extends to a open neighborhood of γ inside G .

Recall that for elements g in the small neighborhood of the identity and $\gamma' \in T^\circ$, we have: $X^{g\gamma'g^{-1}} = gX^{\gamma'} = gX^\gamma$.

Now, let x be a fixed point of $\gamma' \in T^{\circ\circ}$ and let gx the corresponding fixed point of $g\gamma'g^{-1}$ (recall Figure 9). Note that if g fixes the origin o , then we have that $x \in X^r$ if and only if $gx \in X^r$.

Lemma 9.8. *With the notation above, for any $\gamma' \in T^\circ$, any $g \in U^\circ$, and any $e \geq 0$ we have:*

$$\text{trace}(\gamma', V^{U_x^{(e)}}) = \text{trace}(g\gamma'g^{-1}, V^{U_{gx}^{(e)}})$$

Proof. This proof is almost identical to the proof of the previous Lemma. Since $U_{gx}^{(e)} = gU_x^{(e)}g^{-1}$ we have the isomorphism:

$$\begin{array}{ccc} \phi : V^{U_x^{(e)}} & \xrightarrow{\quad} & V^{U_{gx}^{(e)}} \\ v & \mapsto & gv \end{array}$$

for any $\gamma' \in T^\circ$ the following commutative diagram:

$$\begin{array}{ccc} v & \xrightarrow{\phi} & gv \\ \gamma' \downarrow & & \downarrow g\gamma'g^{-1} \\ \gamma'v & \xrightarrow{\phi} & g\gamma'v \end{array}$$

Hence the *trace* of γ' on $V^{U_x^{(e)}}$ is the same as the *trace* of $g\gamma'g^{-1}$ on $V^{U_{gx}^{(e)}}$. \square

Let $g \in U^\circ$ and $\gamma' \in T^{\circ\circ}$. Since γ' and $g\gamma'g^{-1}$ have the same traces on fibers of respective fixed points we see that the local constancy of the expression (2) holds for all such elements $g\gamma'g^{-1}$. Since elements of this form contain an open neighborhood of γ we obtain the following analogue for compact elements of Lemma 8.1:

Corollary 9.9. *For γ a (regular semisimple) compact element there is an open neighborhood $\mathcal{O}_{U^\circ}(T^{\circ\circ})$ of γ in G (which depends on γ and e) such that for all elements γ' in this neighborhood the expression (2) is constant.*

9.5. Main result: the character on a compact element. In this section we extend the Schneider-Stuhler formula to compact elements. We start with a connected reductive group G and a finitely generated admissible representation (π, V) of G , with character function $\Theta_\pi(g)$. We assume that the decomposition (\oplus) is nice (independently of $e \geq e_0$). We have shown the existence of such nice decompositions (independent of $e \geq e_0$) for groups of semisimple rank 1, but in general we don't have the e independence. Everything in this section applies to a general G . It is only in the last corollary, when we remove this assumption, that we will have to restrict ourselves back to groups of semisimple rank 1. Let $\gamma \in G$ be a regular semisimple compact element, with $o \in X^\gamma$, and fix $e \geq e_0$. The approach here is the analogous approach taken in recovering the Schneider-Stuhler result for semisimple rank 1 groups for elliptic elements. [See section 8].

Fix an open compact subgroup K of G with the following properties (these properties are the analogue for compact elements of the properties listed in Claim 8.2):

- (1) The character is locally constant on the neighborhood γK of γ :
 $\Theta_\pi(\gamma) = \Theta_\pi(\gamma K)$
- (2) γK is contained in the open neighborhood U_γ^e of γ described in Corollary 9.9.
- (3) γ normalizes K .

As explained in section 8, it is possible to choose a nice decomposition (\oplus) which is common to all such elements: the decomposition being $T_{\gamma k}$ -equivariant simultaneously for all $\gamma \in K$ (by (3) above). Hence similarly to the above we obtain:

$$\text{trace}(T_{\gamma k} Q_q^r; C_q) = \text{trace}(T_{\gamma k} \overline{Q}_q^r; C_q) \quad d \geq q \geq 0 \quad \text{for all } k \in K.$$

Take $f := \frac{1_{\gamma K}}{\text{vol}(K)} \in C_c^\infty(G)$ to be the characteristic function of the set γK normalized by its volume, and let $r_0 = r_0(f)$ as in Theorem 6.1. For all $r \geq r_0$ we obtain the following:

$$\begin{aligned}
\Theta_\pi(\gamma) &= \text{trace}(\pi(f); V) \\
&= \int_K \sum_{q=0}^d (-1)^q \text{trace}(T_{\gamma k} \overline{Q}_q^r; C_q) dk \\
&= \int_K \sum_{q=0}^d (-1)^q \text{trace}(T_{\gamma k} Q_q^r; C_q) dk \\
&= \int_K \sum_{q=0}^d \sum_{F(\gamma k) \in (X^{\gamma k} \cap X^r)_q} (-1)^q \text{trace}(\gamma k; V^{U_F^{(e)}}) dk \\
&= \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma \cap X^r)_q} (-1)^q \text{trace}(\gamma; V^{U_F^{(e)}}).
\end{aligned}$$

Which gives the main result of this paper:

Theorem 9.10. *Let G be connected reductive, (π, V) a finitely generated admissible representation of G , $e \geq e_0(V)$, and γ regular semisimple compact. Assume the existence of nice decompositions (\oplus) uniformly in e (see Definition 7.1). For all r large enough¹² we can express the character Θ_π of π on the compact element γ using information contained in the truncated fixed point set $X^\gamma \cap X^r$ as follows:*

$$\Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F(\gamma) \in (X^\gamma \cap X^r)_q} (-1)^q \text{trace}(\gamma; V^{U_F^{(e)}})$$

For groups of semisimple rank 1, we know the existence of nice decompositions uniformly in e (Lemma 7.7), so we can remove this assumption in the theorem:

Corollary 9.11. *For G connected reductive of semisimple rank 1, (π, V) a finitely generated admissible representation of G , $e \geq e_0$, γ regular semisimple compact, and r large enough, the following holds:*

$$\Theta_\pi(\gamma) = \sum_{q=0}^d \sum_{F \in (X^\gamma \cap X^r)_q} (-1)^q \text{trace}(\gamma; V^{U_F^{(e)}})$$

¹²Let r' be large enough so that $X^{r'}$ contains a fundamental domain D for X^γ . Let $r_0 = r_0(f)$ as above. The phrase ‘ r large enough’ means $r \geq \max(r', r_0)$.

10. CONCLUDING REMARKS

In Theorem 9.10 we made the assumption about existence of nice decompositions uniformly in e . Existence of nice decompositions is controlled by the exactness of the truncated complex $(\bar{*})$. In section 7 we proved exactness of this complex: in the semisimple rank 1 case, we showed that exactness is independent of e_r , but in general we had only shown exactness with dependence on e_r (see Lemma 7.7). Thus in the semisimple rank 1 case, we are able to remove the assumption and obtain Corollary 9.11. One can hope that it might be possible to show exactness of $(\bar{*})$ independently of e_r and hence remove the assumption from the main theorem.

REFERENCES

- [AB1] M. F. Atiyah; R. Bott, *A Lefschetz fixed point formula for elliptic complexes. I.* Ann. of Math. (2) 86 (1967) 374–407.
- [AB2] M. F. Atiyah; R. Bott, *A Lefschetz fixed point formula for elliptic complexes. II. Applications.* Ann. of Math. (2) 88 (1968) 451–491.
- [Bro] K.S. Brown, *Buildings.* Springer-Verlag 1989
- [Cas] W. Casselman, *Characters and Jacquet modules.* Math. Ann. 230 (1977), no.2, 101–105.
- [H-C] Harish-Chandra, (notes by G. van Dijk) *Harmonic analysis on reductive p -adic groups*, Lecture Notes in Mathematics, vol. 162, Springer, 1970.
- [Kaz] D. Kazhdan, *Cuspidal geometry of p -adic groups.* J.Analyse Math. 47 (1986), 1–36
- [Kot] R. Kottwitz, *Tamagawa numbers.* Ann. Math. 127 (1988) 629–646.
- [Lan] R. Langlands, *Base change for GL_2 .* Annals of Math. Studies, Number 96 (1980), Princeton university press.
- [Moy] A. Moy, *Displacement functions on the Bruhat-Tits building.* The mathematical legacy of *Harish-Chandra* (Baltimore, MD 1998), 483–499, Proc. Sympos. Pure Math. 68.
- [Mur] F. Murnaghan, *Characters of admissible representations of reductive p -adic groups.* Lecture notes for the thematic year (1999) on number theory at the Canadian CRM.
- [Rog] J. Rogawski, *An application of the building to orbital integrals.* Compositop Math. 42 (1981), 417–423.
- [SS1] P. Schneider; U. Stuhler, *Resolutions for smooth representations of the general linear group over a local field.* J. Reine Angew. Math. 436 (1993), 19–32.
- [SS2] P. Schneider; U. Stuhler, *Representation theory and sheaves on the Bruhat-Tits building.* Inst. Hautes tudes Sci. Publ. Math. No. 85 (1997), 97–191.

[Ser] J.-P. Serre, *Trees*. Springer-Verlag 1980

[Tit] J. Tits, *Reductive groups over local fields*. Automorphic Forms, Representation and L -functions. Proc. Symp. Pure Math. 33 Vol.1 (1979), 29–69.

[Vig] M.-F. Vignéras, *Cohomology of Sheaves on the building and R -representations*. Inventiones Math. 127 (1997), 349–373.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR , MICHIGAN
48109

Current address: Department of Mathematics, University of Toronto, Toronto, Ontario,
Canada M5S 3G3

E-mail address: `jkorman@math.toronto.edu`