

CRASH COURSE ON MANIFOLDS

A **manifold** is a (Hausdorff, second countable) topological space M equipped with an equivalence class of **atlases**.

An **atlas** is an open covering $M = \bigcup_i U_i$ and homeomorphisms $\varphi_i: U_i \rightarrow \Omega_i$ where $\Omega_i \subset \mathbb{R}^n$ is open, such that the transition maps $\varphi_j \circ \varphi_i^{-1}: \varphi_i(U_i \cap U_j) \rightarrow \varphi_j(U_i \cap U_j)$ are smooth (that is, are of type C^∞ : all partial derivatives of all orders exist and are continuous).

Two atlases $\{\varphi_i: U_i \rightarrow \Omega_i\}$ and $\{\tilde{\varphi}_j: \tilde{U}_j \rightarrow \tilde{\Omega}_j\}$ are **equivalent** if their union is an atlas, that is, if $\tilde{\varphi}_j \varphi_i^{-1}$ and $\varphi_i \tilde{\varphi}_j^{-1}$ are all smooth.

$\varphi_i: U_i \rightarrow \Omega_i$ is called a **map**.

$\varphi_i^{-1}: \Omega_i \rightarrow U_i$ is called a **parametrization**.

One can write $\varphi_i = (x^1, \dots, x^n)$. $x^j: U_i \rightarrow \mathbb{R}$ are **coordinates**.

Let $X \subset \mathbb{R}^N$ be any subset and $\Omega \subset \mathbb{R}^n$ open. A continuous map $\varphi: X \rightarrow \Omega$ is called **smooth** if every point in X is contained in an open subset $V \subset \mathbb{R}^N$ such that there exists a smooth function $\tilde{\varphi}: V \rightarrow \Omega$ with $\tilde{\varphi}|_{X \cap V} = \varphi$. A continuous map $\psi: \Omega \rightarrow X$ is called **smooth** if it is smooth as a map to \mathbb{R}^N . A **diffeomorphism** $\varphi: X \rightarrow \Omega$ is a homeomorphism such that both φ and φ^{-1} are smooth.

Theorem. Let $M \subset \mathbb{R}^N$ be a subset that is “locally diffeomorphic to \mathbb{R}^n ”: for every point in M there exists a neighborhood $U \subset M$ and there exists an open subset $\Omega \subset \mathbb{R}^n$ and there exists a diffeomorphism $\varphi: U \rightarrow \Omega$. Then M is a manifold with atlas $\{\varphi: U \rightarrow \Omega\}$. (The transition maps will automatically be smooth.) Such an M is called an **embedded submanifold of \mathbb{R}^N** .

Example. $S^2 = \{(x, y, z) \mid x^2 + y^2 + z^2 = 1\}$. For instance, x, y are coordinates on the upper hemisphere.

A continuous function $f: M \rightarrow \mathbb{R}$ is **smooth** if $f \circ \varphi_i^{-1}: \Omega_i \rightarrow \mathbb{R}$ is smooth for all i (as a function of n variables).

$C^\infty(M) := \{ \text{the smooth functions } f: M \rightarrow \mathbb{R} \}$.

A (continuous) curve $\gamma: \mathbb{R} \rightarrow M$ is **smooth** if $\varphi_i \circ \gamma$ is smooth for all i .

We will define the **tangent space** $T_m M =$ “directions along M at the initial point m ”.

A smooth curve $\gamma: \mathbb{R} \rightarrow M$ with $\gamma(0) = m$ defines “differentiation along the curve”, which is the linear functional $C^\infty(M) \rightarrow \mathbb{R}$,

$$D_\gamma: f \mapsto \left. \frac{d}{dt} \right|_{t=0} f(\gamma(t)).$$

We define an equivalence of such curves by $\gamma \sim \tilde{\gamma}$ if $D_\gamma = D_{\tilde{\gamma}}$.

This means that γ and $\tilde{\gamma}$ have the **same direction** at the point $m = \gamma(0) = \tilde{\gamma}(0)$, that is, they are **tangent** to each other at this point.

Geometric definition of the tangent space:

$$T_m M = \{ \text{the equivalence classes of curves in } M \text{ through } m. \}$$

Leibnitz property: $D_\gamma(fg) = (D_\gamma f)g(m) + f(m)(D_\gamma g)$.

Definition. A **derivation at m** is a linear functional $D: C^\infty(M) \rightarrow \mathbb{R}$ that satisfies the Leibnitz property.

Theorem. The derivations at m form a linear vector space: if D_1, D_2 are derivations and $a, b \in \mathbb{R}$ then $aD_1 + bD_2$ is a derivation.

Theorem. For every derivation D there exists a curve γ such that $D = D_\gamma$.

Corollary. $T_m M$ is a linear vector space (identified with the space of derivations at m).

If x^1, \dots, x^n are coordinates near m then $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}$ are a basis of $T_m M$.

Differential of a function: $df|_m \in T_m^* M = (T_m M)^*$ is given by

$$df|_m(v) = vf,$$

the derivative of f in the direction of $v \in T_m M$.

If x^1, \dots, x^n are coordinates then

$$dx^1 \left(\frac{\partial}{\partial x^j} \right) = \frac{\partial x^1}{\partial x^j} = \delta_{1j},$$

so dx^1, \dots, dx^n is the basis of $T_m^* M$ which is dual to the basis $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}$ of $T_m M$.

In coordinates, $df = \sum_i \frac{\partial f}{\partial x^i} dx^i$.

A **differential form of degree 0** is a smooth function.

A **differential form of degree 1**, $\alpha \in \Omega^1(M)$, associates to each $m \in M$ a linear functional $\alpha_m \in T_m^*M$. In coordinates: $\alpha = \sum_i c_i(x) dx^i$. We require that the coefficients $c_i(x)$ be smooth functions of $x = (x^1, \dots, x^n)$.

A **differential form of degree 2**, $\alpha \in \Omega^2(M)$, associates to each $m \in M$ an alternating (i.e., anti-symmetric) bilinear form $\alpha_m: T_m M \times T_m M \rightarrow \mathbb{R}$. In coordinates: $\alpha = \sum_{i,j} c_{ij}(x) dx^i \wedge dx^j$ (where

$$dx^i \wedge dx^j: (u, v) \mapsto \det \begin{bmatrix} u^i & v^i \\ u^j & v^j \end{bmatrix}$$

if $u = \sum u^k \frac{\partial}{\partial x^k}$ and $v = \sum v^k \frac{\partial}{\partial x^k}$).

A **differential form of degree k**:

$$\alpha = \sum_{i_1, \dots, i_k} c_{i_1 \dots i_k}(x) dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

(where $dx^{i_1} \wedge \dots \wedge dx^{i_k}$ is similarly given by a $k \times k$ determinant).

Exterior derivative:

$$d\alpha = \sum_{i_1, \dots, i_k, j} \frac{\partial c_{i_1 \dots i_k}}{\partial x^j} dx^j \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

α is **closed** if $d\alpha = 0$; α is **exact** if there exists β such that $\alpha = d\beta$.

De Rham cohomology: $H_{dR}^k(M) = \{\text{closed } k\text{-forms}\} / \{\text{exact } k\text{-forms}\}$.

An **oriented manifold** is a manifold equipped with an equivalence class of oriented atlases. (Jacobians of transition maps must have positive determinants.)

Integration: Let M be an oriented manifold of dimension n . For an n -form with support in a coordinate neighborhood U_i : write it as $f(x) dx^1 \wedge \dots \wedge dx^n$ where x^1, \dots, x^n are (oriented) coordinates and take the Riemann integral of f on \mathbb{R}^n . For an arbitrary compactly supported form α : choose a **partition of unity** $\rho_i: M \rightarrow \mathbb{R}$, $\sum \rho_i = 1$, $\text{supp } \rho_i \subset U_i$, and define

$$\int_M \alpha = \sum_i \int_M (\rho_i \alpha),$$

noting that $\rho_i \alpha$ is supported in U_i .

Pullback: $f: M \rightarrow N$ induces $f^*: \Omega^k(N) \rightarrow \Omega^k(M)$. This enables us to integrate a k -form over an oriented k -submanifold.

A **manifold with boundary** is defined like a manifold except that the Ω 's are open subsets of the upper half space. Its boundary ∂M is well defined.

Stokes's theorem: $\int_M d\alpha = \int_{\partial M} \alpha$.

$\alpha \in \Omega^k(M)$ is closed iff $\int_N \alpha = 0$ whenever N is the boundary of a compact oriented submanifold-with-boundary of M . If α is exact, $\int_N \alpha = 0$ for every compact oriented submanifold $N \subset M$. (α is exact iff $\int_N \alpha = 0$ for every smooth cycle $N \subset M$.) If the integral of a closed form on N is nonzero, we can think that N “wraps around a hole in M ”.

Theorem: if M is oriented and compact n -manifold then $\alpha \mapsto \int_M \alpha$ induces an isomorphism $H_{dR}^n(M) \rightarrow \mathbb{R}$.

Multiplicative structure: $[\alpha] \cdot [\beta] = [\alpha \wedge \beta]$ is a well defined ring structure on $H_{dR}^*(M)$. $f: M \rightarrow N$ induces a ring homomorphism $f^*: H_{dR}^*(N) \rightarrow H_{dR}^*(M)$.