

The 13 Postulates

Everything you ever wanted to know about the real numbers is summarized as follows. There is a set \mathbb{R} “of real numbers” with two binary operations defined on it, $+$ and \cdot (“addition” and “multiplication”), two different distinct elements 0 and 1 and a subset \mathbb{P} “of positive numbers” so that the following 13 postulates hold:

- P1** Addition is associative: $\forall a, b, c \quad a + (b + c) = (a + b) + c$ (“ \forall ” means “for every”).
- P2** The number 0 is an additive identity: $\forall a \quad a + 0 = 0 + a = a$.
- P3** Additive inverses exist: $\forall a \exists (-a)$ s.t. $a + (-a) = (-a) + a = 0$ (“ \exists ” means “there is” or “there exists”).
- P4** Addition is commutative: $\forall a, b \quad a + b = b + a$.
- P5** Multiplication is associative: $\forall a, b, c \quad a \cdot (b \cdot c) = (a \cdot b) \cdot c$.
- P6** The number 1 is a multiplicative identity: $\forall a \quad a \cdot 1 = 1 \cdot a = a$.
- P7** Multiplicative inverses exist: $\forall a \neq 0 \exists a^{-1}$ s.t. $a \cdot a^{-1} = a^{-1} \cdot a = 1$.
- P8** Multiplication is commutative: $\forall a, b \quad a \cdot b = b \cdot a$.
- P9** The distributive law: $\forall a, b, c \quad a \cdot (b + c) = a \cdot b + a \cdot c$.
- P10** The trichotomy for \mathbb{P} : for every a , exactly one of the following holds: $a = 0$, $a \in \mathbb{P}$ or $(-a) \in \mathbb{P}$.
- P11** Closure under addition: if a and b are in P , then so is $a + b$.
- P12** Closure under multiplication: if a and b are in P , then so is $a \cdot b$.
- P13** The thirteenth postulate is the most subtle and interesting of all. It will await a few weeks.

Here are a few corollaries and extra points:

1. Sums such as $a_1 + a_2 + a_3 + \cdots + a_n$ are well defined.
2. The additive identity is unique. (Also multiplicative).
3. Additive inverses are unique. (Also multiplicative).
4. Subtraction can be defined.
5. $a \cdot b = a \cdot c$ iff (if and only if) $a = 0$ or $b = c$.
6. $a \cdot b = 0$ iff $a = 0$ or $b = 0$.

7. $x^2 - 3x + 2 = 0$ iff $x = 1$ or $x = 2$.
8. $a - b = b - a$ iff $a = b$.
9. A “well behaved” order relation can be defined (i.e., the boolean operations $<$, \leq , $>$ and \geq can be defined and they have some expected properties).
10. The “absolute value” function $a \mapsto |a|$ can be defined and for all numbers a and b we have

$$|a + b| \leq |a| + |b|.$$