

The decomposition of Global Conformal Invariants V.

Spyros Alexakis*

Abstract

This is the fifth in a series of papers where we prove a conjecture of Deser and Schwimmer regarding the algebraic structure of “global conformal invariants”; these are defined to be conformally invariant integrals of geometric scalars. The conjecture asserts that the integrand of any such integral can be expressed as a linear combination of a local conformal invariant, a divergence and of the Chern-Gauss-Bonnet integrand.

The present paper complements [6] in reducing the purely algebraic results that were used in [3, 4] to certain simpler Lemmas, which will be proven in the last paper in this series, [8].

Contents

1	Introduction	2
2	Proof of the technical Lemmas from [6].	8
2.1	Re-statement of the technical Lemmas 4.6–4.9 from [6].	8
2.2	Two more technical Lemmas.	12
2.3	Proof of Lemmas 4.6, 4.8 in [6]: The main part.	27
2.4	The remaining cases of Lemma 2.9.	44
2.5	Proof of Lemmas 2.2, 2.4:	71
3	The proof of Proposition 1.1 in the special cases:	75
3.1	The direct proof of Proposition 1.1 (in case II) in the “special cases”.	75
3.2	The remaining cases of Proposition 1.1 in case III:	83

*University of Toronto, alexakis@math.toronto.edu.

This work has absorbed the best part of the author’s energy over many years. This research was partially conducted during the period the author served as a Clay Research Fellow, an MSRI postdoctoral fellow, a Clay Liftoff fellow and a Procter Fellow.

The author is immensely indebted to Charles Fefferman for devoting twelve long months to the meticulous proof-reading of the present paper. He also wishes to express his gratitude to the Mathematics Department of Princeton University for its support during his work on this project.

1 Introduction

This is the fourth in a series of papers [3, 5, 6, 7, 8] where we prove a conjecture of Deser-Schwimmer [18] regarding the algebraic structure of global conformal invariants. We recall that a global conformal invariant is an integral of a natural scalar-valued function of Riemannian metrics, $\int_{M^n} P(g) dV_g$, with the property that this integral remains invariant under conformal re-scalings of the underlying metric.¹ More precisely, $P(g)$ is assumed to be a linear combination, $P(g) = \sum_{l \in L} a_l C^l(g)$, where each $C^l(g)$ is a complete contraction in the form:

$$\text{contr}^l(\nabla^{(m_1)} R \otimes \dots \otimes \nabla^{(m_s)} R); \quad (1.1)$$

here each factor $\nabla^{(m)} R$ stands for the m^{th} iterated covariant derivative of the curvature tensor R . ∇ is the Levi-Civita connection of the metric g and R is the curvature associated to this connection. The contractions are taken with respect to the quadratic form g^{ij} . In this series of papers we prove:

Theorem 1.1 *Assume that $P(g) = \sum_{l \in L} a_l C^l(g)$, where each $C^l(g)$ is a complete contraction in the form (1.1), with weight $-n$. Assume that for every closed Riemannian manifold (M^n, g) and every $\phi \in C^\infty(M^n)$:*

$$\int_{M^n} P(e^{2\phi} g) dV_{e^{2\phi} g} = \int_{M^n} P(g) dV_g.$$

We claim that $P(g)$ can then be expressed in the form:

$$P(g) = W(g) + \text{div}_i T^i(g) + \text{Pfaff}(R_{ijkl}).$$

Here $W(g)$ stands for a local conformal invariant of weight $-n$ (meaning that $W(e^{2\phi} g) = e^{-n\phi} W(g)$ for every $\phi \in C^\infty(M^n)$), $\text{div}_i T^i(g)$ is the divergence of a Riemannian vector field of weight $-n+1$, and $\text{Pfaff}(R_{ijkl})$ is the Pfaffian of the curvature tensor.

Before we discuss the position of the present paper in the series [3]–[8], we digress to describe the relation between the present series of papers with classical and recent work on scalar local invariants in various geometries.

Broad Discussion: The theory of *local* invariants of Riemannian structures (and indeed, of more general geometries, e.g. conformal, projective, or CR) has a long history. As discussed in [3], the original foundations of this field were laid in the work of Hermann Weyl and Élie Cartan, see [28, 17]. The task of writing out local invariants of a given geometry is intimately connected with understanding polynomials in a space of tensors with given symmetries; these polynomials are required to remain invariant under the action of a Lie group on the components of the tensors. In particular, the problem of writing down all local Riemannian invariants reduces to understanding the invariants of the orthogonal group.

¹See the introduction of [3] for a detailed discussion of the Deser-Schwimmer conjecture, and for background on scalar Riemannian invariants.

In more recent times, a major program was laid out by C. Fefferman in [20] aimed at finding all scalar local invariants in CR geometry. This was motivated by the problem of understanding the local invariants which appear in the asymptotic expansion of the Bergman and Szegő kernels of strictly pseudoconvex CR manifolds, in a similar way to which Riemannian invariants appear in the asymptotic expansion of the heat kernel; the study of the local invariants in the singularities of these kernels led to important breakthroughs in [11] and more recently by Hirachi in [25]. This program was later extended to conformal geometry in [21]. Both these geometries belong to a broader class of structures, the *parabolic geometries*; these admit a principal bundle with structure group a parabolic subgroup P of a semi-simple Lie group G , and a Cartan connection on that principle bundle (see the introduction in [15]). An important question in the study of these structures is the problem of constructing all their local invariants, which can be thought of as the *natural, intrinsic* scalars of these structures.

In the context of conformal geometry, the first (modern) landmark in understanding *local conformal invariants* was the work of Fefferman and Graham in 1985 [21], where they introduced the *ambient metric*. This allows one to construct local conformal invariants of any order in odd dimensions, and up to order $\frac{n}{2}$ in even dimensions. The question is then whether *all* invariants arise via this construction.

The subsequent work of Bailey-Eastwood-Graham [11] proved that this is indeed true in odd dimensions; in even dimensions, they proved that the result holds when the weight (in absolute value) is bounded by the dimension. The ambient metric construction in even dimensions was recently extended by Graham-Hirachi, [24]; this enables them to indentify in a satisfactory way *all* local conformal invariants, even when the weight (in absolute value) exceeds the dimension.

An alternative construction of local conformal invariants can be obtained via the *tractor calculus* introduced by Bailey-Eastwood-Gover in [10]. This construction bears a strong resemblance to the Cartan conformal connection, and to the work of T.Y. Thomas in 1934, [27]. The tractor calculus has proven to be very universal; tractor buncles have been constructed [15] for an entire class of parabolic geometries. The relation between the conformal tractor calculus and the Fefferman-Graham ambient metric has been elucidated in [16].

The present work [3]–[8], while pertaining to the question above (given that it ultimately deals with the algebraic form of local *Riemannian* and *conformal* invariants), nonetheless addresses a different *type* of problem: We here consider Riemannian invariants $P(g)$ for which the *integral* $\int_{M^n} P(g)dV_g$ remains invariant under conformal changes of the underlying metric; we then seek to understand the possible algebraic form of the *integrand* $P(g)$, ultimately proving that it can be de-composed in the way that Deser and Schwimmer asserted. It is thus not surprising that the prior work on the construction and understanding of local *conformal* invariants plays a central role in this endeavor, in the papers [4, 5].

On the other hand, a central element of our proof are the main algebraic Propositions 5.1, 3.1, 3.2 in [3, 4]; these deal *exclusively* with algebraic properties of the *classical* scalar Riemannian invariants.² The “fundamental Proposition 1.1” makes no reference to integration; it is purely a statement concerning *local Riemannian invariants*. Thus, while the author was led to the main algebraic Propositions in [3, 4] out of the strategy that he felt was necessary to solve the Deser-Schwimmer conjecture, they can be thought of as results of an independent interest. The *proof* of these Propositions, presented in the present paper and in [8] is in fact not particularly intuitive. It is the author’s sincere hope that deeper insight (and hopefully a more intuitive proof) will be obtained in the future as to *why* these algebraic Propositions hold.

We now discuss the position of the present paper in this series of papers.

The purpose of the present paper is to complete the part IIA in this series: In [3, 4, 5] we proved that the Deser-Schwimmer conjecture holds, *provided* one can show certain “Main algebraic propositions”, namely 5.2 in [3] and 3.1, 3.2 in [4]. In [6] we claimed a more general Proposition which *implies* Proposition 5.2 in [3] and Propositions 3.1, 3.2 in [4]; this new “fundamental Proposition” 2.1 in [6] is to be proven by an induction of four parameters. In [6] we also reduced the inductive step of Proposition 2.1 to three Lemmas (in particular we distinguished cases I,II,III on Proposition 2.1 by examining the tensor fields appearing in its hypothesis, see (1.7) below; Lemmas 3.1, 3.2, 3.5 in [6] correspond to these three cases). We proved that these three Lemmas *imply* the inductive step of the fundamental Proposition in cases I,II,III respectively, apart from certain *special cases* which were deferred to the present paper. In these special cases we will derive Proposition 2.1 in [6] directly,³ in section 3. Now, in *proving* that the inductive step of Proposition 1.1 follows from Lemmas 3.1, 3.2, 3.5 in [6] we asserted certain *technical Lemmas*, whose proof was deferred to the present paper. These were Lemmas 4.6, 4.8, and 4.7, 4.9 in [6]; also, the proof of Lemma 5.1 in [6] was deferred to the present paper. We prove all these Lemmas from [6] in section 2.

For reference purposes, and for the reader’s convenience, we recall the precise formulation of the “fundamental Proposition” 2.1 in [6], referring the reader to [6] for a definition of many of the terms appearing in the formulation. First however, we will recall (schematically) the “main algebraic Proposition” 5.2 in [3]; this is a *special case* of Proposition 2.1 in [6], and provides a simpler version of it.

A simpler version of Proposition 2.1 in [6]: Given a Riemannian metric g over an n -dimensional manifold M^n and auxiliary C^∞ scalar-valued functions $\Omega_1, \dots, \Omega_p$ defined over M^n , the objects of study are linear combinations of tensor fields $\sum_{l \in L} a_l C_g^{l, i_1 \dots i_\alpha}$, where each $C_g^{l, i_1 \dots i_\alpha}$ is a *partial contraction* with

²These “main algebraic propositions” are discussed in brief below. A *generalization* of these Propositions is the Proposition 1.1 below.

³By this we mean *without* recourse to the Lemmas 3.1, 3.2, 3.5 in [6].

α free indices, in the form:

$$p \text{contr}(\nabla^{(m)} R \otimes \dots \otimes \nabla^{(m_s)} R \otimes \nabla^{(b_1)} \Omega_1 \otimes \dots \otimes \nabla^{(b_m)} \Omega_p); \quad (1.2)$$

here $\nabla^{(m)} R$ stands for the m^{th} covariant derivative of the curvature tensor R ,⁴ and $\nabla^{(b)} \Omega_h$ stands for the b^{th} covariant derivative of the function Ω_h . A *partial contraction* means that we have list of pairs of indices $(a, b), \dots, (c, d)$ in (1.2), which are contracted against each other using the metric g^{ij} . The remaining indices (which are not contracted against another index in (1.2)) are the *free indices* i_1, \dots, i_α .

The ‘‘main algebraic Proposition’’ 5.2 in [3] (roughly) asserts the following: Let $\sum_{l \in L_\mu} a_l C_g^{l, i_1 \dots i_\mu}$ stand for a linear combination of partial contractions in the form (1.2), where each $C_g^{l, i_1 \dots i_\mu}$ has a given number σ_1 of factors and a given number p of factor $\nabla^{(b)} \Omega_h$. Assume also that $\sigma_1 + p \geq 3$, each $b_i \geq 2$,⁵ and that for each contracting pair of indices (a, b) in any given $C_g^{l, i_1 \dots i_\alpha}$, the indices a, b do not belong to the same factor. Assume also the rank $\mu > 0$ is fixed and each partial contraction $C_g^{l, i_1 \dots i_\mu}$, $l \in L_\mu$ has a given *weight* $-n + \mu$.⁶ Let also $\sum_{l \in L_{>\mu}} a_l C_g^{l, i_1 \dots i_{y_l}}$ stand for a (formal) linear combination of partial contractions of weight $-n + y_l$, with all the properties of the terms indexed in L_μ , *except* that now all the partial contractions have a different rank y_l , and each $y_l > \mu$.

The assumption of the ‘‘main algebraic Proposition’’ 5.1 in [3] is a local equation in the form:

$$\sum_{l \in L_\mu} a_l X \text{div}_{i_1} \dots X \text{div}_{i_\mu} C_g^{l, i_1 \dots i_\mu} + \sum_{l \in L_{>\mu}} a_l X \text{div}_{i_1} \dots X \text{div}_{i_{y_l}} C_g^{l, i_1 \dots i_{y_l}} = 0, \quad (1.3)$$

which is assumed to hold *modulo* complete contractions with $\sigma + 1$ factors. Here given a partial contraction $C_g^{l, i_1 \dots i_\alpha}$ in the form (1.2) $X \text{div}_{i_s} [C_g^{l, i_1 \dots i_\alpha}]$ stands for sum of $\sigma - 1$ terms in $\text{div}_{i_s} [C_g^{l, i_1 \dots i_\alpha}]$ where the derivative ∇^{i_s} is *not* allowed to hit the factor to which the free index i_s belongs.⁷

Proposition 5.2 in [3] then asserts that there will exist a linear combination of partial contractions in the form (1.2), $\sum_{h \in H} a_h C_g^{h, i_1 \dots i_{\mu+1}}$ with all the properties of the terms indexed in $L_{>\mu}$, and all with rank $(\mu + 1)$, so that:

$$\sum_{l \in L_1} a_l C_g^{l, (i_1 \dots i_\mu)} + \sum_{h \in H} a_h X \text{div}_{i_{\mu+1}} C_g^{h, (i_1 \dots i_\mu) i_{\mu+1}} = 0; \quad (1.4)$$

⁴In particular it is a tensor of rank $m + 4$; if we write out its free indices it would be in the form $\nabla_{r_1 \dots r_m}^{(m)} R_{ijkl}$.

⁵This means that each function Ω_h is differentiated at least twice.

⁶See [3] for a precise definition of weight.

⁷Recall that given a partial contraction $C_g^{l, i_1 \dots i_\alpha}$ in the form (1.2) with σ factors, $\text{div}_{i_s} C_g^{l, i_1 \dots i_\alpha}$ is a sum of σ partial contractions of rank $\alpha - 1$. The first summand arises by adding a derivative ∇^{i_s} onto the first factor T_1 and then contracting the upper index i_s against the free index i_s ; the second summand arises by adding a derivative ∇^{i_s} onto the second factor T_2 and then contracting the upper index i_s against the free index i_s etc.

the above holds modulo terms of length $\sigma + 1$. The symbol (\dots) means that we are *symmetrizing* over the indices between parentheses.

In [6] we set up a multiple induction by which we will prove Proposition 5.2 in [3] (outlined above) and also the main algebraic Propositions 3.1, 3.2 in [4]. The generalized proposition 1.1 which we formulated in [6] deals with tensor fields in the forms:

$$\begin{aligned} & p\text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \dots \otimes \nabla^{(m_s)} R_{ijkl} \otimes \\ & \nabla^{(b_1)} \Omega_1 \otimes \dots \otimes \nabla^{(b_p)} \Omega_p \otimes \nabla \phi_1 \otimes \dots \otimes \nabla \phi_u), \end{aligned} \quad (1.5)$$

$$\begin{aligned} & p\text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \dots \otimes \nabla^{(m_{\sigma_1})} R_{ijkl} \otimes \\ & S_* \nabla^{(\nu_1)} R_{ijkl} \otimes \dots \otimes S_* \nabla^{(\nu_t)} R_{ijkl} \otimes \\ & \nabla^{(b_1)} \Omega_1 \otimes \dots \otimes \nabla^{(b_p)} \Omega_p \otimes \\ & \nabla \phi_{z_1} \dots \otimes \nabla \phi_{z_w} \otimes \nabla \phi'_{z_{w+1}} \otimes \dots \otimes \nabla \phi'_{z_{w+d}} \otimes \dots \otimes \nabla \tilde{\phi}_{z_{w+d+1}} \otimes \dots \otimes \nabla \tilde{\phi}_{z_{w+d+y}}). \end{aligned} \quad (1.6)$$

(See the introduction in [6] for a detailed description of the above form). We remark that a complete or partial contraction in the above form will be called “acceptable” if each $b_i \geq 2$, for $1 \leq i \leq p$.⁸ This convention was introduced in [6].

The claim of Proposition is a generalization of the “main algebraic Proposition” in [3]:

Proposition 1.1 *Consider two linear combinations of acceptable tensor fields in the form (1.6):*

$$\begin{aligned} & \sum_{l \in L_\mu} a_l C_g^{l, i_1 \dots i_\mu}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u), \\ & \sum_{l \in L_{>\mu}} a_l C_g^{l, i_1 \dots i_{\beta_l}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u), \end{aligned}$$

where each tensor field above has real length $\sigma \geq 3$ and a given simple character $\vec{\kappa}_{\text{simp}}$. We assume that for each $l \in L_{>\mu}$, $\beta_l \geq \mu + 1$. We also assume that none of the tensor fields of maximal refined double character in L_μ are “forbidden” (see Definition (2.12)).

We denote by

$$\sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$$

a generic linear combination of complete contractions (not necessarily acceptable) in the form (1.5) that are simply subsequent to $\vec{\kappa}_{\text{simp}}$.⁹ We assume that:

⁸In other words, we are requiring each function Ω_i is differentiated at least twice.

⁹Of course if $\text{Def}(\vec{\kappa}_{\text{simp}}) = \emptyset$ then by definition $\sum_{j \in J} \dots = 0$.

$$\begin{aligned}
& \sum_{l \in L_1} a_l X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_\alpha} C_g^{l, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) + \\
& \sum_{l \in L_2} a_l X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_{\beta_l}} C_g^{l, i_1 \dots i_{\beta_l}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) = 0.
\end{aligned} \tag{1.7}$$

We draw our conclusion with a little more notation: We break the index set L_μ into subsets L^z , $z \in Z$, (Z is finite) with the rule that each L^z indexes tensor fields with the same refined double character, and conversely two tensor fields with the same refined double character must be indexed in the same L^z . For each index set L^z , we denote the refined double character in question by \vec{L}^z . Consider the subsets L^z that index the tensor fields of maximal refined double character.¹⁰ We assume that the index set of those z 's is $Z_{Max} \subset Z$.

We claim that for each $z \in Z_{Max}$ there is some linear combination of acceptable $(\mu + 1)$ -tensor fields,

$$\sum_{r \in R^z} a_r C_g^{r, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u),$$

where each $C_g^{r, i_1 \dots i_{\mu+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$ has a μ -double character \vec{L}_1^z and also the same set of factors $S_* \nabla^{(\nu)} R_{ijkl}$ as in \vec{L}^z contain special free indices, so that:

$$\begin{aligned}
& \sum_{l \in L^z} a_l C_g^{l, i_1 \dots i_\mu}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} v \dots \nabla_{i_\mu} v - \\
& \sum_{r \in R^z} a_r X \operatorname{div}_{i_{\mu+1}} C_g^{r, i_1 \dots i_{\mu+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} v \dots \nabla_{i_\mu} v = \\
& \sum_{t \in T_1} a_t C_g^{t, i_1 \dots i_\mu}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} v \dots \nabla_{i_\mu} v,
\end{aligned} \tag{1.8}$$

modulo complete contractions of length $\geq \sigma + u + \mu + 1$. Here each

$$C_g^{t, i_1 \dots i_\mu}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$$

is acceptable and is either simply or doubly subsequent to \vec{L}^z .¹¹

(See the first section in [6] for a description of the notions of *real length*, *acceptable tensor fields*, *simple character*, *refined double character*, *maximal*

¹⁰Note that in any set S of μ -refined double characters with the same simple character there is going to be a subset S' consisting of the maximal refined double characters.

¹¹Recall that "simply subsequent" means that the simple character of $C_g^{t, i_1 \dots i_\mu}$ is subsequent to $\operatorname{Simp}(\vec{L}^z)$.

refined double character, simply subsequent, strongly doubly subsequent). The Proposition 1.1 is proven by a multiple induction on the parameters $-n$ (the weight of the complete contractions appearing in (1.7)), σ (the total number of factors in the form $\nabla^{(m)}R_{ijkl}, S_*\nabla^{(\nu)}R_{ijkl}, \nabla^{(A)}\Omega_h$ among the partial contractions in (1.7)),¹² Φ (the number of factors $\nabla\phi_1, \dots, \nabla\phi_u$ appearing in (1.7)), $\tilde{\Phi}$ (the number of factors $\nabla\phi, \nabla\phi', \nabla\tilde{\phi}$ appearing appearing in (1.7)), and $\sigma_1 + \sigma_2$ (the total number of factors $\nabla^{(m)}R_{ijkl}, S_*\nabla^{(\nu)}R_{ijkl}$). Proposition 1.1 when $\tilde{\Phi} = 0$ coincides with the “Main algebraic Proposition” in [3] outlined above.¹³

2 Proof of the technical Lemmas from [6].

2.1 Re-statement of the technical Lemmas 4.6–4.9 from [6].

We start by recalling a definition from [6] that will be used frequently in the present paper:

Definition 2.1 Consider any tensor field in the form (1.6). We consider any set of indices, $\{x_1, \dots, x_s\}$ belonging to a factor T (here T is not in the form $\nabla\phi$). We assume that these indices are neither free nor are contracting against a factor $\nabla\phi_h$.

If the indices belong to a factor T in the form $\nabla^{(B)}\Omega_1$ then $\{x_1, \dots, x_s\}$ are removable provided $B \geq s + 2$.

Now, we consider indices that belong to a factor $\nabla^{(m)}R_{ijkl}$ (and are neither free nor are contracting against a factor $\nabla\phi_h$). Any such index x which is a derivative index will be removable. Furthermore, if T has at least two free derivative indices, then if neither of the indices i, j are free then we will say one of i, j is removable; accordingly, if neither of k, l is free then we will say that one of k, l is removable. Moreover, if T has one free derivative index then: if none of the indices i, j are free then we will say that one of the indices i, j is removable; on the other hand if one of the indices i, j is also free and none of the indices k, l are free then we will say that one of the indices k, l is removable.

Now, we consider a set of indices $\{x_1, \dots, x_s\}$ that belong to a factor $T = S_*\nabla^{(\nu)}R_{ijkl}$ and are not special, and are not free and are not contracting against any $\nabla\phi$. We will say this set of indices is removable if $s \leq \nu$. Furthermore, if none of the indices k, l are free and $\nu > 0$ and at least one of the other indices in T is free, we will say that one of the indices k, l is removable.

For the first two Lemmas, 2.1, 2.2 we will consider tensor fields in the form:

¹²The partial contractions in (1.7) are assumed to all have the same simple character—this implies that they all have the same number of factors $\nabla^{(m)}R_{ijkl}, S_*\nabla^{(\nu)}R_{ijkl}, \nabla^{(A)}\Omega_h$ respectively.

¹³Similarly, the “Main algebraic Propositions” 3.1, 3.2 in [4] coincide with Proposition 1.1 above when $\tilde{\Phi} = 1$.

$$\begin{aligned}
& p\text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \cdots \otimes \nabla^{m_{\sigma_1}} R_{ijkl} \otimes \\
& S_* \nabla^{(\nu_1)} R_{ijkl} \otimes \cdots \otimes S_* \nabla^{(\nu_t)} R_{ijkl} \otimes \nabla Y \otimes \\
& \nabla^{(b_1)} \Omega_1 \otimes \cdots \otimes \nabla^{(b_p)} \Omega_p \otimes \\
& \nabla \phi_{z_1} \cdots \otimes \nabla \phi_{z_w} \otimes \nabla \phi'_{z_{w+1}} \otimes \cdots \otimes \nabla \phi'_{z_{w+d}} \otimes \cdots \otimes \nabla \tilde{\phi}_{z_{w+d+1}} \otimes \cdots \otimes \nabla \tilde{\phi}_{z_{w+d+y}}).
\end{aligned} \tag{2.1}$$

(Notice this is the same as the form (1.6), but for the fact that we have inserted a factor ∇Y in the second line). Our claims are then the following:

Lemma 2.1 *Assume an equation:*

$$\begin{aligned}
& \sum_{h \in H_2} a_h X_* \text{div}_{i_1} \cdots X_* \text{div}_{i_{a_h}} C_g^{h, i_1 \dots i_{a_h}} (\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}) = \\
& \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_{u'}),
\end{aligned} \tag{2.2}$$

where all tensor fields have rank $a_h \geq \alpha$. All tensor fields have a given u -simple character $\bar{\kappa}'_{\text{simp}}$, for which $\sigma \geq 4$. Moreover, we assume that if we formally treat the factor ∇Y as a factor $\nabla \phi_{u'+1}$ in the above equation, then the inductive assumption of Proposition 1.1 can be applied. (See subsection 3.1 in [6] for a strict discussion of the multi-parameter induction by which we prove Proposition 1.1.)

The conclusion (under various assumptions which we will explain below): Denote by $H_{2, \alpha}$ the index set of tensor fields with rank α .

We claim that there is a linear combination of acceptable¹⁴ tensor fields, $\sum_{d \in D} a_d C_g^{d, i_1 \dots i_{\alpha+1}} (\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$, each with a simple character $\bar{\kappa}'_{\text{simp}}$ so that:

$$\begin{aligned}
& \sum_{h \in H_{2, \alpha}} a_h C_g^{h, i_1 \dots i_{\alpha}} (\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}) \nabla_{i_1} v \cdots \nabla_{i_{\alpha}} v - \\
& X_* \text{div}_{i_{\alpha+1}} \sum_{d \in D} a_d C_g^{d, i_1 \dots i_{\alpha+1}} (\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}) \nabla_{i_1} v \cdots \nabla_{i_{\alpha}} v = \\
& + \sum_{t \in T} a_t C_g^t (\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}, v^{\alpha}).
\end{aligned} \tag{2.3}$$

The linear combination on the right hand side stands for a generic linear combination of complete contractions in the form (2.1) with a factor ∇Y and with a simple character that is subsequent to $\bar{\kappa}'_{\text{simp}}$.

¹⁴“Acceptable” in the sense that each factor Ω_i is differentiated at least twice).

The assumptions under which (2.4) will hold: The assumption under which (2.4) holds is that there should be no tensor fields of rank α in (2.2) which are “bad”. Here “bad” means the following:

If $\sigma_2 = 0$ in $\bar{\kappa}'_{simp}$ then a tensor field in the form (2.1) is “bad” provided:

1. The factor ∇Y contains a free index.
2. If we formally erase the factor ∇Y (which contains a free index), then the resulting tensor field should have no removable indices,¹⁵ and no free indices.¹⁶ Moreover, any factors S_*R_{ijkl} should be simple.

If $\sigma_2 > 0$ in $\bar{\kappa}'_{simp}$ then a tensor field in the form (2.1) is “bad” provided:

1. The factor ∇Y should contain a free index.
2. If we formally erase the factor ∇Y (which contains a free index), then the resulting tensor field should have no removable indices, any factors S_*R_{ijkl} should be simple, any factor $\nabla_{ab}^{(2)}\Omega_h$ should have at most one of the indices a, b free or contracting against a factor $\nabla\phi_s$.
3. Any factor $\nabla^{(m)}R_{ijkl}$ can contain at most one (necessarily special, by virtue of 2.) free index.

Furthermore, we claim that the proof of this Lemma will only rely on the inductive assumption of Proposition 1.1. Moreover, we claim that if all the tensor fields indexed in H_2 (in (2.2)) do not have a free index in ∇Y then we may assume that the tensor fields indexed in D in (2.4) have the same property.

Lemma 2.2 We assume (2.2), where $\sigma = 3$. We also assume that for each of the tensor fields in $H_2^{\alpha,*}$ ¹⁷ there is at least one removable index. We then have two claims:

Firstly, the conclusion of Lemma 2.1 holds in this setting also. Secondly, we can write:

$$\begin{aligned} & \sum_{h \in H_2} a_h X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_\alpha} C_g^{h, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}) = \\ & \sum_{q \in Q} a_q X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_{\alpha'}} C_g^{q, i_1 \dots i_{\alpha'}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}) \\ & + \sum_{t \in T} a_t C_g^t(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'}), \end{aligned} \quad (2.4)$$

¹⁵Thus, the tensor field should consist of factors S_*R_{ijkl} , $\nabla^{(2)}\Omega_h$, and factors $\nabla_{r_1 \dots r_m}^{(m)}R_{ijkl}$ with all the indices r_1, \dots, r_m contracting against factors $\nabla\phi_h$.

¹⁶I.e. $\alpha = 1$ in (2.2).

¹⁷Recall from [6] that $H_2^{\alpha,*}$ is the index set of tensor fields of rank α in (2.2) with a free index in the factor ∇Y .

where the linear combination $\sum_{q \in Q} a_q C_g^{q, i_1 \dots i_{a'}}$ stands for a generic linear combination of tensor fields in the form:

$$\begin{aligned}
& p \text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \dots \otimes \nabla^{(m_{\sigma_1})} R_{ijkl} \otimes \\
& S_* \nabla^{(\nu_1)} R_{ijkl} \otimes \dots \otimes S_* \nabla^{(\nu_t)} R_{ijkl} \otimes \nabla^{(B)} Y \otimes \\
& \nabla^{(b_1)} \Omega_1 \otimes \dots \otimes \nabla^{(b_p)} \Omega_p \otimes \\
& \nabla \phi_{z_1} \dots \otimes \nabla \phi_{z_w} \otimes \nabla \phi'_{z_{w+1}} \otimes \dots \otimes \nabla \phi'_{z_{w+d}} \otimes \dots \otimes \nabla \tilde{\phi}_{z_{w+d+1}} \otimes \dots \otimes \nabla \tilde{\phi}_{z_{w+d+y}}),
\end{aligned} \tag{2.5}$$

with $B \geq 2$, with a simple character $\bar{\kappa}'_{simp}$ and with each $a' \geq \alpha$. The acceptable complete contractions $C_g^t(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_{u'})$ are simply subsequent to $\bar{\kappa}'_{simp}$. $X \text{div}_i$ here means that ∇_i is not allowed to hit the factors $\nabla \phi_h$ (but it is allowed to hit $\nabla^{(B)} Y$).

For our next two Lemmas, we will be considering tensor fields in the general form:

$$\begin{aligned}
& \text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \dots \otimes \nabla^{(m_s)} R_{ijkl} \otimes \\
& S_* \nabla^{(\nu_1)} R_{ijkl} \otimes \dots \otimes S_* \nabla^{(\nu_b)} R_{ijkl} \otimes \nabla_{r_1 \dots r_B}^{(B,+)} (\nabla_a \omega_1 \nabla_b \omega_2 - \nabla_b \omega_1 \nabla_a \omega_1) \\
& \otimes \nabla^{(d_1)} \Omega_p \otimes \dots \otimes \nabla^{(d_p)} \Omega_p \otimes \nabla \phi_1 \otimes \dots \otimes \nabla \phi_u);
\end{aligned} \tag{2.6}$$

here $\nabla_{r_1 \dots r_B}^{(B,+)}(\dots)$ stands for the sublinear combination in $\nabla_{r_1 \dots r_B}^{(B)}(\dots)$ where each ∇ is not allowed to hit the factor $\nabla \omega_2$.

Lemma 2.3 Consider a linear combination of partial contractions,

$$\sum_{x \in X} a_x C_g^{x, i_1 \dots i_a}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_{u'}),$$

where each of the tensor fields $C_g^{x, i_1 \dots i_a}$ is in the form (2.6) with $B = 0$ (and is antisymmetric in the factors $\nabla_a \omega_1, \nabla_b \omega_2$ by definition), with rank $a \geq \alpha$ and real length $\sigma \geq 4$.¹⁸ We assume that all these tensor fields have a given simple character which we denote by $\bar{\kappa}'_{simp}$ (we use u' instead of u to stress that this Lemma holds in generality). We assume an equation:

$$\begin{aligned}
& \sum_{x \in X} a_x X_* \text{div}_{i_1} \dots X_* \text{div}_{i_a} C_g^{x, i_1 \dots i_a}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) = 0,
\end{aligned} \tag{2.7}$$

¹⁸ Recall that in the definition of “real length” in this setting, we count each factor $\nabla^{(m)} R, \nabla^{(\nu)} R, \nabla^{(B)} \Omega_x$ once, the two factors ω_1, ω_2 for one, and the factors $\nabla \phi, \nabla \phi', \nabla \tilde{\phi}$ nor nothing.

where $X_* \text{div}_i$ stands for the sublinear combination in $X \text{div}_i$ where ∇_i is in addition not allowed to hit the factors $\nabla\omega_1, \nabla\omega_2$. The contractions C^j here are simply subsequent to $\vec{\kappa}'_{simp}$. We assume that if we formally treat the factors $\nabla\omega_1, \nabla\omega_2$ as factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ (disregarding whether they are contracting against special indices) in the above, then the inductive assumption of Proposition 1.1 applies.

The conclusion we will draw (under various hypotheses that we will explain below) is that we can write:

$$\begin{aligned} & \sum_{x \in X} a_x X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_a} C_g^{x, i_1 \dots i_a}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) = \\ & \sum_{x \in X'} a_x X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_a} C_g^{x, i_1 \dots i_a}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) + \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) = 0, \end{aligned} \quad (2.8)$$

where the tensor fields indexed in X' on the right hand side are in the form (2.6) with $B > 0$. All the other sublinear combinations are as above. We recall from [6] that $X_+ \text{div}_i$ stands for the sublinear combination in $X \text{div}_i$ where ∇_i is in addition not allowed to hit the factor $\nabla\omega_2$ (it is allowed to hit the factor $\nabla^{(B)}\omega_1$).

Assumptions needed for (2.8): We claim (2.8) under certain assumptions on the α -tensor fields in (2.7) which have rank α and have a free index in one of the factors $\nabla\omega_1, \nabla\omega_2$ (say to $\nabla\omega_1$ wlog)—we denote the index set of those tensor fields by $X^{\alpha,*} \subset X$.

The assumption we need in order for the claim to hold is that no tensor field indexed in $X^{\alpha,*}$ should be “bad”. A tensor field is “bad” if it has the property that when we erase the expression $\nabla_{[a}\omega_1 \nabla_{b]}\omega_2$ (and make the index that contracted against b into a free index) then the resulting tensor field will have no removable indices, and all factors $S_* R_{ijkl}$ will be simple.

Lemma 2.4 *We assume (2.7), where now the tensor fields have length $\sigma = 3$. We also assume that for each of the tensor fields indexed in X , there is a removable index in each of the real factors. We then claim that the conclusion of Lemma 2.3 is still true in this setting.*

For the most part, the remainder of this paper is devoted to proving the above Lemmas. However, we first state and prove some further technical claims, one of which appeared as Lemma 5.1 in [6].¹⁹

2.2 Two more technical Lemmas.

We claim an analogue of Lemma 4.10 in [6] can be derived when we have tensor fields with a given simple character $\vec{\kappa}_{simp}$, and where rather than having one

¹⁹Its proof was also deferred to the present paper.

additional factor $\nabla\phi_{u+1}$ (which is not encoded in the simple character $\vec{\kappa}_{simp}$), we have *two additional factors* $\nabla_a\phi_{u+1}, \nabla_b\phi_{u+2}$.

Lemma 2.5 *Consider a linear combination of acceptable tensor fields in the form (1.6) with a given u -simple character $\vec{\kappa}_{simp}$:*

$\sum_{l \in L} a_l C_g^{l, i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$. Assume that the minimum rank among those tensor fields above is $\alpha \geq 2$. Assume an equation:

$$\begin{aligned} & \sum_{l \in L} a_l X_* \text{div}_{i_3} \dots X_* \text{div}_{i_\beta} C_g^{l, i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} + \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) = 0 \end{aligned} \quad (2.9)$$

(here $X_* \text{div}_i$ means that ∇^i is in addition not allowed to hit the factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$). We also assume that if we formally treat the factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ as factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ then (2.9) falls under the inductive assumption of Proposition 1.1 (with respect to the parameters (n, σ, Φ, u)). Denote by $L^\alpha \subset L$ the index set of terms with rank α . We additionally assume that none of the tensor fields $C_g^{l, i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$ are “forbidden”, in the sense defined above Proposition 2.1 in [6].

We then claim that there exists a linear combination of $(\alpha + 1)$ -tensor fields with a u -simple character $\vec{\kappa}_{simp}$ (indexed in Y below) so that:

$$\begin{aligned} & \sum_{l \in L^\alpha} a_l C_g^{l, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} \nabla_{i_3} v \dots \nabla_{i_\alpha} v \\ & + X_* \text{div}_{i_{\alpha+1}} \sum_{y \in Y} a_y C_g^{y, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} \nabla_{i_3} v \dots \nabla_{i_\alpha} v \\ & + \sum_{j \in J} a_j C_g^{j, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} \nabla_{i_3} v \dots \nabla_{i_\alpha} v. \end{aligned} \quad (2.10)$$

Furthermore, we also claim that we can write:

$$\begin{aligned} & \sum_{l \in L} a_l X \text{div}_{i_3} \dots X \text{div}_{i_\beta} C_g^{l, i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} = \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) + \\ & \sum_{q \in Q_1} a_q X \text{div}_{i_3} \dots X \text{div}_{i_\alpha} C_g^{q, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_{u+2}) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} + \\ & \sum_{q \in Q_2} a_q X \text{div}_{i_3} \dots X \text{div}_{i_\alpha} C_g^{q, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_{u+2}), \end{aligned} \quad (2.11)$$

where the tensor fields indexed in Q_1 are acceptable with a u -simple character $\vec{\kappa}_{simp}$ and with a factor $\nabla^{(2)}\phi_{u+1}$ and a factor $\nabla\phi_{u+2}$. The tensor fields indexed in Q_2 are acceptable with a u -simple character $\vec{\kappa}_{simp}$ and with a factor $\nabla^{(2)}\phi_{u+2}$ and a factor $\nabla\phi_{u+1}$.

Proof of Lemma 2.5: We may divide the index set L^α into subsets $L_I^\alpha, L_{II}^\alpha$ according to whether the two factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ are contracting against the same factor or not—we will then prove our claim for those two index sets separately. Our claim for the index set L_{II}^α follows by a straightforward adaptation of the proof of Lemma 4.10 in [3]. (Notice that the forbidden cases of the present Lemma are exactly in correspondence with the forbidden cases of that Lemma). Therefore, we now prove our claim for the index set L_I^α :

We denote by $L_I \subset L, J_I \subset J$ the index set of terms for which the two factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ are contracting against the same factor. It then follows that (2.9) holds with the index sets L, J replaced by L_I, J_I —denote the resulting new equation by $\text{New}[(2.9)]$. Now, for each tensor field $C_g^{l, i_1 \dots i_\beta}$ and each complete contraction C_g^j , we let $\text{Sym}[C_g^{l, i_1 \dots i_\beta}], \text{Sym}[C_g^{l, i_1 \dots i_\beta}], \text{AntSym}[C_g^j], \text{AntSym}[C_g^j]$ stand for the tensor field/complete contraction that arises from $C_g^{l, i_1 \dots i_\beta}, C_g^j$ by symmetrizing (resp. anti-symmetrizing) the indices a, b in the two factors $\nabla_a\phi_{u+1}, \nabla_b\phi_{u+2}$. We accordingly derive two new equations from $\text{New}[(2.9)]$, which we denote by $\text{New}[(2.9)]_{\text{Sym}}$ and $\text{New}[(2.9)]_{\text{AntSym}}$.

We will then prove the claim separately for the tensor fields in the sublinear combination $\sum_{l \in L_I^\alpha} a_l \text{Sym}[C_g^{l, i_1 \dots i_\alpha}]$ and the tensor fields in the sublinear combination $\sum_{l \in L_I^\alpha} a_l \text{AntSym}[C_g^{l, i_1 \dots i_\alpha}]$.

The claim (2.10) for the sublinear combination $\sum_{l \in L_I^\alpha} a_l \text{AntSym}[C_g^{l, i_1 \dots i_\alpha}]$ follows directly from the arguments in the proof of Lemma 2.3. Therefore it suffices to show our claim for the sublinear combination $\sum_{l \in L_I^\alpha} a_l \text{Sym}[C_g^{l, i_1 \dots i_\alpha}]$.

We prove this claim as follows: We divide the index set L_I^α according to the *form* of the factor against which the two factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ are contracting: List out the non-generic factors in $\vec{\kappa}_{simp}$,²⁰ $\{T_1, \dots, T_a\}$. Then, for each $k \leq a$ we let $L_{I,k}^\alpha$ stand for the index set of terms for which the factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ are contracting against the factor T_k . We also let $L_{I,a+1}^\alpha$ stand for the index set of terms for which the factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ are contracting against a generic factor $\nabla^{(m)}R_{ijkl}$. We will prove our claim for each of the sublinear combinations $\sum_{l \in L_{I,a+1}^\alpha} a_l \text{Sym}[C_g^{l, i_1 \dots i_\alpha}]$ separately.

We firstly observe that for each $k \leq a+1$, we may obtain a new true equation from (2.9) by replacing L by $L_{I,a+1}$ —denote the resulting equation by $(2.9)_{I, \text{Sym}, k}$. Therefore, for each $k \leq a+1$ for which T_k is in the form $\nabla^{(p)}\Omega_h$, our claim follows straightforwardly by applying Corollary 1 from [3].²¹

²⁰Recall from the introduction in [6] that the non-generic factors in $\vec{\kappa}_{simp}$ are all the factors in the form $\nabla^{(A)}\Omega_h, S_*\nabla^{(\nu)}R_{ijkl}$, and also all the factors $\nabla^{(m)}R_{ijkl}$ that contract against at least one factor $\nabla\phi_s$.

²¹There is no danger of falling under a “forbidden case”, since we started with tensor fields which were not forbidden.

Now, we consider the case where the factor T_k is in the form $S_* \nabla^{(\nu)} R_{abcd}$: In that case we denote by $L_{I,k,\sharp}$ the index set of terms for which one of the factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$ is contracting against a special index in T_k . In particular, we will let $L_{I,k,\sharp}^\alpha \subset L_{I,k,\sharp}$ stand for the index set of terms with rank α . We will then show two equations:

Firstly, that there exists a linear combination of tensor fields as claimed in (2.10) so that:

$$\begin{aligned}
& \sum_{l \in L_{I,k,\sharp}^\alpha} a_l \text{Sym}[C]_g^{l,i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v - \\
& \sum_{y \in Y} a_y X \text{div}_{i_{\alpha+1}} C_g^{y,i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v \\
& = \sum_{l \in L_{\mathcal{O}K}^\alpha} a_l X \text{div}_{i_{\alpha+1}} C_g^{l,i_1 \dots i_{\alpha+1}} \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v + \\
& \sum_{j \in J} a_j C_g^{j,i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v,
\end{aligned} \tag{2.12}$$

where the tensor fields in $L_{\mathcal{O}K}^\alpha$ have all the properties of the terms in $L_{I,k}$, rank α and furthermore none of the factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$ are contracting against a special index.

Then (under the assumption that $L_{I,k,\sharp}^\alpha = \emptyset$) we claim that we can write:

$$\begin{aligned}
& \sum_{l \in L_{I,k,\sharp}} a_l X \text{div}_{i_3} \dots X \text{div}_{i_\beta} \text{Sym}[C]_g^{l,i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} \\
& = \sum_{l \in L_{I,k,\mathcal{O}K}} a_l X \text{div}_{i_3} \dots X \text{div}_{i_\beta} \text{Sym}[C]_g^{l,i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \\
& \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} + \sum_{j \in J} a_j \text{Sym}[C]_g^{j,i_1 i_2}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2},
\end{aligned} \tag{2.13}$$

where the tensor fields in $L_{I,k,\mathcal{O}K}$ have all the properties of the terms in $L_{I,k}$, but they additionally have rank $\geq \alpha + 1$ and furthermore none of the factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$ are contracting against a special index.

If we can show the above two equations, then we are reduced to showing our claim under the additional assumption that no tensor field indexed in L in $\text{Sym}(2.9)$ has any factor $\nabla \phi_{u+1}, \nabla \phi_{u+1}$ contracting against a special index in T_k . Under that assumption, we may additionally assume that none of the complete contractions indexed in J in (2.9) have that property.²² Therefore, we may then *erase* the factor $\nabla \phi_{u+1}$ from all the complete contractions and tensor fields in $(2.9)_k$ by virtue of the operation *Erase*, introduced in the Appendix of [3]—our claim then follows by applying Corollary 1 from [3] to the resulting equation and then re-introducing the erased factor $\nabla \phi_{u+1}$.

²²This can be derived by repeating the *proof* of (2.12), (2.13).

Outline of the proof of (2.12), (2.13): Firstly we prove (2.12): Suppose wlog T_k is contracting against $\nabla\phi_1$ and $\nabla\phi'_2, \dots, \nabla\phi'_h$; then replace the two factors $\nabla_a\phi_1, \nabla_b\phi_{u+1}$ by g_{ab} and then apply *Ricto* Ω_{p+1} ,²³ (obtaining a new true equation) and then apply the eraser to the resulting true equation. We then apply Corollary 1 from [3] to the resulting equation,²⁴ and finally we replace the factor $\nabla_{r_1 \dots r_b}^{(b)}\Omega_{p+1}$ by an expression

$$S_* \nabla_{y_2 \dots y_h r_1 \dots r_{b-1}}^{(b+h-1)} R_{ijkr_b} \nabla^i \tilde{\phi}_1 \nabla^j \phi_{u+2} \nabla^k \phi_{u+1} \nabla^{y_2} \phi'_2 \dots \nabla^{y_h} \phi'_h.$$

As in the proof of Lemma 4.10 in [6], we derive our claim. Then, (2.13) is proven by iteratively applying this step and making ∇v 's into $Xdiv$'s at each stage.

We analogously show our claim when the factor T_k is in the form $\nabla^{(m)}R_{ijkl}$: In that case we denote by $L_{I,k,\sharp}$ the index set of terms for which *both* the factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ are contracting against a special index in T_k . We will then show two equations:

Firstly, that there exists a linear combination of tensor fields as claimed in (2.10) so that:

$$\begin{aligned} & \sum_{l \in L_{I,k,\sharp}^\alpha} a_l \text{Sym}[C]_g^{l, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v - \\ & \sum_{y \in Y} a_y Xdiv_{i_{\alpha+1}} C_g^{y, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v = \\ & \sum_{l \in L_{OK}^\alpha} a_l Xdiv_{i_{\alpha+1}} C_g^{l, i_1 \dots i_{\alpha+1}} \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v + \\ & \sum_{j \in J} a_j C_g^{j, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \omega \nabla_{i_3} v \dots \nabla_{i_\alpha} v, \end{aligned} \tag{2.14}$$

where the tensor fields in L_{OK}^α have all the properties of the terms in $L_{I,k}$, but they additionally have rank α and furthermore one of the factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ does not contract against a special index. Then (under the assumption that $L_{I,k,\sharp}^\alpha = \emptyset$) we denote by $L_{I,k,\sharp}$ the sublinear combination of terms in $L_{I,k}$ with *both* factors $\nabla\phi_{u+1}$ or $\nabla\phi_{u+2}$ contracting against a special index in T_k . We claim that we can write:

²³See the relevant Lemma in the Appendix of [3].

²⁴Since the factor $\nabla\phi_{u+2}$ survives this operation, and since we started out with terms that were not “forbidden”, there is no danger of falling under a “forbidden case” of Corollary 1 from [3].

$$\begin{aligned}
& \sum_{l \in L_{I,k,\#}} a_l X \operatorname{div}_{i_3} \dots X \operatorname{div}_{i_\beta} \operatorname{Sym}[C]_g^{l, i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \\
\nabla_{i_2} \phi_{u+2} &= \sum_{l \in L_{I,k,OK}} a_l X \operatorname{div}_{i_3} \dots X \operatorname{div}_{i_\beta} \operatorname{Sym}[C]_g^{l, i_1 \dots i_\beta}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \\
\nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2} &+ \sum_{j \in J} a_j \operatorname{Sym}[C]_g^{j, i_1 i_2}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \nabla_{i_2} \phi_{u+2},
\end{aligned} \tag{2.15}$$

where the tensor fields in $L_{I,k,OK}$ have all the properties of the terms in $L_{I,k}$, but they additionally have rank $\geq \alpha + 1$ and furthermore one of the factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$ does not contract against a special index.

If we can show the above two equations, then we are reduced to showing our claim under the additional assumption that no tensor field indexed in L in $\operatorname{Sym}(2.9)$ has the two factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$, contracting against a special index in T_k . Under that assumption, we may additionally assume that none of the complete contractions indexed in J in (2.9) have that property. Therefore, we may then *erase* the factor $\nabla \phi_{u+1}$ from all the complete contractions and tensor fields in $(2.9)_k$ —our claim then follows by applying Lemma 4.10 in [6] to the resulting equation²⁵ and then re-introducing the erased factor $\nabla \phi_{u+1}$.

Outline of the proof of (2.14), (2.15): Firstly we prove (2.14). Suppose wlog T_k is contracting against $\nabla \phi_1, \dots, \nabla \phi_h$ (possibly with $h = 0$); then replace the two factors $\nabla_a \phi_1, \nabla_b \phi_{u+1}$ by g_{ab} and then apply *Ricto* Ω_{p+1} (obtaining a new true equation) and then apply the eraser to the factors $\nabla \phi_1, \dots, \nabla \phi_h$ in the resulting true equation. Then (apart from the cases, discussed below, where the above operation may lead to a “forbidden case” of Corollary 1 in [6]), we apply Corollary 1 from [6] to the resulting equation, and finally we replace the factor $\nabla_{r_1 \dots r_b}^{(b)} \Omega_{p+1}$ by an expression

$$\nabla_{s_1 \dots s_h r_1 \dots r_{b-2}}^{(b+h)} R_{i r_{b-1} k r_b} \nabla^i \phi_{u+1} \nabla^k \phi_{u+2} \nabla^{s_1} \phi_1 \dots \nabla^{s_h} \phi_h.$$

As in the proof of Lemma 4.10 in [6], we derive our claim. Then, (2.14) is proven by iteratively applying this step and making ∇v 's into $X \operatorname{div}$'s at each stage (again, provided we never encounter “forbidden cases”). If we do encounter forbidden cases, then our claims follow by just making the factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$ into $X \operatorname{div}$'s and then applying Corollary 1 in [3] to the resulting equation (the resulting equation is not forbidden, since it will contain a factor $\nabla^{(m)} R_{ijkl}$ with two free indices), and in the end re-naming two factors ∇v into $\nabla \phi_{u+1}, \nabla \phi_{u+2}$. \square

A Further Generalization: Proof of Lemma 5.1 from [6].

²⁵Notice that there is no danger of falling under a “forbidden case” of that Lemma, since there will be a non-simple factor $S_* \nabla^{(v)} R_{ijkl}$, by virtue of the factor $\nabla \phi_{u+2}$.

We remark that on a few occasions later in this series of papers we will be using a generalized version of the Lemma 2.5. The generalized version asserts that the claim of Lemma 2.5 remains true, for the general case where rather than one or two “additional” factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$ we have $\beta \geq 3$ “additional” factors $\nabla\phi_{u+1}, \dots, \nabla\phi_{u+\beta}$. Moreover, in that case there are no “forbidden cases”.

Lemma 2.6 *Let $\sum_{l \in L_1} a_l C_g^{l, i_1 \dots i_{\mu}, i_{\mu+1} \dots i_{\mu+\beta}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$, $\sum_{l \in L_2} a_l C_g^{l, i_1 \dots i_{b_l}, i_{b_l+1} \dots i_{b_l+\beta}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$ stand for two linear combinations of acceptable tensor fields in the form (1.6), with a u -simple character $\vec{\kappa}_{simp}$. We assume that the terms indexed in L_1 have rank $\mu + \beta$, while the ones indexed in L_2 have rank greater than $\mu + \beta$.*

Assume an equation:

$$\begin{aligned} & \sum_{l \in L_1} a_l X \operatorname{div}_{i_{\beta+1}} \dots X \operatorname{div}_{i_{\mu+\beta}} C_g^{l, i_1 \dots i_{\mu+\beta}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_{\beta}} \phi_{u+\beta} \\ & + \sum_{l \in L_2} a_l X \operatorname{div}_{i_{\beta+1}} \dots X \operatorname{div}_{i_{b_l}} C_g^{l, i_1 \dots i_{b_l+\beta}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_{\beta}} \phi_{u+\beta} \\ & + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_{u+\beta}) = 0, \end{aligned} \tag{2.16}$$

modulo terms of length $\geq \sigma + u + \beta + 1$. Furthermore, we assume that the above equation falls under the inductive assumption of Proposition 2.1 in [6] (with regard to the parameters weight, σ, Φ, p). We are not excluding any “forbidden cases”.

We claim that there exists a linear combination of $(\mu + \beta + 1)$ -tensor fields in the form (1.6) with u -simple character $\vec{\kappa}_{simp}$ and length $\sigma + u$ (indexed in H below) such that:

$$\begin{aligned} & \sum_{l \in L_1} a_l C_g^{l, i_1 \dots i_{\mu+\beta}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_{\beta}} \phi_{u+\beta} \nabla_{i_{\beta+1}} v \dots \nabla_{i_{\beta+\mu}} v \\ & + \sum_{h \in H} a_h X \operatorname{div}_{i_{\mu+\beta+1}} C_g^{h, i_1 \dots i_{\mu+\beta+1}}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_{\beta}} \phi_{\beta+1} \\ & \nabla_{i_1} v \dots \nabla_{i_{\beta+\mu}} v + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_{u+\beta}, v^\mu) = 0, \end{aligned} \tag{2.17}$$

modulo terms of length $\geq \sigma + u + \beta + \mu + 1$. The terms indexed in J here are u -simply subseqent to $\vec{\kappa}_{simp}$.

Proof of Lemma 2.6: The proof of the above is a straightforward adaptation of the proof of Lemma 2.5, *except* for the cases where the tensor fields

$C_g^{l, i_1 \dots i_\mu, i_{\mu+1} \dots i_{\mu+\beta}}$ are “bad”, where “bad” in this case would mean that all factors are in the form R_{ijkl} , $S_* R_{ijkl}$, $\nabla^{(2)} \Omega_h$,²⁶ and in addition each factor $\nabla^{(2)} \Omega_h$ contracts against at most one factor $\nabla \phi_h$, $1 \leq h \leq u + \beta$. So we now focus on that case:

Let us observe that by weight considerations, *all* tensor fields in (2.9) must now have rank μ .

We recall that this special proof applies only in the case where there are special free indices in factors $S_* R_{ijkl}$ among the tensor fields of minimum rank in (2.9). (If there were no such terms, then the regular proof of Lemma 2.5 applies). We distinguish three cases: Either $p > 0$, or $p = 0, \sigma_1 > 0$ or $p = \sigma_1 = 0$ and $\sigma_2 > 0$. We will prove the above by an induction on the parameters (*weight*), σ : Suppose that the weight of the terms in (2.16) is $-K$ and the real length is $\sigma \geq 3$. We assume that the Lemma holds when the equation (2.16) consists of terms with weight $-K'$, $K' < K$, or of terms with weight $-K$ and real length $\sigma', 3 \leq \sigma' < \sigma$.

The case $p > 0$: We first consider the μ -tensor fields in (2.9) with the extra factor $\nabla \phi_{u+1}$ contracting against a factor $\nabla^{(2)} \Omega_h$. Denote the index set of those terms by \bar{L}_μ . We will firstly prove that:

$$\sum_{l \in \bar{L}_\mu} a_l C_g^{l, i_1 \dots i_{\mu+\beta}} (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_\beta} \phi_{u+\beta} \nabla_{i_{\beta+1}} v \dots \nabla_{i_{\beta+\mu}} v = 0. \quad (2.18)$$

It suffices to prove the above for the sublinear combination of μ -tensor fields where $\nabla \phi_{u+1}$ contracts against $\nabla^{(2)} \Omega_1$. (2.18) will then follow by repeating this step p times.

We start by a preparatory claim: Let us denote by $\bar{L}_{\mu, \#} \subset \bar{L}_\mu$ the index set of μ -tensor fields for which the factor $\nabla^{(2)} \Omega_1$ contains a free index, say the index i_1 wlog. We will firstly prove that:

$$\sum_{l \in \bar{L}_{\mu, \#}} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_\beta} \phi_{u+\beta} \nabla_{i_{\beta+1}} v \dots \nabla_{i_{\beta+\mu}} v = 0. \quad (2.19)$$

Proof of (2.19): We will use the technique (introduced in subsection 3.1 of [7]) of “inverse integration by parts” followed by the silly divergence formula.

Let us denote by \hat{C}_g^l the complete contraction that arises from each $C_g^{l, i_1 \dots i_{\mu+\beta}}$ by formally erasing the expression $\nabla_{s_{i_1}}^{(2)} \Omega_1 \nabla^s \phi_{u+1}$ and then making all free indices $i_{\beta+1}, \dots, i_{\beta+\mu}$ into internal contractions.²⁷ Then, the “inverse integration by parts” implies a new integral equation:

²⁶Notice that by weight considerations, if this property holds for one of the terms $C_g^{l, i_1 \dots i_\mu, i_{\mu+1} \dots i_{\mu+\beta}}$, then it will hold for all of them.

²⁷We recall that to “make a free index i_y into an internal contraction” means that we add a derivative ∇_{i_y} onto the factor T_{i_y} to which the free index i_y belongs. The new derivative index ∇^{i_y} is then contracted against the index i_y in T_{i_y} .

$$\int_{M^n} \sum_{l \in L_\mu} a_l \hat{C}_g^l + \sum_{j \in J} a_j C_g^j + \sum_{z \in Z} a_z C_g^z dV_g = 0. \quad (2.20)$$

Here the complete contractions indexed in J have length $\sigma + u$, u factors $\nabla \phi_u$ but are simply subsequent to the simple character $\vec{\kappa}_{simp}$. The terms indexed in Z either have length $\geq \sigma + u + 1$ or have length $\sigma + u$, but also at least one factor $\nabla^{(B)} \phi_h$ with $B \geq 2$.

Now, in the above, we consider the complete contractions indexed in $\bar{L}_{\mu, \#} \subset \bar{L}_\mu$ and we “pull out” the expression $\Delta \nabla_t \Omega_1 \nabla^t \phi_{u+1}$ to write:

$$\sum_{l \in L_{\mu, \#}} a_l \hat{C}_g^l = \sum_{l \in L_{\mu, \#}} a_l \bar{C}_g^l \cdot (\Delta \nabla_t \Omega_1 \nabla^t \phi_1).$$

Now, we consider the silly divergence formula applied to (2.20) obtained by integrating by parts with respect to the function Ω_1 . If we denote the integrand in (2.20) by F_g , we denote the resulting (local) equation by $silly[F_g] = 0$. We consider the sublinear combination $silly^*[F_g]$ which consists of terms with length $\sigma + u$, μ internal contractions and $u - 1 + \beta$ factors $\nabla \phi_h$, $h \geq 2$ and a factor $\Delta \phi_{u+1}$. Clearly, this sublinear combination must vanish separately modulo longer terms:

$$silly^*[F_g] = 0.$$

The above equation can be expressed as:

$$Spread^{\nabla^s, \nabla^s} \left[\sum_{l \in \bar{L}_{\mu, \#}} a_l \bar{C}_g^l \right] \cdot \Omega_1 \cdot \Delta \phi_{u+1} = 0. \quad (2.21)$$

(Here $Spread^{\nabla^s, \nabla^s}$ is a formal operation that acts on complete contractions in the form (1.5) by hitting a factor T in the form $\nabla^{(m)} R_{ijkl}$ or $\nabla^{(p)} \Omega_h$ with a derivative ∇^s and then hitting another factor $T' \neq T$ in the form $\nabla^{(m)} R_{ijkl}$ or $\nabla^{(p)} \Omega_h$ by a derivative ∇_s which contracts against ∇^s and then adding over all the terms we can thus obtain.) Now, using the fact that (2.21) holds formally, we derive:²⁸

$$\sum_{l \in L_{\mu, \#}} a_l \bar{C}_g^l = 0. \quad (2.22)$$

Thus, applying the operation Sub_ν $\mu - 1$ times to the above and then multiplying by $\nabla_{i_1 i_2} \Omega_1 \nabla^{i_1} \nu \nabla^{i_2} \phi_{u+1}$ we derive (2.19). So for the rest of this proof we may assume that $\bar{L}_{\mu, \#} = \emptyset$.

Now we prove our claim under the additional assumption that for the tensor fields indexed in \bar{L}_μ , the factor $\nabla^{(2)} \Omega_1$ contains no free index.

We again refer to the equation (2.20) and perform integrations by parts with respect to the factor $\nabla^{(B)} \Omega_1$. We denote the resulting local equation by $silly[L_g] = 0$. We pick out the sublinear combination $silly^*[L_g]$ of terms

²⁸This can be proven by using the operation $Erase[...]$, see the Appendix in [6].

with $\sigma + u$ factors, $u + \beta$ factors $\nabla\phi_h$, μ internal contractions, with $u + \beta - 1$ factors $\nabla\phi_h$, $h \geq 2$ and a factor $\Delta\phi_1$. This sublinear combination must vanish separately, $silly^*[L_g] = 0$; the resulting new true equation can be described easily: Let us denote by \hat{C}_g^{l,j_1} the 1-vector field that arises from $C_g^{l,i_1\dots i_\mu}$, $l \in L_{\mu,*}$ by formally erasing the factor $\nabla_{j_s}^{(2)}\Omega_1\nabla^s\phi_1$, making the index j that contracted against j into a free index j_1 , and making all the free indices i_1, \dots, i_μ into internal contractions. (Denote by $\bar{\kappa}'_{simp}$ the simple character of these vector fields). Then the equation $silly^*[L_g] = 0$ can be expressed in the form:

$$\sum_{l \in L_{\mu,*}} a_l \{Xdiv_{j_1} \hat{C}_g^{l,j_1}\} \Delta\phi_1 + \sum_{j \in J} a_j C_g^j \Delta\phi_1 = 0; \quad (2.23)$$

here the complete contractions C_g^j are simply subsequent to $\bar{\kappa}'_{simp}$. The above holds modulo terms of length $\geq \sigma + u + 1$. Now, we apply the operation Sub_ω μ times (see the Appendix in [3]). In the case $\sigma > 3$, applying the inductive assumption of our Lemma 2.6 to the resulting equation (notice that the above falls under the inductive assumption of this Lemma since we have lowered the weight in absolute value; we ensure that Lemma 2.6 can be applied by just labeling one of the factors $\nabla\omega$ into $\nabla\phi_{u+1}$). We derive (due to weight considerations) that there *can not be* tensor fields of higher rank, thus:

$$\sum_{l \in L_\mu} a_l Sub_\omega^{\mu-1} [\hat{C}_g^{l,j_1}] \nabla_{i_1} v \Delta\phi_1 = 0. \quad (2.24)$$

Now, formally replacing the factor $\nabla_{i_1} v$ by $\nabla_{j_1 t}^{(2)}\Omega_1\nabla^t\phi_1$, and then setting $\omega = v$, we derive the claim of our Lemma. In the case $\sigma = 3$ (2.24) follows by inspection, since the only two possible cases are $\sigma_2 = 2$ and $\sigma_1 = 2$; in the first case there are only two possible tensors field in \bar{L}_μ while in the second there are four. The equation (2.23) (by inspection) implies that the coefficients of all these tensor fields must vanish, which is equivalent to (2.24).

Now, we will prove our claim under the additional assumption $\bar{L}_\mu = \emptyset$ (still for $p > 0$). We again refer to (2.20) and again consider the same equation $silly[L_g] = 0$ as above. We now pick out the sublinear combination of terms with $\sigma + u$ factors, $u + \beta$ factors $\nabla\phi_h$, and μ internal contractions. We derive:

$$\sum_{l \in L_\mu} a_l Xdiv_{j_1} Xdiv_{j_2} \hat{C}_g^{l,j_1 j_2} + \sum_{j \in J} a_j C_g^j = 0; \quad (2.25)$$

here the terms $\hat{C}_g^{l,j_1 j_2}$ arise from the μ -tensor fields $C_g^{l,i_1\dots i_\mu}$ by replacing all μ free indices by internal contractions, erasing the factor $\nabla_{j^k}^{(2)}\Omega_1$ and making the indices j, k into free indices j_1, j_2 . Now, applying Sub_ω μ times, and then applying the inductive assumption of Lemma 4.10 2.6 (this applies by length considerations as above for $\sigma > 3$; while if $\sigma = 3$ the claim (2.26) will again follow by inspection) we derive:

$$\sum_{l \in L_\mu} a_l \hat{C}_g^{l, j_1 j_2} \nabla_{j_1} v \nabla_{j_2} v = 0; \quad (2.26)$$

Replacing the expression $\nabla_{j_1} v \nabla_{j_2} v$ by a factor $\nabla_{j_1 j_2}^{(2)} \Omega_2$ and then setting $\omega = v$, we derive our claim in this case $p > 0$.

The case $p = 0, \sigma_1 > 0$: We will reduce ourselves to the previous case: We let L_μ^1 the index set of μ -tensor fields where the factor $T_1 = S_* R_{ijkl} \nabla^i \tilde{\phi}_1$ contains a special free index (say the index k is the free index $i_{\beta+1}$ wlog). We will prove our claim for the index set L_μ^1 ; if we can prove this, then clearly our Lemma will follow by induction.

To prove this claim, we consider the first conformal variation of our hypothesis, $Image_Y^1[L_g] = 0$, and we pick out the sublinear combination of terms with length $\sigma + u + \beta$, with the factor $\nabla^{(\nu)} S_* R_{ijkl} \nabla^i \tilde{\phi}_1$ has been replaced by a factor $\nabla^{(\nu+2)} Y$, and the factor $\nabla \phi_1$ now contracts against a factor $T_2 = R_{ijkl}$. This sublinear combination vanishes separately, thus we derive a new local equation. To describe the resulting equation, we denote by

$\hat{C}_g^{l, i_1 \dots i_{\beta+1} \dots i_{\mu+\beta}}(Y, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_\beta} \phi_{u+\beta}$ the $(\mu - 1)$ -tensor field that arises from by formally replacing the factor by and also adding a derivative index ∇_{i_*} onto the factor $T_2 = R_{ijkl}$ and then contracting that index against a factor $\nabla \phi_1$. Denote the $(u - 1)$ -simple character of the above (the one defined by $\nabla \phi_2, \dots, \nabla \phi_u$) by $\tilde{\kappa}'_{simp}$. We then have an equation:

$$\begin{aligned} & \sum_{l \in L_\mu^1} a_l X \operatorname{div}_{i_{\beta+2}} \dots X \operatorname{div}_{i_{\beta+\mu}} \hat{C}_g^{l, i_1 \dots i_{\beta+1} \dots i_{\mu+\beta}}(Y, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_\beta} \phi_{u+\beta} \\ & + \sum_{h \in H} a_h X \operatorname{div}_{i_{\beta+1}} \dots X \operatorname{div}_{i_{\beta+\mu}} C_g^{l, i_2 \dots i_{\mu+\beta}}(Y, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_{\beta+\mu}} \phi_{u+\beta} \\ & + \sum_{j \in J} a_j C_g^{j, i_{\mu+1} \dots i_{\mu+\beta}}(Y, \phi_1, \dots, \phi_u) \nabla_{i_{\mu+1}} \phi_{u+1} \dots \nabla_{i_{\mu+\beta}} \phi_{u+\beta}. \end{aligned} \quad (2.27)$$

The terms indexed in H are acceptable, have a $(u - 1)$ -simple character $\tilde{\kappa}'_{simp}$ and the factor $\nabla \phi_1$ contracts against the index i in the factor $T_2 = R_{ijkl}$; writing that factor as $S_* R_{ijkl} \nabla^i \tilde{\phi}_1$, we denote the resulting u -simple factor by $\tilde{\kappa}_{simp}$. The terms indexed in J are simply subsequent to $\tilde{\kappa}'_{simp}$. Now, applying the inductive assumption of Lemma 2.6,²⁹ we derive that:

$$\sum_{h \in H} a_h C_g^{l, i_2 \dots i_{\mu+\beta}}(Y, \phi_1, \dots, \phi_u) \nabla_{i_1} \phi_{u+1} \dots \nabla_{i_{\beta+1}} \phi_{u+\beta} \nabla_{i_{\beta+1}} v \dots \nabla_{i_{\beta+\mu}} v = 0. \quad (2.28)$$

²⁹The terms indexed in L_μ^1 are now simply subsequent to $\tilde{\kappa}_{simp}$.

Thus, we may assume wlog that $H = \emptyset$ in (2.27). Now, we again apply Lemma 2.6 to (2.27) (under that additional assumption), and we derive that:

$$\sum_{l \in L_\mu^1} a_l \hat{C}_g^{l, i_1 \dots i_{\beta+1} \dots i_{\mu+\beta}}(Y, \phi_1, \dots, \phi_u) \nabla_1 \phi_{u+1} \dots \nabla_{i_{\beta+1}} \phi_{u+\beta} \nabla_{i_{\beta+2}} v \dots \nabla_{i_{\beta+\mu}} v = 0. \quad (2.29)$$

Now, erasing the factor $\nabla \phi_1$ from the above, and then formally replacing the factor $\nabla_{ab}^{(2)} Y$ by $S_* R_{i(ab)l} \nabla^i \tilde{\phi}_1 \nabla^l v$, we derive our claim.

The case $p = 0, \sigma_1 = 0$: In this case $\sigma = \sigma_2$. In other words, all factors in $\vec{\kappa}_{simp}$ are simple factors in the form $S_* R_{ijkl} \nabla^i \tilde{\phi}_h$. We recall that in this case all μ -tensor fields in (2.9) must have at most one free index in any factor $S_* R_{ijkl}$. In that case, we will prove our claim in a more convoluted manner, again reducing ourselves to the inductive assumption of Proposition 2.1 in [6].

A key observation is the following: By the definition of the special cases, $\mu + \beta \leq \sigma_2$. In the case of strict inequality, we see (by a counting argument) that at least one of the special indices in one of the factors $S_* R_{ijkl}$ must contract against a special index in another factor $S_* R_{abcd}$. In the case $\mu + \beta = \sigma_2$ this remains true, *except* for the terms for which the β factors $\nabla \phi_{u+h}$ contract against special indices, say the indices k , in β factors $T_y = S_* R_{ikl} \nabla^i \tilde{\phi}_y$, and moreover these factors must not contain a free index, and all other factors $S_* R_{ikl}$ contain exactly one free index, which must be special. In this subcase, we will prove our claim for all μ -tensor fields *excluding* this particular “bad” sublinear combination; we will prove our claim for this sublinear combination in the end.

We will now proceed to *normalize* the different $(\mu + \beta)$ -tensor fields in (2.9). A *normalized* tensor field will be in the form (1.6), with possibly certain pairs of indices in certain of the factors $S_* R_{ijkl}$ being symmetrized.

Let us first introduce some definitions: Given each $C_g^{l, i_1 \dots i_\mu}$, we list out the factors T_1, \dots, T_{σ_2} in the form $S_* R_{ikl}$. Here T_a is the factor for which the index i is contracting against the factor $\nabla \tilde{\phi}_a$. We say that factors $S_* R_{ikl}$ are type I if they contain no free index. We say they are of type II if they contain a special free index. We say they are of type III if they contain a non-special free index.

Given any tensor field $C_g^{l, i_1 \dots i_\mu}$ in the form (1.6), pick out the pairs of factors, T_α, T_β in the form $S_* R_{ijkl}$ for which a special index in T_α contracts against a special index in T_β . (Call such particular contractions “special-to-special” particular contractions). Now, in any $C_g^{l, i_1 \dots i_\mu}$ we define an ordering among all its factors $S_* R_{ijkl}$: The factor $T_a = S_* R_{ikl} \nabla^i \tilde{\phi}_a$ is more important than $T_b = S_* R_{i'j'k'l'} \nabla^{i'} \tilde{\phi}_b$ if $a < b$.

Now, consider a tensor field $C_g^{l, i_1 \dots i_\mu}$ and list out all the pairs of factors T_a, T_b with a special-to-special particular contraction. We say that (T_a, T_b) is the *most important* pair of factors with a special-to-special particular contraction³⁰ if any other such pair (T_c, T_d) ,³¹ has either T_c being less important than T_a or $T_a = T_c$

³⁰Assume wlog that T_a is more important than T_b

³¹Again assume wlog that T_c is more important than T_d .

and T_d less important than T_b .

Now, consider a tensor field $C_g^{l,i_1 \dots i_\mu}$ and consider the most important pair of factors (T_a, T_b) with a special-to-special particular contraction. Assume wlog that the index l in $T_a = S_* R_{ijkl} \nabla^i \tilde{\phi}_a$ contracts against the index l' in $T_b = S_* R_{i'j'k'l'} \nabla^{i'} \tilde{\phi}_b$. We say that $C_g^{l,i_1 \dots i_\mu}$ is normalized if both factors T_a, T_b are normalized. The factor $T_a = S_* R_{ijkl} \nabla^i \tilde{\phi}_a$ is normalized if: Either the index j contracts against a factor T_c which is more important than T_b , or if the indices j, k are symmetrized. If T_a is of type II, then we require that the index j in $T_b = S_* R_{ij(\text{free})l}$ must be contracting against a special index of some other factor T_c , and moreover T_c must be more important than T_b . If T_a is of type III, then it is automatically normalized. The same definition applies to T_b , where any reference to T_b must be replaced by a reference to T_a .

Let us now prove that we may assume wlog that all μ -tensor fields in (2.9) are normalized: Consider a $C_g^{l,i_1 \dots i_\mu}$ in (2.9) for which the most important pair of factors with a special-to-special particular contraction is the pair (T_a, T_b) . We will prove that we can write:

$$C_g^{l,i_1 \dots i_{\mu+\beta}} = \tilde{C}_g^{l,i_1 \dots i_{\mu+\beta}} + \sum_{t \in T} a_t C_g^{t,i_1 \dots i_{\mu+\beta}}; \quad (2.30)$$

here the term $\tilde{C}_g^{l,i_1 \dots i_{\mu+\beta}}$ is normalized, the most important pair of factors with a special-to-special particular contraction is the pair (T_a, T_b) , and moreover its refined double character is the same as for $C_g^{l,i_1 \dots i_{\mu+\beta}}$. Each $C_g^{t,i_1 \dots i_{\mu+\beta}}$ has either the same, or a doubly subsequent refined double character to $C_g^{l,i_1 \dots i_{\mu+\beta}}$; moreover in the first case its most important pair of factors with a special-to-special particular contraction will be less important than the pair (T_a, T_b) . In the second case the most important pair will either be (T_a, T_b) or a less important pair.

Clearly, if we can prove the above, then by iterative repetition we may assume wlog that all $(\mu + \beta)$ -tensor fields in (2.9) are normalized.

Proof of (2.30): Pick out the most important pair of factors with a special-to-special particular contraction is the pair (T_a, T_b) in $C_g^{l,i_1 \dots i_{\mu+\beta}}$. Let us first normalize T_a . If T_a is of type III, there is nothing to do. If it is of type II and already normalized, there is again nothing to do. If it is of type II and not normalized, then we *interchange* the indices j, k . The resulting factor is normalized. The correction term we obtain by virtue of the first Bianchi identity is also normalized (it is of type III). Moreover, the resulting tensor field is doubly subsequent to $C_g^{l,i_1 \dots i_{\mu+\beta}}$. Finally, if the factor T_a is of type I, we inquire on the factor T_c against which j in $T_a = S_* R_{ijkl}$ contracts: If it is more important than T_b , then we leave T_a as it is; it is already normalized. If not, we symmetrize j, k . The resulting tensor field is normalized. The correction term we obtain by virtue of the first Bianchi identity will then have the same refined double character as $C_g^{l,i_1 \dots i_{\mu+\beta}}$, and moreover its most important pair of factors with a special-to-special particular contraction is less important than that pair (T_a, T_b) .

We now prove the claim of Lemma 2.6 in this special case, under the additional assumption that all tensor fields in (2.9) are normalized. We list out the most important pair of special-to-special particular contractions in each $C_g^{l, i_1 \dots i_{\mu+\beta}}$, and denote it by $(a, b)_l$. We let (α, β) stand for the lexicographically minimal pair among the list $(a, b)_l, l \in L_\mu$. We denote by $L_\mu^{(\alpha, \beta)} \subset L_\mu$ the index set of terms with a special-to-special particular contraction among the terms T_α, T_β . We will prove that:

$$\sum_{l \in L_\mu^{(\alpha, \beta)}} a_l C_g^{l, i_1 \dots i_{\mu+\beta}} \nabla_{i_1} v \dots \nabla_{i_\mu} v = 0. \quad (2.31)$$

Clearly, the above will imply our claim, by iterative repetition.³²

Proof of (2.31): Consider $Image_{Y_1, Y_2}^2[L_g] = 0$ and pick out the sublinear combination where the factors T_α, T_β are replaced by $\nabla^{(A)} Y_1 \otimes g, \nabla^{(B)} Y_2 \otimes g$, and the two factors $\nabla \tilde{\phi}_\alpha, \nabla \tilde{\phi}_\beta$ contract against each other. The resulting sublinear combination must vanish separately. We erase the expression $\nabla_t \tilde{\phi}_\alpha \nabla^t \tilde{\phi}_\beta$,³³ and derive a new true equation which will be in the form:

$$\sum_{l \in L_\mu^{(\alpha, \beta)}} a_l X \text{div}_{i_1} \dots X \text{div}_{i_\mu} \tilde{C}_g^{l, i_1 \dots i_{\mu+\beta}}(\Omega_1, Y_1, Y_2) + \sum_{j \in J} a_j C_g^j(\Omega_1, Y_1, Y_2) = 0; \quad (2.32)$$

here the tensor fields $\tilde{C}_g^{l, i_1 \dots i_{\mu+\beta}}(\Omega_1, Y_1, Y_2)$ arise from the tensor fields $C_g^{l, i_1 \dots i_{\mu+\beta}}$ by replacing the expression $\nabla^i \tilde{\phi}_\alpha S_* R_{ijkl} \otimes S_* R_{i'jk}{}^l \nabla^{i'} \tilde{\phi}_\beta$ by an expression $\nabla_{jk} Y_1 \otimes \nabla_{j'k'} Y_2$ (notice we have lowered the weight in absolute value).

Now, applying the inductive assumption of Lemma 2.6 to the above,³⁴ we derive:

$$\sum_{l \in L_\mu^{(\alpha, \beta)}} a_l \tilde{C}_g^{l, i_1 \dots i_{\mu+\beta}}(\Omega_1, Y_1, Y_2) \nabla_{i_1} v \dots \nabla_{i_\mu} v = 0; \quad (2.33)$$

The proof of (2.31) is only one step away. Let us start with an important observation: For each given complete contraction above, examine the factor $\nabla_{zx}^{(2)} Y_1$; it either contracts against no factor ∇v or one factor ∇v .³⁵ In the first case, the factor $\nabla_{zx}^{(2)} Y_1$ must have arisen from a factor $S_* R_{ijkl}$ of type I. In fact, the indices z, x correspond to the indices j, k in the original factor, and we can even determine their position: Since the pair (α, β) is the most important pair in (2.9), at most one of the indices z, x can contract against a special index in a more important factor than T_β . If one of them does (say z),

³²In the subcase $\mu + \beta = \sigma_2$ it will only imply it for the “excluded” sublinear combination defined above.

³³Denote the resulting $(u - 2)$ -simple character by $\tilde{\kappa}_{simp}'''$.

³⁴We have lowered the weight in absolute value.

³⁵The two corresponding sublinear combinations vanish separately, of course.

then that index must have been the index j in $T_\alpha = S_*R_{ikl}$. If none of them does, then the two indices z, x must be symmetrized over, since the two indices j, k in T_α to which they correspond were symmetrized over. Now, these two separate sublinear combinations in (2.33) must vanish separately (this can be proven using the eraser from the Appendix in [3]), and furthermore in the first case, we may assume that the index z (which contracts against a special index in a more important factor than T_β) occupies the leftmost position in $\nabla_{zx}^{(2)}Y_1$ and is *not permuted* in the formal permutations of indices that make (2.33) hold *formally*).

On the other hand, consider the terms in (2.33) with the factor $\nabla^{(2)}Y_1$ contracting against a factor ∇v . By examining the index y in the factor $\nabla_{yt}^{(2)}Y_1\nabla^t v$, we can determine the *type* of factor in $C_g^{l, i_1 \dots i_{\mu+\beta}}$ from which the factor $\nabla^{(2)}Y_1$ arose: If the index y is contracting against a special index in a factor S_*R_{ijkl} which is *more important* than T_β , then $\nabla^{(2)}Y_1$ can only have arisen from a factor of type II in $C_g^{l, i_1 \dots i_{\mu+\beta}}$. In fact, the index y in $\nabla^{(2)}Y_1$ must correspond to the index j in $S_*R_{ij(\text{free})l}$ in T_α . If the index y in $\nabla_{yt}^{(2)}Y_1\nabla^t v$ does not contract against a special index in a factor T_c which is more important than T_β , then the factor $\nabla^{(2)}Y_1$ can only have arisen from a factor of type III in $C_g^{l, i_1 \dots i_{\mu+\beta}}$. In fact, the index y in $\nabla^{(2)}Y_1$ must correspond to the index k in $S_*R_{i(\text{free})kl}$ in T_α .

The same analysis can be repeated for the factor $\nabla^{(2)}Y_2$, with any reference to the factor T_β now replaced by the factor T_α .

In view of the above analysis, we can break the LHS of (2.33) into four sublinear combinations which vanish separately (depending on whether $\nabla^{(2)}Y_1, \nabla^{(2)}Y_2$ contract against a factor ∇v or not). Then in each of the four sublinear combinations, we can arrange that in the formal permutations that make the LHS of (2.33) formally zero, the two indices in the factors $\nabla^{(2)}Y_1, \nabla^{(2)}Y_2$ are *not permuted* (by virtue of the remarks above). In view of this and the analysis in the previous paragraph, we can then *replace* the two factors $\nabla_{zx}^{(2)}Y_1, \nabla_{qw}^{(2)}Y_2$ by an expression $\nabla^i \tilde{\phi}_\alpha S_*R_{izxl} \otimes S_*R_{i'qw} \nabla^{i'} \tilde{\phi}_\beta$, in such a way that the resulting linear combination *vanishes formally without permuting the two indices* q, w, q', w' . This proves our claim, except for the subcase $\mu + \beta = \sigma_2$ where we only derive our claim for all terms except for the “bad sublinear combination”. We now prove our claim for that.

We then break up the LHS of (2.16) according to which factor T_s the factor $\nabla\phi_{u+1}$ contracts—denote the index set of those terms by L_μ^K . Denote the resulting sublinear combinations by $L_g^K, K = 1, \dots, \sigma_2$. Given any K , we consider the equation $\text{Image}_Y^\perp[L_g] = 0$, and we pick out the sublinear combination where the term $\nabla^{(B)}S_*R_{ijkl}\nabla^i\tilde{\phi}_K$, is replaced by $\nabla^{(B+2)}Y$, and the factor $\nabla\phi_K$ now contracts against the factor $\nabla\phi_{u+1}$. This sublinear combination must vanish separately. We then again perform the “inverse integration by parts” to this true equation (deriving an integral equation), and then we consider the silly divergence formula for this integral equation, obtained by integrating by parts with

respect to $\nabla^{(B)}Y$. We pick out the sublinear combination with $\sigma + u + \beta$ factors, μ internal contractions and $u + \beta$ factors $\nabla\phi_h$, and a expression $\nabla_s\phi_{u+1}\nabla^s\tilde{\phi}_K$. This gives us a new true local equation:

$$\sum_{l \in L_\mu^K} a_l X_* \text{div}_{j_1} X_* \text{div}_{j_2} \tilde{C}_g^{l, j_1 j_2} + \sum_{j \in J} a_j C_g^j = 0. \quad (2.34)$$

Here the tensor fields $\tilde{C}_g^{l, j_1 j_2}$ arise from $C_g^{l, i_1 \dots i_\mu}$ by formally replacing all μ free indices by internal contractions, and also replacing the expression $\nabla_x \phi_{u+1} \otimes S_* R_{i(jk)}^x \nabla^i \tilde{\phi}_K$ by $\nabla_x \phi_{u+1} \nabla^s \tilde{\phi}_K \otimes Y$, and then making the indices j, k that contracted against j, k into free indices j^1, j^2 . $X_* \text{div}_j$ stands for the sublinear combination in $X \text{div}_j$ where ∇^j is not allowed to hit the factor Y . Now, applying the inductive assumption of Lemma 2.6 to the above,³⁶ we derive that:

$$\sum_{l \in L_\mu^K} a_l \tilde{C}_g^{l, j_1 j_2} \nabla_{j_1} \omega \nabla_{j_2} \omega = 0.$$

Now, we replace the expression $\nabla^x \phi_K \nabla_x \phi_{u+1} \nabla_{j_1} \omega \nabla_{j_2} \omega Y$ by $\nabla^l \phi_{u+1} S_* R_{i(j_1 j_2)l} \nabla^i \tilde{\phi}_K$ and then replacing all internal contractions by factors ∇v (applying the operation $Su b_v$ from the Appendix in [3]). The resulting (true) equation is precisely our remaining claim for the “bad” sublinear combination. \square

2.3 Proof of Lemmas 4.6, 4.8 in [6]: The main part.

We first write down the form of the complete and partial contractions that we are dealing with in Lemmas 2.1 and 2.3. In the setting of Lemma 2.1 we recall that the tensor fields $C^{h, i_1 \dots i_\alpha}$ indexed in H_2 (in the hypothesis of Lemma 2.1) are all partial contractions in the form:

$$\begin{aligned} & p \text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \dots \otimes \nabla^{(m_{\sigma_1})} R_{ijkl} \otimes S_* \nabla^{(\nu_1)} R_{ijkl} \otimes \dots \otimes S_* \nabla^{(\nu_t)} R_{ijkl} \otimes \\ & \nabla^{(b_1)} \Omega_1 \otimes \dots \otimes \nabla^{(b_p)} \Omega_p \otimes \nabla Y \otimes \\ & \nabla \phi_{z_1} \dots \otimes \nabla \phi_{z_f} \otimes \nabla \phi'_{z_{f+1}} \otimes \dots \otimes \nabla \phi'_{z_{f+d}} \otimes \dots \otimes \nabla \tilde{\phi}_{z_{f+d+1}} \otimes \dots \otimes \nabla \tilde{\phi}_{z_{f+d+y}}), \end{aligned} \quad (2.35)$$

where we let $f + d + y = u'$. The main assumption here is that all tensor fields have the same u' -simple character (the one defined by $\nabla\phi_1, \dots, \nabla\phi_{u'}$), which we denote by $\tilde{\kappa}_{simp}^+$. The other main assumption is that if we formally treat the factor ∇Y as a function $\nabla\phi_{u+1}$, then the hypothesis of Lemma 2.1 falls under the inductive assumptions of Proposition 1.1 (i.e. the weight, real length, Φ and p are as in our inductive assumption of Proposition 1.1).

In the setting of Lemma 2.3 we recall that we are dealing with complete and partial contractions in the form:

³⁶We have lowered the weight in absolute value.

$$\begin{aligned}
& \text{contr}(\nabla^{(m_1)} R_{ijkl} \otimes \cdots \otimes \nabla^{(m_{\sigma_1})} R_{ijkl} \otimes S_* \nabla^{(\nu_1)} R_{ijkl} \otimes \cdots \otimes S_* \nabla^{(\nu_t)} R_{ijkl} \otimes \\
& \nabla^{(b_1)} \Omega_1 \otimes \cdots \otimes \nabla^{(b_p)} \Omega_p \otimes [\nabla \omega_1 \otimes \nabla \omega_2] \otimes \\
& \nabla \phi_{z_1} \cdots \otimes \nabla \phi_{z_f} \otimes \nabla \phi'_{z_{f+1}} \otimes \cdots \otimes \nabla \phi'_{z_{f+d}} \otimes \cdots \otimes \tilde{\phi}_{z_{f+d+1}} \otimes \cdots \otimes \tilde{\phi}_{z_{f+d+y}}),
\end{aligned} \tag{2.36}$$

where we let $f + d + y = u'$. The main assumption here is that all partial contractions have the same u' -simple character (the one defined by $\nabla \phi_1, \dots, \nabla \phi_{u'}$), which we denote by $\vec{\kappa}_{simp}^+$. The other main assumption is that if we formally treat the factors $\nabla \omega_1, \nabla \omega_2$ as factors $\nabla \phi_{u+1}, \nabla \phi_{u+2}$, then the hypothesis of Lemma 2.3 falls under the inductive assumptions of Proposition 1.1 (i.e. the weight, real length, Φ and p are as in our inductive assumption of Proposition 1.1).

Note: From now on, we will be writing $u' = u$, for simplicity. We will also be writing $\vec{\kappa}_{simp}^+ = \vec{\kappa}_{simp}$, for simplicity. We will also be labelling the indices i_1, \dots, i_α as $i_{\pi+1}, \dots, i_{\alpha+1}$.

New induction: We will now prove the two Lemmas 2.1, 2.3 by a new induction on the weight of the complete contractions in the hypotheses of those Lemmas. We will assume that these two Lemmas are true when the weight of the complete contractions in their hypotheses is $-W$, for any $W < K \leq n$. We will then show our Lemmas for weight $-K$.

Reduce Lemma 2.1 to two Lemmas: In order to show Lemma 2.1, we further break up H_2 into subsets: We say that $h \in H_2^a$ if and only if $C^{h, i_{\pi+1} \dots i_{\alpha+1}}$ has a free index (say the free index $i_{\alpha+1}$ wlog) belonging to the factor ∇Y . On the other hand, we say that $h \in H_2^b$ if the index in the factor ∇Y is not free. Lemma 2.1 will then follow from Lemmas 2.7, 2.8 below:

Lemma 2.7 *There exists a linear combination of acceptable $(\alpha - \pi + 1)$ -tensor fields, $\sum_{v \in V} a_v C_g^{v, i_{\pi+1} \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$, where the index $i_{\alpha+1}$ belongs to the factor ∇Y , with a simple character $\vec{\kappa}_{simp}$, so that:*

$$\begin{aligned}
& \sum_{h \in H_2^a} a_h C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v = \\
& \sum_{v \in V} a_v X_* \text{div}_{i_{\alpha+2}} C_g^{v, i_{\pi+1} \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v + \\
& \sum_{j \in J} a_j C_g^{j, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v.
\end{aligned} \tag{2.37}$$

Each C^j is simply subsequent to $\vec{\kappa}_{simp}$.

We observe that if we can show our first claim, then we can assume, with no loss of generality, that $H_2^a = \emptyset$, since it immediately follows from the above that:

$$\begin{aligned}
& \sum_{h \in H_2^a} a_h X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) = \\
& \sum_{v \in V} a_v X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} X_* \operatorname{div}_{i_{\alpha+2}} C_g^{v, i_{\pi+1} \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \\
& + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.38}$$

where each complete contraction C^j is subsequent to $\vec{\kappa}_{simp}$. (Note that one of the free indices in the tensor fields $C_g^{v, i_{\pi+1} \dots i_{\alpha+2}}$ will belong to the factor ∇Y).

Second claim, in the setting of Lemma 2.1:

Lemma 2.8 *We assume $H_2^a = \emptyset$. We then claim that modulo complete contractions of length $\geq \sigma + u + 1$:*

$$\begin{aligned}
& \sum_{h \in H_2} a_h C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v = \\
& \sum_{t \in T} a_t X_* \operatorname{div}_{i_{\alpha+2}} C_g^{t, i_{\pi+1} \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.39}$$

where each C^j is acceptable and subsequent to $\vec{\kappa}_{simp}$.

We observe that if we can show the above two Lemmas then Lemma 2.1 will follow. (Notice that replacing by the RHS of (2.38) into the hypothesis of Lemma 2.1, we do not introduce 1-forbidden terms).

We make two analogous claims for Lemma 2.3:

Reduce Lemma 2.3 to two Lemmas: We say that $h \in H_2^a$ if and only if $C^{h, i_{\pi+1} \dots i_{\alpha+1}}$ has a free index belonging to one of the factors $\nabla \omega_1, \nabla \omega_2$. On the other hand, we say that $h \in H_2^b$ if in none of the factors $\nabla \omega_1, \nabla \omega_2$ in $C^{h, i_{\pi+1} \dots i_{\alpha+1}}$ contains a free index. (Observe that we may assume with no loss of generality that there are no tensor fields $C^{h, i_{\pi+1} \dots i_{\alpha+1}}$ with free indices in both factors $\nabla \omega_1, \nabla \omega_2$ -this is by virtue of the anti-symmetry of the factors $\nabla \omega_1, \nabla \omega_2$). We make two claims. Firstly:

Lemma 2.9 *There is a linear combination of acceptable $(\alpha - \pi + 1)$ -tensor fields, $\sum_{v \in V} a_v C_g^{v, i_{\pi+1} \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u)$, in the form (2.35) with a simple character $\vec{\kappa}_{simp}$, so that:*

$$\begin{aligned}
& \sum_{h \in H_2^a} a_h X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{\alpha+1}} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) = \\
& \sum_{v \in V} a_v X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{\alpha+1}} X_+ \operatorname{div}_{i_{\alpha+2}} C_g^{v, i_{\pi+1} \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{\alpha+1}} C_g^{q, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \nabla_+ [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u).
\end{aligned} \tag{2.40}$$

(Recall that by definition the complete contractions indexed in Q have a factor $\nabla^{(2)}\omega_1$).

We observe that if we can show our first claim, then we can, with no loss of generality, assume that $H_2^a = \emptyset$.

Second claim:

Lemma 2.10 *We assume $H_2^a = \emptyset$, and that for some $k \geq 1$, we can write:*

$$\begin{aligned}
& \sum_{h \in H_2^b} a_x X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{\alpha+1}} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{t \in T_k} a_t X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{\alpha+k}} C_g^{t, i_{\pi+1} \dots i_{\alpha+k}}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{\alpha+1}} C_g^{q, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \nabla_+ [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.41}$$

where the last two linear combinations on the right hand side of the above are generic linear combinations in the form described in the claim of Lemma 2.3.³⁷ On the other hand, $\sum_{t \in T_k} a_t C_g^{t, i_{\pi+1} \dots i_{\alpha+k}}(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u)$ is a linear combination of acceptable $(\alpha - \pi + k)$ -tensor fields in the form (2.36) with a simple character $\vec{\kappa}_{simp}$, and with two anti-symmetric factors $\nabla\omega_1, \nabla\omega_2$ which do not contain a free index. We then claim that modulo complete contractions of length $\geq \sigma + u + 1$ we can write:

³⁷In Lemma 2.3, Q is called V .

$$\begin{aligned}
& \sum_{t \in T_k} a_t X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_{a+k}} C_g^{t, i_1 \dots i_{a+k}} (\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) = \\
& \sum_{t \in T_{k+1}} a_t X_+ \operatorname{div}_{i_{\pi+1}} \dots X_+ \operatorname{div}_{i_{a+k+1}} C_g^{t, i_{\pi+1} \dots i_{a+k+1}} (\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_{a+1}} C_g^{q, i_1 \dots i_{a+1}} (\Omega_1, \dots, \Omega_p, \nabla_+ [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \\
& + \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.42}$$

with the same notational conventions as above.

We observe that if we can show the above two claims, then Lemma 2.3 will follow by iterative repetition of the second claim.

We will now show the four Lemmas above.

Proof of Lemmas 2.8 and 2.10: Lemma 2.8 is a direct consequence of Lemma 4.10 in [6].³⁸ Lemma 2.10 can be proven in two steps: Firstly, by Lemma 2.5 we derive that there exists a linear combination of acceptable $(a+k+1)$ -tensor fields (indexed in X below) with a u-simple character $\vec{\kappa}_{simp}$ so that:

$$\begin{aligned}
& \sum_{t \in T_k} a_t C_g^{t, i_1 \dots i_{a+k}} (\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \nabla_{i_1} v \dots \nabla_{i_{a+k}} v - \\
& \sum_{t \in T_{k+1}} a_t X_* \operatorname{div}_{i_{a+k+1}} C_g^{t, i_{\pi+1} \dots i_{a+k+1}} (\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u) \nabla_{i_1} v \dots \nabla_{i_{a+k}} v \\
& + \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u, v^{a+k}),
\end{aligned} \tag{2.43}$$

where the complete contractions indexed in J have length $\sigma + a + k + 1$ and are simply subsequent to $\vec{\kappa}_{simp}$.

Then, making the factors ∇v in the above into $X_+ \operatorname{div}s$, we derive Lemma 2.10. \square

Proof of Lemma 2.7:

We have denoted by $\vec{\kappa}_{simp}$ the simple character of our tensor fields. We distinguish two cases: In case A there is a factor $\nabla^{(m)} R_{ijkl}$ in $\vec{\kappa}_{simp}$, and in case B there is no such factor.

We denote $\alpha + 1 = \gamma$, for brevity.

³⁸Observe that our hypotheses on the tensor fields in the equation in Lemma 2.1 not being “bad” ensure that we do not fall under the “forbidden” cases of Lemma 4.10 in [6].

Now we break the set H_2^b into subsets: In case A we say that $h \in H_2^{b,+}$ if and only if ∇Y is contracting against an internal index of a factor $\nabla^{(m)}R_{ijkl}$. In case B we say that $h \in H_2^{b,+}$ if and only if ∇Y is contracting against one of the indices k, l in a factor $S_*\nabla^{(\nu)}R_{ijkl}$.

We define $H_2^{b,-} = H_2^b \setminus H_2^{b,+}$.

In each of the above cases and subcases we treat the function ∇Y as a function $\nabla\phi_{u+1}$ in our Lemma hypothesis. Then, by applying the first claim in Lemma 4.10 in [6]³⁹ to our Lemma hypothesis and then making the ∇ vs into X_*divs , we derive that we can write:

$$\begin{aligned} & X_*div_{i_{\pi+1}} \dots X_*div_{i_\gamma} \sum_{h \in H_2^{b,+}} a_h C_g^{h, i_{\pi+1} \dots i_\alpha, i_\gamma}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) = \\ & X_*div_{i_{\pi+1}} \dots X_*div_{i_\gamma} \sum_{h \in H_2^{b,*,-}} a_h C_g^{h, i_{\pi+1} \dots i_\alpha, i_\gamma}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \\ & + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u), \end{aligned} \tag{2.44}$$

where $\sum_{h \in H_2^{b,*,-}} a_h C_g^{h, i_{\pi+1} \dots i_\alpha, i_\gamma}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$ stands for a *generic* linear combination as defined above (i.e. it is in the general form $\sum_{h \in H_2^b} \dots$ but the factor ∇Y is *not* contracting against a special index in any factor $\nabla^{(m)}R_{ijkl}$ or $S_*\nabla^{(\nu)}R_{ijkl}$.⁴⁰ On the other hand, each $C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$ is a complete contraction with a simple character that is subsequent to $\vec{\kappa}_{simp}$.

Thus, by virtue of (2.44), we reduce ourselves to the case where $H_2^{b,+} = \emptyset$. We will then show Lemma 2.7 separately in cases A and B, under the assumption that $H_2^{b,+} = \emptyset$.

Proof of Lemma 2.7 in case A: We will define the C-crucial factor, for the purposes of this proof only: We denote by *Set* the set of numbers u for which $\nabla\phi_u$ is contracting against one of the factors $\nabla^{(m)}R_{ijkl}$. If $Set \neq \emptyset$, we define u_+ to be the minimum element of *Set*, and we pick out the factor $\nabla^{(m)}R_{ijkl}$ in each C^h against which $\nabla\phi_{u_+}$ contracts. We call that factor $\nabla^{(m)}R_{ijkl}$ *C-crucial*. If $Set = \emptyset$, we will say *the C-crucial factors* and will mean *any* of the factors $\nabla^{(m)}R_{ijkl}$.

Now, we pick out the subset $H_2^{b,*} \subset H_2^b$, which is defined by the rule: $h \in H_2^{b,*}$ if and only if ∇Y is contracting against the (one of the) C-crucial factor(s).

Now, for each $h \in H_2^a$ we denote by

$$Hitdiv_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$$

³⁹By weight considerations, since we started out with no “bad terms” in Lemma 2.1, we will not encounter no “forbidden tensor fields” for Lemma 4.10 in [6].

⁴⁰Recall that a special index in a factor $\nabla^{(m)}R_{ijkl}$ is an internal index, while a special index in a factor $S_*\nabla^{(\nu)}R_{ijkl}$ is an index k, l .

the sublinear combination in $X_*div_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$ that arises when ∇_{i_γ} hits the (one of the) C-crucial factor.⁴¹ It then follows that:

$$\begin{aligned} & \sum_{h \in H_2^a} a_h X_*div_{i_{\pi+1}} \dots X_*div_{i_\alpha} Hitdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \\ & + \sum_{h \in H_2^{b,*}} a_h X_*div_{i_{\pi+1}} \dots X_*div_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u), \end{aligned} \tag{2.45}$$

where each C_g^j has the factor ∇Y contracting against the C-crucial factor $\nabla^{(m)}R_{ijkl}$ and is simply subsequent to $\vec{\kappa}_{simp}$.

We now denote the $(u+1)$ -simple character (the one defined by $\nabla\phi_1, \dots, \nabla\phi_{u+1} = \nabla Y$) of the tensor fields $Hitdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}, i_\gamma}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$ by $\vec{\kappa}'_{simp}$. (Observe that they all have the same $(u+1)$ -simple character).

We observe that just applying Corollary 1 in [6] to (2.45) (all tensor fields are acceptable and have the same simple character $\vec{\kappa}'_{simp}$),⁴² we obtain an equation:

$$\begin{aligned} & \sum_{h \in H_2^a} a_h Hitdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v + \\ & \sum_{u \in U} a_u Xdiv_{i_{\alpha+1}} C_g^{u,i_{\pi+1}\dots i_\alpha, i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v = \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v = 0, \end{aligned} \tag{2.46}$$

where the tensor fields indexed in U are acceptable (we are treating ∇Y as a factor $\nabla\phi_{u+1}$), have a simple character $\vec{\kappa}'_{simp}$ and each C^j is simply subsequent to $\vec{\kappa}'_{simp}$.

But then, our first claim follows almost immediately. We recall the operation $Erase_{\nabla Y}[\dots]$ from the Appendix in [3] which acts on the complete contractions in the above by erasing the factor ∇Y and the (derivative) index that it contracts against. Then, since (2.46) holds formally, we have that the tensor field required for Lemma 2.7 is:

$$\sum_{u \in U} a_u Erase_{\nabla Y}[C_g^{u,i_{\pi+1}\dots i_\alpha, i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)] \cdot \nabla_{i_\gamma} Y.$$

⁴¹Recall that i_γ is the free index that belongs to ∇Y .

⁴²Notice that by weight considerations, since we started out with no “bad” terms in the hypothesis of Lemma 2.1, there is no danger of falling under a “forbidden case” of that Corollary.

Proof of Lemma 2.7 in case B: We again distinguish two subcases: In subcase (i) there is some non-simple factor $S_* \nabla^{(\nu)} R_{ijkl}$ in $\vec{\kappa}_{simp}$ or a non-simple factor $\nabla^{(B)} \Omega_x$ contracting against two factors $\nabla \phi'_h$ in $\vec{\kappa}_{simp}$. In subcase (ii) there are no such factors.

In the subcase (i), we arbitrarily pick out one factor $S_* \nabla^{(\nu)} R_{ijkl}$ or $\nabla^{(B)} \Omega_x$ with the properties described above and call it the D -crucial factor. In this first subcase we will show our claim for the whole sublinear combination $\sum_{h \in H_2^a} \dots$ in one piece.

In the subcase (ii), we will introduce some notation: We will examine each factor $T = S_* \nabla^{(\nu)} R_{ijkl}$, $T = \nabla^{(B)} \Omega_x$ in each tensor field $C_g^{h, i_{\pi+1} \dots i_{\alpha}, i_{\alpha+1}}$ and define its “measure” as follows: If $T = S_* \nabla^{(\nu)} R_{ijkl}$ then its “measure” will stand for its total number of free indices *plus* $\frac{1}{2}$. If $T = \nabla^{(B)} \Omega_x$ then its “measure” will stand for its total number of free indices *plus* the number of factors $\nabla \phi_h$ against which it is contracting.

We divide the index set H_2^a into subsets according to the measure of any given factor. We denote by M the maximum measure among all factors among the tensor fields $C_g^{h, i_{\pi+1} \dots i_{\alpha}, i_{\alpha+1}}$, $h \in H_2^a$. We denote by $H_a^{2,*} \subset H_2^a$ the index set of the tensor fields which contain a factor of maximum measure. We will show the claim of Lemma 2.7 for the sublinear combination $\sum_{h \in H_a^{2,*}} \dots$. Clearly, if we can do this, then Lemma 2.7 will follow by induction.

We will prove Lemma 2.7 in the second subcase (which is the hardest). The proof in the first subcase follows by the same argument, only by disregarding any reference to M free indices belonging to a given factor etc.

Proof of Lemma 2.7 in case B for the sublinear combination $\sum_{h \in H_a^{2,}} \dots$:*

We will further divide $H_a^{2,*}$ into subsets, $H_a^{2,*,k}$, $k = 1, \dots, \sigma$, according to the factor of maximum measure: Firstly, we order the factors $S_* \nabla^{(\nu)} R_{ijkl}, \dots, \nabla^{(p)} \Omega_h$ in $\vec{\kappa}_{simp}$, and label them T_1, \dots, T_σ (observe each factor is well-defined in $\vec{\kappa}_{simp}$, because we are in case B). We then say that $h \in H_2^{a,*,1}$ if in $C_g^{u, i_{\pi+1} \dots i_{\alpha}}$ the factor T_1 has measure M . We say that $h \in H_2^{a,*,2}$ if in $C_g^{u, i_{\pi+1} \dots i_{\alpha}}$ the factor T_2 has measure M and T_1 has measure less than M , etc. We will then prove our claim for each of the index sets $h \in H_2^{a,*,k}$.⁴³ We arbitrarily pick a $k \leq K$ and show our claim for $\sum_{h \in H_a^{2,*,k}} \dots$.

For the purposes of this proof, we call the factor T_k the D -crucial factor.

Now, we pick out the subset $H_2^{b,k} \subset H_2^b$, which is defined by the rule: $h \in H_2^{b,k}$ if and only if ∇Y is contracting against the D -crucial factor T_k .

Now, for each $h \in H_2^a$ we denote by

$$Hitdiv_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$$

the sublinear combination in $Xdiv_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$ that arises when ∇_{i_γ} hits the D -crucial factor.⁴⁴ It then follows that:

⁴³Again we observe that if we can prove this then Lemma 2.7 in case B will follow by induction.

⁴⁴Recall that $i_\gamma = i_{\alpha+1}$ belongs to ∇Y by hypothesis.

$$\begin{aligned}
& \sum_{h \in H_2^a} a_h X \operatorname{div}_{i_{\pi+1}} \dots X \operatorname{div}_{i_\alpha} \operatorname{Hitdiv}_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \\
& + \sum_{h \in H_2^{b,k}} a_h X \operatorname{div}_{i_{\pi+1}} \dots X \operatorname{div}_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.47}$$

where each C_g^j has the factor ∇Y contracting against the D -crucial factor and is simply subsequent to $\vec{\kappa}_{simp}$.

We now denote the $(u+1)$ -simple character (the one defined by $\nabla \phi_1, \dots, \nabla \phi_{u+1} = \nabla Y$) of the tensor fields $\operatorname{Hitdiv}_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u)$ by $\vec{\kappa}'_{simp}$. (Observe that they all have the same $(u+1)$ -simple character).

We apply Corollary 1 in [6] to (2.47) (all tensor fields are acceptable and have the same simple character $\vec{\kappa}'_{simp}$) and then pick out the sublinear combination where there are M factors ∇v or $\nabla \phi_h$ or $\nabla \phi'_h$ contracting against T_k , we obtain an equation:

$$\begin{aligned}
& \sum_{h \in H_2^{a,*,k}} a_h \operatorname{Hitdiv}_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v + \\
& \sum_{u \in U} a_u X \operatorname{div}_{i_{\alpha+1}} C_g^{h, i_{\pi+1} \dots i_\alpha, i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v = \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v = 0,
\end{aligned} \tag{2.48}$$

where the tensor fields indexed in U are acceptable and have a simple character $\vec{\kappa}'_{simp}$ and each C^j is simply subsequent to $\vec{\kappa}'_{simp}$.

Now, observe that if $M \geq \frac{3}{2}$, we can apply the eraser to ∇Y (see the Appendix of [3]) and the index it is contracting against in the D -crucial factor and derive our conclusion as in case A.

On the other hand, in the remaining cases⁴⁵ the above argument cannot be directly applied. In those cases, we derive our claim as follows:

In the case $M = 1$ the D -crucial factor is of the form $\nabla^{(p)} \Omega_h$, then we cannot directly derive our claim by the above argument, because if for some tensor fields in U above we have ∇Y contracting according to the pattern $\nabla_i Y \nabla^{ij} \Omega_h \nabla_j \psi$ (where $\psi = v$ or $\psi = \phi_h$), then we will not obtain acceptable tensor fields after we apply the eraser. Therefore, if $M = 1$ and the D -crucial factor is of the form $\nabla^{(p)} \Omega_h$, we apply Lemma 4.6 in [6] to (2.48) (treating the factors ∇v as factors

⁴⁵Observe that the remaining cases are when $M = 0$, $M = \frac{1}{2}$, $M = 1$.

$\nabla\phi$)⁴⁶ to obtain a new equation in the form (2.48), where each tensor field in U has the factor ∇Y is contracting against a factor $\nabla^{(l)}\Omega_h$, $l \geq 3$.⁴⁷ Then, applying the eraser as explained, we derive our Lemma 2.7 in this case.

When $M = \frac{1}{2}$ or $M = 0$, then we first apply the inductive assumptions of Corollaries 3,2 in [6] (respectively) to (2.48),⁴⁸ in order to assume with no loss of generality that for each tensor field indexed in U there, the factor ∇Y is either contracting against a factor $\nabla^{(B)}\Omega_h$, $B \geq 3$ or a factor $S_*\nabla^{(\nu)}R_{ijkl}$, $\nu \geq 1$. Then the eraser can be applied and produces acceptable tensor fields. Hence, applying $Erase_{\nabla Y}$ to (2.48) we derive our claim. \square

Proof of Lemma 2.9:

We re-write the hypothesis of Lemma 2.3 (which is also the hypothesis of Lemma 2.9) in the following form:

$$\begin{aligned} & \sum_{h \in H_2} a_h X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} \{C_g^{h, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) - \\ & \operatorname{Switch}[C]_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} \\ & = \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u). \end{aligned} \tag{2.49}$$

Here the operation *Switch* interchanges the indices a and b in the two factors $\nabla_a \omega_1$, $\nabla_b \omega_2$.

Notational conventions: We have again denoted by $H_2^a \subset H_2$ the index set of those vector fields for which one of the free indices (say $i_{\alpha+1}$) belongs to a factor $\nabla \omega_1$ or $\nabla \omega_2$. With no loss of generality we assume that for each $h \in H_2^a$ $i_{\alpha+1}$ belongs to the factor $\nabla \omega_1$. We can clearly do this, due to the antisymmetry of the factors $\nabla \omega_1, \nabla \omega_2$.

We have defined $H_2^b = H_2 \setminus H_2^a$. For each $h \in H_2^b$ we denote by $T_{\omega_1}, T_{\omega_2}$ the factors against which $\nabla \omega_1, \nabla \omega_2$ are contracting. Also, for each $h \in H_2^a$ we will denote by T_{ω_2} the factor against which $\nabla \omega_2$ is contracting.⁴⁹

For each $h \in H_2$, we will call the factors $T_{\omega_1}, T_{\omega_2}$ against which $\nabla \omega_1$ or $\nabla \omega_2$ are contracting “problematic” in the following cases: If T_{ω_1} or T_{ω_2} is of the form $\nabla^{(m)}R_{ijkl}$ and $\nabla \omega_1$ or $\nabla \omega_2$ is contracting against an internal index. Alternatively, if T_{ω_1} or T_{ω_2} is of the form $S_*\nabla^{(\nu)}R_{ijkl}$ and the factor $\nabla \omega_1$ or $\nabla \omega_2$ is contracting one of the indices k or l .

We then define a few subsets of H_2^a, H_2^b :

⁴⁶Furthermore, we can observe that we do not fall under a “forbidden case” of Lemma 4.1 in [6], by weight considerations, and since the tensor fields in our Lemma assumption are not “bad”.

⁴⁷Note that the weight becomes less negative, hence Lemma 4.10 in [6] applies.

⁴⁸By our assumptions there will be a removable index in these cases. Hence our extra requirements of those Lemmas are fulfilled.

⁴⁹Note that the definition of $T_{\omega_1}, T_{\omega_2}$ depends on h ; however, to simplify notation we suppress the index h that should appear in $T_{\omega_1}, T_{\omega_2}$.

Definition 2.2 We define $H_{2,**}^b$ to stand for the index set of the tensor fields $C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}$'s for which $\nabla_{\omega_1}, \nabla_{\omega_2}$ are contracting against different factors and both $T_{\omega_1}, T_{\omega_2}$ are problematic.

We define $H_{2,*}^a \subset H_2^a$ to be the index set of the tensor fields $C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}$'s for which T_{ω_2} is problematic.

We define $H_{2,*}^b$ to stand for the index set of the tensor fields $C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}$'s for which either $T_{\omega_1} = T_{\omega_2}$ or $T_{\omega_1} \neq T_{\omega_2}$ and one of the factors $T_{\omega_1}, T_{\omega_2}$ is problematic.

Abusing notation, we will be using the symbols $\sum_{h \in H_{2,*}^b} \dots$ etc to denote generic linear combinations as explained above, when these symbols appear in the right hand sides of the equations below.

We then state three preparatory claims:

Firstly, we claim that we can write:

$$\begin{aligned}
& \sum_{h \in H_{2,**}^b} a_h X_+ \text{div}_{i_{\pi+1}} \dots X_+ \text{div}_{i_{\alpha+1}} \{C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} \\
& - \text{Switch}[C]_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \} = \\
& \sum_{h \in H_{2,*}^b} a_h X_+ \text{div}_{i_{\pi+1}} \dots X_+ \text{div}_{i_{\alpha+1}} \{C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} \\
& - \text{Switch}[C]_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \} \\
& + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.50}$$

where the linear combination $\sum_{h \in H_{2,*}^b} \dots$ on the RHS stands for a generic linear combination in the form described above. Observe that if we can show (2.50) then we may assume with no loss of generality that $H_{2,**}^b = \emptyset$ in our Lemma hypothesis.

Then, assuming that $H_{2,**}^b = \emptyset$ in our Lemma hypothesis we will show that there exists a linear combination of $(\alpha - \pi + 1)$ -tensor fields (indexed in X below) which are in the form (2.5) with a simple character $\vec{\kappa}_{simp}$ so that:

$$\begin{aligned}
& \sum_{h \in H_{2,*}^a} a_h \{ C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) - \\
& \text{Switch}[C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)] \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v - \\
& X_* \text{div}_{i_{\alpha+2}} \sum_{x \in X} a_x \{ C_g^{x, i_1 \dots i_{\alpha+2}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) - \\
& \text{Switch}[C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)] \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v \\
& + \sum_{h \in H_{2,*}^b} a_h \{ C_g^{h, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) - \\
& \text{Switch}[C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)] \nabla_{i_{\pi+1}} v \dots \nabla_{i_{\alpha+1}} v \\
& = \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u, v^{\alpha-\pi}).
\end{aligned} \tag{2.51}$$

We observe that if we can show the above, we may then assume that $H_{2,*}^a = \emptyset$ (and $H_{2,**}^b = \emptyset$) in the hypothesis of Lemma 2.9.

Finally, under the assumption that $H_{2,**}^b = H_{2,*}^a = \emptyset$ in our Lemma hypothesis, we will show that we can write:

$$\begin{aligned}
& \sum_{h \in H_{2,*}^b} a_h X_+ \text{div}_{i_{\pi+1}} \dots X_+ \text{div}_{i_{\alpha+1}} \{ C_g^{h, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) - \\
& \text{Switch}[C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)] \} = \\
& \sum_{h \in H_{2,OK}^b} a_h X_+ \text{div}_{i_{\pi+1}} \dots X_+ \text{div}_{i_{\alpha+1}} \{ C_g^{h, i_1 \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) - \\
& \text{Switch}[C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)] \} \\
& + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, [\omega_1, \omega_2], \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.52}$$

where the sublinear combination $\sum_{h \in H_{2,OK}^b} \dots$ on the right hand side stands for a generic linear combination of acceptable tensor fields in the form (2.5) with simple character \vec{k}_{simp} , with no free indices in the factors $\nabla \omega_1, \nabla \omega_2$ and where the factors $T_{\omega_1}, T_{\omega_2}$ are not problematic. Therefore, if we can show the above equations, we are reduced to showing Lemma 2.9 under the additional assumptions that $H_{a,*}^2 = H_{b,**}^2 = H_{b,*}^2 = \emptyset$.

(Sketch of the) Proof of (2.50), (2.51), (2.52): (2.50) follows by re-iterating the proof of the first claim of Lemma 4.10 in [6].⁵⁰ (2.51) follows by re-iterating

⁵⁰By the additional restrictions imposed on the assumption of Lemma 2.3 there is no danger of falling under a “forbidden case” of Corollary 1 in [6].

the proof of the first claim of Lemma 4.10 in [6], but rather than applying Corollary 1 [6] in that proof, we now apply Lemma 2.7 (which we have shown).⁵¹ Finally, the claim of (2.52) for the sublinear combination in $H_{2,*}^b$ where $T_{\omega_1} \neq T_{\omega_2}$ follows by applying Lemma 2.5.⁵² We can then show that the remaining sublinear combination in $\sum_{h \in H_{2,*}^b} \dots$ must vanish separately (modulo a linear combination $\sum_{j \in J} \dots$) by just picking out the sublinear combination in the hypothesis of Lemma 2.10 where both factors $\nabla\omega_1, \nabla\omega_2$ are contracting against the same factor. \square

Now, under these additional assumptions that $H_{a,*}^2 = H_{b,**}^2 = H_{b,*}^2 = \emptyset$, we will show our claim by distinguishing two cases: In case A there is a factor $\nabla^{(m)}R_{ijkl}$ in $\vec{\kappa}_{simp}$; in case B there is no such factor. An important note: We may now use Lemma 2.7, which we have proven earlier in this section.

Proof of Lemma 2.9 in case A.

We define the (set of) C-crucial factor(s) (which will necessarily be of the form $\nabla^{(m)}R_{ijkl}$) as in the setting of Lemma 2.7. Firstly a mini-claim which only applies to the case where the C-crucial factor is unique:

Mini-claim, when the C-crucial factor is unique: We then consider the tensor fields $C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}$, $h \in H_2^a$ for which $\nabla\omega_2$ is contracting against the C-crucial factor. Notice that by our hypothesis that $H_{a,*}^2 = \emptyset$, we will have that $\nabla\omega_2$ is contracting against a derivative index in the C-crucial factor. Denote by $H_2^{a,+} \subset H_2^a$ the index set of these tensor fields.

We observe that for each $h \in H_2^{a,+}$ we can now construct a tensor field by erasing the index in the factor $\nabla^{(m)}R_{ijkl}$ that contracts against the factor $\nabla\omega_2$ and making the index in $\nabla\omega_2$ into a free index i_{β} . We denote this tensor field by $C_g^{h, i_{\pi+1} \dots i_{\alpha+1} i_{\beta}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)$. By the analogous operation we obtain a tensor field $Switch[C_g^{h, i_{\pi+1} \dots i_{\alpha+1} i_{\beta}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)]$.

It follows that in the case where the C-crucial factor is unique, for each $h \in H_2^{a,+}$:

⁵¹Observe that the assumption that Lemma 2.3 does not include “forbidden cases” ensures that we will not need to apply Lemma 2.7 in a “forbidden case”.

⁵²In this case there will be a factor $\nabla\omega_1$ or $\nabla\omega_2$ contracting against a non-special index; therefore there is no danger of falling under a “forbidden” case of Lemma 2.7.

$$\begin{aligned}
& X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} \{C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\
& - \operatorname{Switch}[C]_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} = \\
& X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} X_* \operatorname{div}_{i_{\beta}} \{C_g^{h, i_{\pi+1} \dots i_{\alpha+1} i_{\beta}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\
& - \operatorname{Switch}[C]_g^{h, i_{\pi+1} \dots i_{\alpha+1} i_{\beta}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} + \\
& \sum_{r \in R} a_r X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} \{C_g^{r, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\
& - \operatorname{Switch}[C]^{r, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.53}$$

where each tensor field $C_g^{r, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)$ has the factor $\nabla \omega_2$ contracting against some factor other than the C-crucial factor.

But we observe that:

$$\begin{aligned}
& X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} X_* \operatorname{div}_{i_{\beta}} \{C_g^{h, i_{\pi+1} \dots i_{\alpha+1} i_{\beta}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\
& - \operatorname{Switch}[C]_g^{h, i_{\pi+1} \dots i_{\alpha+1} i_{\beta}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\} = 0.
\end{aligned} \tag{2.54}$$

Therefore, in the case $Set \neq \emptyset$ or $Set = \emptyset$ and $\sigma_1 = 1$, we have now reduced Lemma 2.9 to the case where $H_2^{a,+} = \emptyset$.

Now, (under the assumption that $H_2^{a,+} = \emptyset$ when the C-crucial factor is unique) we consider the sublinear combination *Special* in the equation hypothesis of Lemma 2.9 that consists of complete contractions with $\nabla \omega_1$ contracting against the C-crucial factor while the factor $\nabla \omega_2$ is contracting against some other factor. (If $Set = \emptyset$ and $\sigma_1 > 1$ *Special* stands for the sublinear combination where $\nabla \omega_1$ is contracting against a generic C-crucial factor and $\nabla \omega_2$ is contracting against some other factor). In particular, for each $h \in H_2^a$, since $H_2^{a,+} = \emptyset$ we see that the sublinear combination in

$$\begin{aligned}
& \sum_{h \in H_2^a} a_h X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha+1}} \{C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\
& - \operatorname{Switch}[C]_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)\}
\end{aligned} \tag{2.55}$$

that belongs to *Special* is precisely:

$$\sum_{h \in H_2^a} a_h X_* \operatorname{div}_{i_{\pi+1}} \dots X_* \operatorname{div}_{i_{\alpha}} \operatorname{Hit} \operatorname{div}_{i_{\alpha+1}} C_g^{h, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u);$$

(in the case $Set = \emptyset$ and $\sigma_1 > 1$ $Hitdiv_{i_{\alpha+1}}$ just means that ∇_{i_γ} can hit any factor $\nabla^{(m)}R_{ijkl}$ that is not contracting against $\nabla\omega_2$; recall that in the other cases it means that it must hit the unique C-crucial factor).

We also consider the tensor fields $C^{h,i_{\pi+1}\dots i_{\alpha+1}}$, $Switch[C]^{h,i_{\pi+1}\dots i_{\alpha+1}}$, $h \in H_2^b$ for which $\nabla\omega_1$ is contracting against the C-crucial factor and $\nabla\omega_2$ is not (or, if there are multiple C-crucial factors, where $\nabla\omega_1, \nabla\omega_2$ are contracting against different C-crucial factors). For this proof, we index all those tensor fields in $H_2^{b,\Psi}$ and we will denote them by $C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}$.

Thus we derive an equation:

$$\begin{aligned} & \sum_{h \in H_2^a} a_h X_* div_{i_{\pi+1}} \dots X_* div_{i_\alpha} Hitdiv_{i_{\alpha+1}} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\ & + \sum_{h \in H_2^{b,\Psi}} a_h X_* div_{i_{\pi+1}} \dots X_* div_{i_\alpha} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) = \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u). \end{aligned} \tag{2.56}$$

We group up the vector fields on the left hand side according to their *weak* $(u+2)$ -characters⁵³ (defined by $\nabla\phi_1, \dots, \nabla\phi_u, \nabla\omega_1, \nabla\omega_2$). (Recall that we started off with complete contractions with the same u -simple characters-so the only new information that we are taking into account is what *type* of factor is $\nabla\omega_2$ contracting against). We consider the set of weak simple characters that we have obtained. We denote the set by $\{\vec{\kappa}_1, \dots, \vec{\kappa}_B\}$, and we respectively have the index sets $H_2^{a,\vec{\kappa}_f}$ and $H_2^{b,\vec{\kappa}_f}$.

We will show our Lemma 2.9 by replacing the index set H_2^a by any $H_2^{a,\vec{\kappa}_f}$, $f \leq B$.

It follows that for each $f \leq B$:

$$\begin{aligned} & \sum_{h \in H_2^{a,\vec{\kappa}_f}} a_h X_* div_{i_{\pi+1}} \dots X_* div_{i_\alpha} Hitdiv_{i_{\alpha+1}} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}} \\ & (\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) + \\ & \sum_{h \in H_2^{b,\vec{\kappa}_f}} a_h X_* div_{i_{\pi+1}} \dots X_* div_{i_\alpha} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) = \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u), \end{aligned} \tag{2.57}$$

where the complete contractions C_g^j have a u -simple character that is subsequent to $\vec{\kappa}_{simp}$. We will show our claim for each of the index sets $H_2^{b,\vec{\kappa}_f}$ separately.

⁵³See [6] for a definition of this notion.

Now, we treat the factors $\nabla\omega_1, \nabla\omega_2$ in the above as factors $\nabla\phi_{u+1}, \nabla\phi_{u+2}$. We see that since $H_{2,**}^b = H_{b,*}^2 = H_{a,*}^2 = \emptyset$, all the tensor fields in the above have the same $(u+2)$ -simple character.

Our claim (Lemma 2.9) for the index set $H_2^{a,\vec{\kappa}_f}$ then follows: Firstly, apply the operator $Eraser_{\nabla\omega_1}[\dots]$ to (2.57).⁵⁴ We are then left with tensor fields (denote them by

$$C_g^{h,i_{\pi+1}\dots i_\alpha}(\Omega_1, \dots, \Omega_p, \omega_2, \phi_1, \dots, \phi_u), h \in H_2^{a,\vec{\kappa}_f},$$

$$C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_2, \phi_1, \dots, \phi_u), h \in H_2^{b,\vec{\kappa}_f},$$

respectively) with the same $(u+1)$ -simple character say $\vec{\kappa}_{simp,f}$. We can then apply Corollary 1 from [6] (since we have weight $-n+2k$, $k > 0$ by virtue of the eraser—notice that by weight considerations, since we started out with no “bad” tensor fields, there is no danger of falling under a “fobidden case”). to derive that there is a linear combination of acceptable α -tensor fields indexed in V below, with $(u+1)$ -simple character $\vec{\kappa}_{simp,f}$ so that:

$$\begin{aligned} & \sum_{h \in H_2^{a,\vec{\kappa}_f}} a_h C_g^{h,i_{\pi+1}\dots i_\alpha}(\Omega_1, \dots, \Omega_p, \omega_2, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v - \\ & \sum_{v \in V} a_v X_* \text{div}_{i_{\alpha+1}} C_g^{v,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_2, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v = \\ & \sum_{j \in J} a_j C_g^{j,i_{\pi+1}\dots i_\alpha}(\Omega_1, \dots, \Omega_p, \omega_2, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v, \end{aligned} \tag{2.58}$$

where each complete contraction indexed in J is $(u+1)$ -subsequent to $\vec{\kappa}_{simp,f}$. In this setting $X_* \text{div}_i$ just means that in addition to the restrictions imposed on $X \text{div}_i$ we are not allowed to hit the factor $\nabla\omega_2$.

Then, if we multiply the above equation by an expression $\nabla_i \omega_1 \nabla^i v$ and then anti-symmetrize the indices a, b in the factors $\nabla_a \omega_1, \nabla_b \omega_2$ and finally make all ∇v s into $X_+ \text{div}s$, we derive our claim. \square

Proof of Lemma 2.9 in case B (when $\sigma_1 = 0$).

Our proof follows the same pattern as the proof of Lemma 2.7 in case B.

We again define the “measure” of each factor in each tensor field $C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}$ as in the proof of case B in Lemma 2.7. Again, let M stand for the maximum measure among all factors in all tensor fields $C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}$, $h \in H_2^a$. We denote by $H_2^{a,M} \subset H_2^a$ the index set of the tensor fields for which some factor has measure M .

We will further divide $H_2^{a,M}$ into subsets, $H_2^{a,M,k}$, $k = 1, \dots, \sigma$, according to the factor which has measure M : Firstly, we order the factors $S_* \nabla^{(\nu)} R_{ijkl}, \dots \nabla^{(p)} \Omega_h$

⁵⁴See the relevant Lemma in the Appendix of [3].

in $\vec{\kappa}_{simp}$, and label them T_1, \dots, T_σ (observe each factor is well-defined in $\vec{\kappa}_{simp}$, because we are in case B). We then say that $h \in H_2^{a,M,1}$ if in $C_g^{h,i_{\pi+1}\dots i_\alpha}$ T_1 has measure M . We say that $h \in H_2^{a,M,2}$ if in $C_g^{h,i_{\pi+1}\dots i_\alpha}$ T_2 has measure M and T_1 has measure less than M , etc. We will then prove our claim for each of the index sets $h \in H_2^{a,M,k}$.⁵⁵ We arbitrarily pick a $k \leq \sigma$ and show our claim for $\sum_{h \in H_2^{a,M,k}} \dots$

For the purposes of this proof, we call the factor T_k the D -crucial factor (in this setting the D -crucial factor is unique).

Now, we pick out the subset $H_2^{b,k,+} \subset H_2^b$, which is defined by the rule: $h \in H_2^{b,k,+}$ if and only if $\nabla\omega_1$ is contracting against the D -crucial factor T_k . We also pick out the subset $H_2^{b,k,-} \subset H_2^b$, which is defined by the rule: $h \in H_2^{b,k,-}$ if and only if $\nabla\omega_2$ is contracting against the D -crucial factor T_k . Finally, we define $H_2^{a,-} \subset H_2^a$, $H_2^{a,+} \subset H_2^a$ to stand for the index set of tensor fields for which $\nabla\omega_2$ contracts against the D -crucial factor.

Now, for each $h \in H_2^a$ we denote by

$$Hitdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)$$

the sublinear combination in $Xdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)$ that arises when ∇_{i_γ} hits the D -crucial factor. It then follows that:

$$\begin{aligned} & \sum_{h \in H_2^a} a_h X_* div_{i_{\pi+1}} \dots X_* div_{i_\alpha} Hitdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\ & - \sum_{h \in H_2^{a,-}} a_h X_* div_{i_{\pi+1}} \dots X_* div_{i_{\alpha+1}} Switch[C]_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\ & + \sum_{h \in H_2^{b,k,+}} a_h X div_{i_{\pi+1}} \dots X div_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\ & - \sum_{h \in H_2^{b,k,-}} a_h X div_{i_{\pi+1}} \dots X div_{i_\gamma} Switch[C]_g^{h,i_{\pi+1}\dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \\ & + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u), \end{aligned} \tag{2.59}$$

where each C_g^j has the factor $\nabla\omega_1$ contracting against the D -crucial factor and is simply subsequent to $\vec{\kappa}_{simp}$.

We now denote the $(u+1)$ -simple character (the one defined by $\nabla\phi_1, \dots, \nabla\omega_1$) of the tensor fields $Hitdiv_{i_\gamma} C_g^{h,i_{\pi+1}\dots i_\alpha, i_\gamma}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u)$ by $\vec{\kappa}'_{simp}$. (Observe that they all have the same $(u+1)$ -simple character).

We observe that just applying Lemma 2.1 to (2.59) (all tensor fields are acceptable and have the same simple character $\vec{\kappa}'_{simp}$ —we treat $\nabla\omega_1$ as a factor

⁵⁵Again we observe that if we can prove this then Lemma 2.9 in case B will follow by induction.

$\nabla\phi_{u+1}$ and the factor $\nabla\omega_2$ as a factor ∇Y) and we then pick out the sublinear combination where there are M factors ∇v contracting against T_k , we obtain an equation:

$$\begin{aligned} & \sum_{h \in H_2^{\alpha, *, k}} a_h \text{Hitdiv}_{i_\gamma} C_g^{h, i_{\pi+1} \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v + \\ & \sum_{x \in X} a_x X \text{div}_{i_{\alpha+1}} C_g^{x, i_{\pi+1} \dots i_{\alpha+1}}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v + \\ & \sum_{j \in J} a_j C_g^{j, i_{\pi+1} \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \omega_1, \omega_2, \phi_1, \dots, \phi_u) \nabla_{i_{\pi+1}} v \dots \nabla_{i_\alpha} v = 0, \end{aligned} \tag{2.60}$$

where the tensor fields indexed in X are acceptable and have a $(u+1)$ -simple character $\bar{\kappa}'_{simp}$ and each C^j is simply subsequent to $\bar{\kappa}'_{simp}$.

Now, observe that if $M \geq \frac{3}{2}$ then we can apply the Eraser (from the Appendix in [3]) to $\nabla\omega_1$ and the index it is contracting against in the D -crucial factor and derive our conclusion as in case A.

The remaining cases are when $M = 1$, $M = \frac{1}{2}$ and $M = 0$. The first one is easier, so we proceed to show our claim in that case. The two subcases $M = \frac{1}{2}$, $M = 0$ will be discussed in the next subsection.

In the case $M = 1$, i.e. the D -crucial factor is of the form $\nabla^{(p)}\Omega_h$, then we cannot derive our claim, because if for some tensor fields in X above we have $\nabla\omega_1$ contracting according to the pattern: $\nabla_i\omega_1\nabla^{ij}\Omega_h\nabla_j\psi$, where $\psi = v$ or $\psi = \phi_h$. Therefore, in this setting, we first apply the eraser twice to remove the expression $\nabla_{ij}^{(2)}\Omega_h\nabla^i\psi\nabla^j\omega_1$ and then apply Corollary 2 from [6]⁵⁶ to (2.60) (observe that (2.60) now falls under the inductive assumption of Lemma 4.6 in [6] since we have lowered the weight⁵⁷ to obtain a new equation in the form (2.60), where each tensor field in X has the factor $\nabla\omega_1$ contracting against a factor $\nabla^{(l)}\Omega_h$, $l \geq 3$. Then, applying the eraser as explained, we derive our Lemma 2.9 in this case.

The cases $M = \frac{1}{2}$, $M = 0$: Notice that in this case we must have $\alpha = \pi$, by virtue of the the definition of maximal “measure” above. We will then prove our claim by proving a more general claim by induction, in the next subsection. \square

2.4 The remaining cases of Lemma 2.9.

We prove our claim in this case by an induction. In order to give a detailed proof, we will re-state our Lemma hypothesis in this case (with a slight change

⁵⁶Recall that we showed in [6] that this is a Corollary of Lemma 4.6 in [6], which we have now shown.

⁵⁷There is no danger of falling under a “forbidden case” of Lemma 2.1 by weight considerations since we are assuming that none of the tensor fields of minimum rank in the assumption of Lemma 2.3 are “bad”.

of notation):

The hypothesis of the remaining cases of Lemma 2.9: We are assuming an equation:

$$\begin{aligned}
& \sum_{x \in X_a} a_x X_* \text{div}_{i_1} C_g^{x, i_1}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, [\omega_1, \omega_2]) + \\
& \sum_{x \in X_b} a_x X_* \text{div}_{i_1} C_g^{x, i_1}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, [\omega_1, \omega_2]) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) = 0,
\end{aligned} \tag{2.61}$$

which holds modulo complete contractions of length $\geq \sigma + u + 3$ ($\sigma \geq 3$ —here σ stands for $u + p$ —see the next equation). We denote the weight of the complete contractions in the above by $-K$. The tensor fields in the above equation are each in the form:

$$\begin{aligned}
& p\text{contr}(S_* \nabla^{(\nu_1)} R_{x_1 j k l} \otimes \dots \otimes S_* \nabla^{(\nu_u)} R_{x_z j' k' l'} \otimes \\
& \nabla^{(a_1)} \Omega_1 \otimes \dots \otimes \nabla^{(a_p)} \Omega_p \otimes [\nabla \omega_1 \otimes \nabla \omega_2] \\
& \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \dots \otimes \nabla^{x_u} \tilde{\phi}_z).
\end{aligned} \tag{2.62}$$

We recall that the u -simple character of the above has been denoted by $\vec{\kappa}_{simp}$. Recall that we are now assuming that all the factors $\nabla^{(a_i)} \Omega_x$ in $\vec{\kappa}_{simp}$ are acceptable.⁵⁸ The complete contractions indexed in J in (2.61) are simply subsequent to $\vec{\kappa}_{simp}$. We also recall that $X_* \text{div}_i$ stands for the sublinear combination in $X \text{div}_i$ where ∇_i is not allowed to hit either of the factors $\nabla \omega_1, \nabla \omega_2$.

We recall that the tensor fields indexed in X_a have the free index i_1 belonging to the factor $\nabla \omega_1$. The tensor fields indexed in X_b have the free index i_1 *not* belonging to any of the factors $\nabla \omega_1, \nabla \omega_2$.

We recall the key assumption that for each of the tensor fields indexed in X_a , there is at least one removable index in each tensor field $C_g^{x, i_1}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, [\omega_1, \omega_2])$, $x \in X_a$.⁵⁹

In order to complete our proof of Lemma 2.9, we will show that we can write:

⁵⁸meaning that each $a_i \geq 2$.

⁵⁹Recall the definition of a “removable” index from Definition 2.1.

$$\begin{aligned}
& \sum_{x \in X_a} a_x C_g^{x, i_1}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, [\omega_1, \omega_2]) \nabla_{i_1} v = \\
& \sum_{x \in X'} a_x X_* \operatorname{div}_{i_2} \dots X_* \operatorname{div}_{i_a} C_g^{x, i_1 \dots i_a}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, [\omega_1, \omega_2]) \nabla_{i_1} v + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u),
\end{aligned} \tag{2.63}$$

where the tensor fields indexed in X' are acceptable in the form (2.62), each with rank $a \geq 2$. Note that this will imply the remaining cases of Lemma 2.9, completing the proof of Lemma 2.3.

We recall that we are proving this claim when the assumption (2.61) formally falls under our inductive assumption of Proposition 1.1 (if we formally treat $\nabla\omega_1, \nabla\omega_2$ as factors $\nabla\phi_{z+1}, \nabla\phi_{z+2}$).

We will prove (2.63) by inductively proving a more general statement. The more general statement is as follows:

The general statement:

Assumptions: We consider vector fields $C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau)$, $C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau)$ in the following forms, respectively:

$$\begin{aligned}
& p\operatorname{contr}(S_* \nabla^{(\nu_1)} R_{x_1 j k l} \otimes \dots \otimes S_* \nabla^{(\nu_v)} R_{x_v j' k' l'} \otimes \\
& \nabla^{(a_1)} \Omega_1 \otimes \dots \otimes \nabla^{(a_b)} \Omega_b \otimes \nabla Y \otimes \nabla \psi_1 \otimes \dots \otimes \nabla \psi_\tau \\
& \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \dots \otimes \nabla^{x_v} \tilde{\phi}_v),
\end{aligned} \tag{2.64}$$

$$\begin{aligned}
& p\operatorname{contr}(S_* \nabla^{(\nu_1)} R_{x_1 j k l} \otimes \dots \otimes S_* \nabla^{(\nu_v)} R_{x_v j' k' l'} \otimes \\
& \nabla^{(a_1)} \Omega_1 \otimes \dots \otimes \nabla^{(a_b)} \Omega_b \otimes [\nabla \chi_1 \otimes \nabla \chi_2] \otimes \nabla \psi_1 \otimes \dots \otimes \nabla \psi_\tau \\
& \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \dots \otimes \nabla^{x_v} \tilde{\phi}_v),
\end{aligned} \tag{2.65}$$

for which the weight is $-W + 1, W \leq K$. We also assume $v + b \geq 2$. *Note:* the bracket $[\dots]$ stands for the anti-symmetrization of the indices a, b in the expression $\nabla_a \omega_1 \nabla_b \omega_2$.

We assume (respectively) the equations:

$$\begin{aligned}
& \sum_{\zeta \in Z_a} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in \bar{Z}_a} a_\zeta X_* \text{div}_{i_1} \dots X_* \text{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = 0,
\end{aligned} \tag{2.66}$$

$$\begin{aligned}
& \sum_{\zeta \in Z_a} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in \bar{Z}_a} a_\zeta X_* \text{div}_{i_1} \dots X_* \text{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = 0,
\end{aligned} \tag{2.67}$$

which holds modulo complete contractions of length $\geq v + b + \tau + 3$.

The tensor fields indexed in Z_a are assumed to have a free index in one of the factors $\nabla Y, \nabla \psi_1, \dots, \nabla \psi_\tau$, or one of the factors $\nabla \chi_1, \nabla \chi_2, \nabla \psi_1, \dots, \nabla \psi_\tau$, respectively. The tensor fields indexed in \bar{Z}_a have rank $\gamma \geq 2$ and all their free indices belong to the factors $\nabla Y, \nabla \psi_1, \dots, \nabla \psi_\tau$, or the factors $\nabla \chi_1, \nabla \chi_2, \nabla \psi_1, \dots, \nabla \psi_\tau$, respectively. The tensor fields indexed in Z_b have the property that i_1 does not belong to any of the factors $\nabla Y, \nabla \psi_1, \dots, \nabla \psi_\tau, \nabla \chi_1, \nabla \chi_2, \nabla \psi_1, \dots, \nabla \psi_\tau$, respectively. We furthermore assume that for the tensor fields indexed in $Z_a \cup Z_b \cup \bar{Z}_a$, none of the factors $\nabla \psi_1, \dots, \nabla \psi_\tau$ are contracting against a special index in any factor $S_* \nabla^{(\nu)} R_{ijkl}$ and none of them are contracting against the rightmost index in each $\nabla^{(a_h)} \Omega_h$ (we will refer to this property as the @-property). We assume that $v + b \geq 2$, and furthermore if $v + b = 2$ then for each $\zeta \in Z_a \cup Z_b$, the factor(s) ∇Y (or $\nabla \chi_1, \nabla \chi_2$) are also not contracting against a special index in any $S_* \nabla^{(\nu)} R_{ijkl}$ and are not contracting against the rightmost index in any $\nabla^{(a_h)} \Omega_h$. Finally (and importantly) we assume that for the tensor fields indexed in Z_a , there is at least one removable index in each C_g^{ζ, i_1} . (In this setting, for a tensor field indexed in Z_a , a “removable” index is either a non-special index in a factor $S_* \nabla^{(\nu)} R_{ijkl}$, with $\nu > 0$ or an index in a factor $\nabla^{(B)} \Omega_h$, $B \geq 3$).

Convention: In this subsection only, for tensor fields in the forms (2.66), (2.67) we say then an index is *special* if it is one of the indices k, l in a fac-

tor $S_* \nabla^{(\nu)} R_{ijkl}$ (this is the usual convention), or if it is an index in a factor $\nabla_{r_1 \dots r_B}^{(B)} \Omega_h$ for which all the other indices are contracting against factors $\nabla \psi_1, \dots, \nabla \psi_\tau$.

All tensor fields in (2.66), (2.67) have a given ν -simple character $\bar{\kappa}_{simp}$. The complete contractions indexed in J are assumed to have a weak ν -character $Weak(\bar{\kappa}_{simp})$ and to be simply subsequent to $\bar{\kappa}_{simp}$. Here $X_* div_i$ stands for the sublinear combination in $X div_i$ where ∇_i is not allowed to hit any of the factors $\nabla Y, \nabla \psi_1, \dots, \nabla \psi_\tau$ or $\nabla \chi_1, \nabla \chi_2, \nabla \psi_1, \dots, \nabla \psi_\tau$, respectively.

The Claims of the general statement: We claim that under the assumption (2.67), there exists a linear combination of acceptable 2-tensor fields in the form (2.64), (2.65) respectively (indexed in W below), for which the @-property is satisfied, so that (respectively):

$$\begin{aligned} & \sum_{\zeta \in Z_a} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_\nu, Y, \psi_1, \dots, \psi_\tau) \nabla_{i_1} v - \\ & \sum_{w \in W} a_w X_* div_{i_2} C_g^{w, i_1 i_2}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_\nu, Y, \psi_1, \dots, \psi_\tau) \nabla_{i_1} v + \quad (2.68) \\ & \sum_{j \in J} a_j C_g^{j, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_\nu, Y, \psi_1, \dots, \psi_\tau) \nabla_{i_1} v = 0, \end{aligned}$$

$$\begin{aligned} & \sum_{\zeta \in Z_a} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_\nu, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v + \\ & \sum_{w \in W} a_w X_* div_{i_2} C_g^{w, i_1 i_2}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_\nu, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v + \\ & \sum_{j \in J} a_j C_g^{j, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_\nu, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v = 0. \quad (2.69) \end{aligned}$$

We observe that when $\tau = 0$ and $\nu + b \geq 3$, (2.69) coincides with (2.63).⁶⁰ Therefore, if we can prove this general statement, we will have shown Lemma 2.9 in full generality, thus also completing the proof of Lemma 2.3.

We also have a further claim, when we assume (2.66), (2.67) with $\nu + b = 2$. In that case, we also claim that we can write:

⁶⁰Also, the assumption of existence of a non-removable index coincides with the corresponding assumption of Lemma 2.3.

$$\begin{aligned}
& X_+ \operatorname{div}_{i_1} \sum_{\zeta \in Z_a \cup Z_b \cup \bar{Z}_a} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.70}$$

$$\begin{aligned}
& X_+ \operatorname{div}_{i_1} \sum_{\zeta \in Z_a \cup \bar{Z}_a \cup Z_b} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.71}$$

where the tensor fields indexed in Q are in the same form as (2.64) or (2.65) respectively, but have a factor (expression) $\nabla^{(2)}Y$ or $\nabla_{a[i}^{(2)}\omega_1 \nabla_{j]}\omega_2$, respectively, and satisfy all the other properties of the tensor fields in Z_a .

Consequence of (2.68), (2.69) when $v + b \geq 3$: We here codify a conclusion one can derive from (2.68), (2.69). This implication will be useful further down in this subsection. We see that by making the factors ∇v into $X_* \operatorname{div}$'s in (2.66), (2.67) and replacing into (2.68), (2.69), we obtain new equations:

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = 0,
\end{aligned} \tag{2.72}$$

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = 0,
\end{aligned} \tag{2.73}$$

where here the tensor fields indexed in Z'_a are like the tensor fields indexed in Z_a in (2.66), (2.67) but have the additional feature that no free index belongs

to the factor $\nabla\psi_1$ (and all the other assumptions of equations (2.66), (2.67) continue to hold).

We then claim that we can derive new equations:

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.74}$$

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.75}$$

where here $X_+ \text{div}_i$ stands for the sublinear combination in $X \text{div}_i$ where ∇_i is allowed to hit the factor ∇Y or $\nabla \chi_1$ (respectively), but not the factors $\nabla \psi_1, \dots, \nabla \phi_\tau, (\nabla \chi_2)$. Furthermore, the linear combinations indexed in Q stand for generic linear combinations of vector fields in the form (2.64) or (2.65), only with the expressions ∇Y or $\nabla_{[a]\omega_1} \nabla_{[b]}\omega_2$ replaced by expressions $\nabla^{(2)} Y$, $\nabla_{c[a]\omega_1}^{(2)} \nabla_{[b]}\omega_2$.

Proof that (2.74), (2.75) follow from (2.68), (2.69): We prove the above by an induction. We will firstly subdivide Z'_a, Z_b into subsets as follows: $\zeta \in Z'_{a, \textcircled{a}}$ or $\zeta \in Z_{b, \textcircled{a}}$ if the factor ∇Y (or one of the factors $\nabla \chi_1, \nabla \chi_2$) is contracting against a special index in the same factor against which $\nabla \psi_1$ is contracting.

Now, if $Z'_{a, \textcircled{a}} \cup Z_{b, \textcircled{a}} \neq \emptyset$ our inductive statement will be the following:

We inductively assume that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z'_{a, \otimes}} a_\zeta X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_{b, \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^k} a_t X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_k} C_g^{t, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z'_{a, N \circ \otimes}} a_\zeta X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.76}$$

and

$$\begin{aligned}
& \sum_{\zeta \in Z'_{a, \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_{b, \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^k} a_t X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_k} C_g^{t, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z'_{a, N \circ \otimes}} a_\zeta X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.77}$$

where the tensor fields indexed in T^k have all the properties of the tensor fields indexed in $Z'_{a, \otimes}$ (in particular the index in $\nabla \psi_1$ is not free) and in addition have rank k . The tensor fields indexed in $Z'_{a, N \circ \otimes}$ in the RHS have all the regular features of the terms indexed in Z'_a (in particular rank $\gamma \geq 1$ and the factor $\nabla \psi_1$ does not contain a free index) and in addition none of the factors ∇Y (or $\nabla \chi_1, \nabla \chi_2$) are contracting against a special index.

Our inductive claim is that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^{k+1}} a_t X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_{k+1}} C_g^{t, i_1 \dots i_{k+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z'_{a, N \circ \mathbb{Q}}} a_\zeta X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.78}$$

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^{k+1}} a_t X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_k} C_g^{t, i_1 \dots i_{k+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z'_{a, N \circ \mathbb{Q}}} a_\zeta X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_\gamma} C_g^{\zeta, i_1 \dots i_\gamma}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \\
& + \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = 0.
\end{aligned} \tag{2.79}$$

We will derive (2.78), (2.79) momentarily. For now, we observe that by iterative repetition of the above inductive step we are reduced to showing (2.74), (2.75) under the additional assumption that $Z'_{a, \mathbb{Q}} = \emptyset$.

Under that assumption, we denote by $Z_{b, \mathbb{Q}} \subset Z_b$ the index set of vector fields for which the factor ∇Y (or one of the factors $\nabla \chi_1, \nabla \chi_2$) is contracting against a special index. We will then assume that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z_{b, \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{t \in V^k} a_t X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_k} C_g^{t, i_1 \dots i_k}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_{b, N \circ \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.80}$$

$$\begin{aligned}
& \sum_{\zeta \in Z_{b, \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{t \in V^k} a_t X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_k} C_g^{t, i_1 \dots i_k}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_{b, N \circ \otimes}} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.81}$$

where the tensor fields indexed in V^k have all the features of the tensor fields indexed in $Z_{b, \otimes}$ but in addition have all the k free indices *not* belonging to factors $\nabla\psi_1, \dots, \nabla\psi_\tau$. The tensor fields indexed in $Z_{b, N \circ \otimes}$ have all the regular features of the tensor fields in Z_b and in addition have the factor ∇Y (or the factors $\nabla\chi_1, \nabla\chi_2$) *not* contracting against special indices. The terms indexed in Q are as required in the RHS of (2.74), (2.75) (which are the equations that we are proving).

We will then show that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z_{b, \mathbb{Q}}} a_{\zeta} X_{+} \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_{\tau}) = \\
& \sum_{t \in V^{k+1}} a_t X_{+} \operatorname{div}_{i_1} \dots X_{+} \operatorname{div}_{i_{k+1}} C_g^{t, i_1 \dots i_{k+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_{\tau}) + \\
& \sum_{\zeta \in Z_{b, N \circ \mathbb{Q}}} a_{\zeta} X_{+} \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_{\tau}) + \\
& \sum_{q \in Q} a_q X_{+} \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_{\tau}) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_{\tau}),
\end{aligned} \tag{2.82}$$

$$\begin{aligned}
& \sum_{\zeta \in Z_{b, \mathbb{Q}}} a_{\zeta} X_{+} \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_{\tau}) = \\
& \sum_{t \in V^{k+1}} a_t X_{+} \operatorname{div}_{i_1} \dots X_{+} \operatorname{div}_{i_{k+1}} C_g^{t, i_1 \dots i_{k+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_{\tau}) \\
& + \sum_{\zeta \in Z_{b, N \circ \mathbb{Q}}} a_{\zeta} X_{+} \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_{\tau}) + \\
& \sum_{q \in Q} a_q X_{+} \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_{\tau}) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_{\tau}).
\end{aligned} \tag{2.83}$$

(Here the tensor fields indexed in V^{k+1} have all the features described above and moreover have rank $k+1$).

Thus, by iterative repetition of this step we are reduced to showing our claim under the additional assumption that $Z'_{a, \mathbb{Q}} = Z_{b, \mathbb{Q}} = \emptyset$.

We prove (2.82), (2.83) below. Now, we present the rest of our claims under the assumption that $Z'_{a, \mathbb{Q}} = Z_{b, \mathbb{Q}} = \emptyset$. For the rest of this proof we will be assuming that all tensor fields have the factor ∇Y (or the factors $\nabla \chi_1, \nabla \chi_2$) not contracting against special indices.

We then perform a new induction: We assume that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^k} a_t X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_k} C_g^{t, i_1 \dots i_k}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.84}$$

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^k} a_t X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_k} C_g^{t, i_1 \dots i_k}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.85}$$

where the tensor fields indexed in T^k have all the properties of the tensor fields indexed in Z'_a (in particular the index in $\nabla \psi_1$ is not free) and in addition have rank k . We then show that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \operatorname{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^{k+1}} a_t X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_{k+1}} C_g^{t, i_1 \dots i_{k+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.86}$$

$$\begin{aligned}
& \sum_{\zeta \in Z'_a} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{\zeta \in Z_b} a_\zeta X_+ \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{t \in T^{k+1}} a_t X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_{k+1}} C_g^{t, i_1 \dots i_{k+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \\
& + \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau).
\end{aligned} \tag{2.87}$$

We will derive (2.86), (2.87) momentarily. For now, we observe that by iterative repetition of the above we are reduced to showing (2.74), (2.75) under the additional assumption that $Z'_a = \emptyset$. In that setting, we can just repeatedly apply the eraser (see the Appendix in [6] for a definition of this notion) to as many factors $\nabla\psi_\tau$ as needed in order to reduce ourselves to a new true equation where each of the real factors is contracting against at most one of the factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla Y$ (or $\nabla\chi_1, \nabla\chi_2$).⁶¹ Then, by invoking Corollary 1 from [6]⁶² and then re-introducing the factors we erased, we derive our claim.

Proof of (2.86), (2.87): Picking out the sublinear combination in (2.84), (2.85) with one derivative on ∇Y or $\nabla\chi_1$ and substituting into (2.72), (2.73) we derive a new equation:

$$\begin{aligned}
& \sum_{t \in T^k} a_t X_* \text{div}_{i_1} \dots X_* \text{div}_{i_k} C_g^{t, i_1 \dots i_k}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau),
\end{aligned} \tag{2.88}$$

⁶¹All remaining factors $\nabla\psi_1, \dots, \nabla\psi_\tau$ and also the factor(s) ∇Y (or $\nabla\chi_1, \nabla\chi_2$) are treated as factors $\nabla\phi_h$

⁶²Notice that there will necessarily be at least one non-simple factor $S_* \nabla^{(\nu)} R_{ijkl}$ or $\nabla^{(B)} \Omega_h$, by virtue of the factor(s) ∇Y (or $\nabla\omega_1, \nabla\omega_2$), therefore that Corollary can be applied.

$$\begin{aligned}
& \sum_{t \in T^k} a_t X_* \operatorname{div}_{i_1} \dots X_* \operatorname{div}_{i_k} C_g^{t, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{\zeta \in Z_b} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau);
\end{aligned} \tag{2.89}$$

(the sublinear combination $\sum_{\zeta \in Z_b} \dots$ above is *generic*).

We now divide the index set T^k according to which of the factors $\nabla\psi_2, \dots, \nabla\psi_\tau, \nabla Y$ (or $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla\chi_1$) contain the k free indices. Thus we write: $T^k = \bigcup_{\alpha \in A} T^{k, \alpha}$ (each $\alpha \in A$ corresponds to a k -subset of the set of factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla Y$ or $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla\chi_1$). We will then show that for each $\alpha \in A$ there exists a tensor field $\sum_{b \in B^\alpha} a_b C_g^{b, i_1 \dots i_{k+1}}$ in the form (2.64) or (2.65) with the first k free indices belonging to the factors in the set α , and the free index i_{k+1} *not* belonging to $\nabla\psi_1$, so that:

$$\begin{aligned}
& \sum_{t \in T^{k, \alpha}} a_t C_g^{t, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \dots \nabla_{i_k} v - \\
& X_* \operatorname{div}_{i_{k+1}} \sum_{b \in B^\alpha} a_b C_g^{b, i_1 \dots i_{k+1}} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \dots \nabla_{i_k} v \\
& = \sum_{j \in J} a_j C_g^{j, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \dots \nabla_{i_k} v,
\end{aligned} \tag{2.90}$$

$$\begin{aligned}
& \sum_{t \in T^{k, \alpha}} a_t C_g^{t, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \dots \nabla_{i_k} v - \\
& X_* \operatorname{div}_{i_{k+1}} \sum_{b \in B^\alpha} a_b C_g^{b, i_1 \dots i_{k+1}} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \dots \nabla_{i_k} v \\
& = \sum_{j \in J} a_j C_g^{j, i_1 \dots i_k} (\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \dots \nabla_{i_k} v.
\end{aligned} \tag{2.91}$$

If we can show the above for every $\alpha \in A$, then replacing the factor ∇v by $X_+ \operatorname{div}$'s we can derive our claim (2.86), (2.87).

Proof of (2.90), (2.91): Refer to (2.88) and (2.89). Denote Y or χ_1 by $\psi_{\tau+1}$ for uniformity. We pick out any $\alpha \in A$; assume that $\alpha = \{\nabla\psi_{x_1}, \dots, \nabla\psi_{x_k}\}$.

Pick out the sublinear combination where the factors $\nabla\psi_{x_1}, \dots, \nabla\psi_{x_k}$ which belong to α are contracting against the same factor as $\nabla\psi_1$. This sublinear combination Z_g vanishes separately (i.e. $Z_g = 0$). We then apply the eraser to

the factors $\nabla\psi_2, \dots, \nabla Y \in A$ (notice this is *well-defined*, since all the above factors *and* the factor $\nabla\psi_1$ are contracting against non-special indices). We obtain a new true equation, which we denote by $Eraser[Z_g] = 0$. It then follows that $Eraser[Z_g] \cdot (\nabla_{i_1}\psi_{x_1}\nabla^{i_1}v \dots \nabla_{i_k}\psi_{x_k}\nabla^{i_k}v) = 0$ is our desired conclusion (2.90), (2.91). \square

(*Sketch of Proof of (2.78), (2.79) (2.82), (2.83)*): These equations can be proven by only a slight modification of the idea above. We again subdivide the index sets T^k, V^k according to the set of factors $\nabla\psi_2, \dots, \nabla\psi_\tau$ or $\nabla\psi_2, \dots, \nabla\psi_\tau, \nabla\omega_1$ which contain the k free indices (so we write $T^k = \bigcup_{\alpha \in A} T^{k,\alpha}$ and $V^k = \bigcup_{\alpha \in A} V^{k,\alpha}$) and we prove the claims above separately for those sublinear combinations.

To prove this, we pick out the sublinear combination in our hypotheses with the factors $\nabla\psi_h, h \in \alpha$ contracting against the same factor against which $\nabla\psi_1$ and ∇Y (or $\nabla\psi_1$ and $\nabla\omega_1$) are contracting. Say $\alpha = \{h_1, \dots, h_k\}$; we then formally replace the expressions $S_* \nabla_{r_1 \dots r_\mu l_1 \dots l_k}^{(\nu)} R_{ijkl} \nabla_{l_1} \psi_{h_1} \dots \nabla_{l_k} \psi_{h_k} \nabla^i \tilde{\phi}_1 \nabla^j \psi_1 \nabla^k Y$ or $\nabla_{r_1 \dots r_\mu l_1 \dots l_k st}^{(A)} \Omega_1 \nabla^{l_1} \psi_{h_1} \dots \nabla_{l_k} \psi_{h_k} \nabla^s \psi_1 \nabla^t Y$ etc, by expressions $S_* \nabla_{r_1 \dots r_\mu}^{(\nu-k)} R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^j \psi_1 \nabla^k Y, \nabla_{r_1 \dots r_\mu st}^{(A-k)} \Omega_1 \nabla^s \psi_1 \nabla^t Y$ and derive our claims (2.78), (2.79) (2.82), (2.83) as above. \square

(*Proof of the claims of our general statement (i.e. (2.68), (2.69) by induction*): We will prove these claims by an induction. Our inductive assumptions are that (2.68), (2.69) follow from (2.66), (2.67) for any weight $-W', W' < K$ and when $W' = K$ they hold for any length $v + b \geq \gamma \geq 2$. We will then show the claim when the weight is $-K$, and $v + b = \gamma + 1$. In the end, we will check our claims for the base case $v + b = 2$.

(*Proof of the inductive step*): Refer back to (2.66), (2.67). We will prove this claim in four steps.

Step 1: Firstly, we will denote by $Z_a^{spec}, \overline{Z}_a^{spec}, Z_b^{spec}$ the index sets of the tensor fields for which ∇Y or one of the factors $\nabla\chi_1, \nabla\chi_2$ (respectively) is contracting against a special index. Then *using the inductive assumptions of our general claim*, we will show that there exists a linear combination of 2-tensor fields (indexed in W below) which satisfies all the requirements of (2.66), (2.68) so that:

$$\begin{aligned} \sum_{\zeta \in Z_a^{spec}} a_\zeta C_g^{\zeta, i_1} \nabla_{i_1} v - X_* \text{div}_{i_2} \sum_{w \in W} a_w C_g^{w, i_1 i_2} \nabla_{i_1} v = \\ \sum_{\zeta \in Z_a^{OK}} a_\zeta C_g^{\zeta, i_1} \nabla_{i_1} v + \sum_{j \in J} a_j C_g^{j, i_1} \nabla_{i_1} v, \end{aligned} \quad (2.92)$$

where the tensor fields in Z_a^{OK} are *generic* linear combinations of tensor fields of the same general type as the ones indexed in Z_a in (2.66), (2.68) and where in

addition none of the factors ∇Y or $\nabla\chi_1, \nabla\chi_2$ are contracting against a special index.

Thus, if we can show the above, by replacing ∇v by an X_*div_i , and substituting back into (2.66), (2.68), we are reduced to showing (2.67), (2.69) under the additional assumption that $Z_a^{spec} = \emptyset$.

Step 2: Then, under the assumption that $Z_a^{spec} = \emptyset$, we will show that we can write:

$$\begin{aligned} & \sum_{\zeta \in Z_b^{spec}} a_\zeta X_*div_{i_1} C_g^{\zeta, i_1} + \sum_{\zeta \in \bar{Z}_a^{spec}} a_\zeta X_*div_{i_1} \dots X_*div_{i_c} C_g^{\zeta, i_1 \dots i_c} = \\ & X_*div_{i_1} \dots X_*div_{i_b} \sum_{c \in C} a_c C_g^{c, i_1 \dots i_b} + \sum_{j \in J} a_j C_g^{j, i_1}, \end{aligned} \quad (2.93)$$

where the tensor fields on the RHS are of the general form as the ones indexed in Z_b, \bar{Z}_a in our hypothesis, and moreover the factor ∇Y (or the factors $\nabla\chi_1, \nabla\chi_2$) is (are) not contracting against special indices.

Notice that if we can show (2.92), (2.93) then we are reduced to showing our claim under the additional assumption that for each $\zeta \in Z_a \cup \bar{Z}_a \cup Z_b$ the factor(s) ∇Y (or $\nabla\chi_1, \nabla\chi_2$) are not contracting against special indices. We will show (2.92), (2.93) below.

Proof of (2.67), (2.69) under the additional assumption that for each $\zeta \in Z_a \cup \bar{Z}_a \cup Z_b$ the factor ∇Y or $(\nabla\chi_1, \nabla\chi_2)$ is not contracting against special indices:

Step 3: Proof of (2.94) below:

We note that for all the tensor fields in the rest of this proof will *not* have the factor ∇Y (or any of the factors $\nabla\chi_1, \nabla\chi_2$) contracting against a special index in any factor $S_*\nabla^{(\nu)}R_{ijkl}$ or $\nabla^{(B)}\Omega_h$. Now, we arbitrarily pick out one factor $T = S_*\nabla^{(\nu)}R_{ijkl}$ or $T = \nabla^{(B)}\Omega_x$ in $\bar{\kappa}_{simp}$ and call it the ‘‘chosen factor’’ for the rest of this subsection.

We will say that the factor ∇Y (or $\nabla\omega_2$) is contracting against a good index in T , if it is contracting against a non-special index in T when T is of the form $S_*\nabla^{(\nu)}R_{ijkl}$ with $\nu > 0$; when T is of the form $\nabla^{(B)}\Omega_x$, then it is contracting against a good index provided $B \geq 3$.

We will say that the factor ∇Y (or $\nabla\omega_2$) is contracting against a bad index if it is contracting against the index j in a factor $T = S_*R_{ijkl}$ or an index in a factor $T = \nabla^{(2)}\Omega_x$. We denote by $Z_a^{BAD} \subset Z_a$ the index set of tensor fields for which ∇Y (or $\nabla\omega_2$) is contracting against a bad index. We also denote by $Z_b^{BAD} \subset Z_b$ the index set of the vector fields for which ∇Y is contracting against a bad index in T and T also contains a free index. We will show that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z_a^{BAD} \cup Z_b^{BAD}} a_\zeta C_g^{\zeta, i_1} \nabla_{i_1} v - X_* \text{div}_{i_2} \sum_{h \in H} a_h C_g^{i_1 i_2} \nabla_{i_1} v = \\
& \sum_{\zeta \in Z_a^{GOOD} \cup Z_b^{GOOD}} a_\zeta C_g^{\zeta, i_1} \nabla_{i_1} v + \sum_{j \in J} a_j C_g^j,
\end{aligned} \tag{2.94}$$

where all the tensor fields indexed in $Z_a^{GOOD} \cup Z_b^{GOOD}$ are *generic* vector fields of the forms indexed in Z_a, Z_b , only with the factors ∇Y or $\nabla \omega_2$ contracting against a *good* index in the factor T . The tensor fields indexed in H are as required in the claim of our general statement (they correspond to the index set W in our general statement).

Step 4: Proof that (2.94) implies our claims (2.68), (2.69).

We start by proving (2.94) (i.e. we prove Step 3). Then, we will show how we can derive our claim from (2.94) (i.e. we then prove Step 4).

Proof of Step 3: Proof of (2.94): We can prove this equation by virtue of our inductive assumption on our general claim. First, we define $\bar{Z}_a^{BAD} \subset \bar{Z}_a$ to stand for the index set of tensor fields where the factor ∇Y (or $\nabla \omega_2$) is contracting against a bad index in the chosen factor. We pick out the sublinear combination in our Lemma assumption where ∇Y (or $\nabla \omega_2$) are contracting against the chosen factor $T = S_* R_{ijkl}$ or $T = \nabla^{(2)} \Omega_x$. This sublinear combination must vanish separately, and we thus derive an equation:

$$\begin{aligned}
& \sum_{\zeta \in Z_a^{BAD} \cup Z_b^{BAD}} a_\zeta X_{**} \text{div}_{i_1} C_g^{\zeta, i_1} + \sum_{\zeta \in \bar{Z}_a^{BAD}} a_\zeta X_{**} \text{div}_{i_1} \dots X_{**} \text{div}_{i_c} C_g^{\zeta, i_1 \dots i_c} + \\
& \sum_{\zeta \in Z_b^{nvBAD}} a_\zeta C_g^{f, i_1} = \sum_{j \in J} a_j C_g^j,
\end{aligned} \tag{2.95}$$

where $X_{**} \text{div}_{i_1}$ stands for the sublinear combination for which ∇_{i_1} is not allowed to hit the chosen factor T . $Z_b^{nvBAD} \subset Z_b$ stands for the index set of tensor fields indexed in Z_b with the free index i_1 *not* belonging to the chosen factor and also with the factor ∇Y (or $\nabla \omega_2$) contracting against a bad index.

Now, define an operation $Op[\dots]$ which acts on the complete contractions above by formally replacing any expression $\nabla_{ij}^{(2)} \Omega_x \nabla^i Y$ (or $\nabla_{ij}^{(2)} \Omega_x \nabla^i \chi_2$) by $\nabla_j D$ (D is a scalar function), or any expression $S_* R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^j Y$ (or $S_* R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^j \chi_2$) by $\nabla_{[k} \theta_1 \nabla_{l]} \theta_2$. (Denote by $\tilde{\kappa}_{simp}$ the simple character of these resulting vector fields). Acting on (2.95) by $Op[\dots]$ produces a true equation, which we may write out as:

$$\begin{aligned}
& \sum_{\zeta \in Z_a^{BAD} \cup Z_b^{BAD}} a_\zeta X_{**} \text{div}_{i_1} \text{Op}[C]_g^{\zeta, i_1} + X_{**} \text{div}_{i_1} \sum_{f \in F} a_f C_g^{f, i_1} \\
& + \sum_{\zeta \in \bar{Z}_a^{BAD}} a_\zeta X_{**} \text{div}_{i_1} \dots X_{**} \text{div}_{i_c} C_g^{\zeta, i_1 \dots i_c} = \sum_{j \in J} a_j C_g^j.
\end{aligned} \tag{2.96}$$

Here $X_{**} \text{div}_i$ stands for the sublinear combination in div_i where ∇_i is not allowed to hit the factor to which ∇_i belongs, nor any of the factors $\nabla\phi_1, \dots, \nabla\phi_u, \nabla\psi_1, \dots, \nabla\psi_\tau$, nor any factors $\nabla D, \nabla\theta_1, \nabla\theta_2$. The vector fields indexed in F are generic vector fields with a simple character $\tilde{\kappa}_{simp}$, for which the free index i_1 does not belong to any of the factors $\nabla\psi_1, \dots, \nabla\psi_\tau$ or any of the factors $\nabla D, (\nabla\chi_1), \nabla\theta_1, \nabla\theta_2$.

Now, observe that the above equation falls under our inductive assumption of the general statement we are proving: We now either have factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla D$, or $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla\chi_1, \nabla D$ or $\nabla\psi_1, \dots, \nabla\psi_\tau, [\nabla\theta_1, \nabla\theta_2]$ or $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla\chi_1, [\nabla\theta_1, \nabla\theta_2]$. Notice that the tensor fields indexed in H_a^{BAD}, H_b^{BAD} are precisely the ones that contain a free index in one of these factors. Therefore, by our inductive assumption of the “general claim” we derive that there exists a linear combination of 2-tensor fields, $\sum_{v \in V} \dots$, (with factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla D$ etc, and which satisfy the @-property for the factors $\nabla\psi_1, \dots, \nabla\psi_\tau$) so that:

$$\begin{aligned}
& \sum_{\zeta \in Z_a^{BAD} \cup Z_b^{BAD}} a_\zeta \text{Op}[C]_g^{\zeta, i_1} \nabla_{i_1} v - X_{**} \text{div}_{i_2} \sum_{v \in V} a_v C_g^{v, i_1 i_2} \nabla_{i_1} v = \\
& \sum_{j \in J} a_j C_g^{j, i_1} \nabla_{i_1} v.
\end{aligned} \tag{2.97}$$

Now, we define an operation $\text{Op}^{-1}[\dots]$, which acts on the complete contractions in the above equation by replacing the factor $\nabla_j D$ by an expression $\nabla_{ij} \Omega_x \nabla^j Y$ (or $\nabla_{ij} \Omega_x \nabla^j \omega_2$) or the expression $\nabla_{[a} \theta_1 \nabla_{b]} \theta_2$ by $S_* R_{ijab} \nabla^i \tilde{\phi}_1 \nabla^j Y$ (or $S_* R_{ijab} \nabla^i \tilde{\phi}_1 \nabla^j \omega_2$). The operation Op^{-1} clearly produces a true equation, which is our desired conclusion, (2.94). \square

Proof of Step 4: We derive our conclusions (2.68), (2.69) in pieces. Firstly, we show these equations with the sublinear combinations Z_a replaced by the index set $Z_{a, spec}$, which index the terms with the free index i_1 belonging to the factor ∇Y or $\nabla\omega_1$ (this will be sub-step A). After proving this claim, we will show (2.68), (2.69) under the additional assumption that $Z_{a, spec} = \emptyset$ (this will be sub-step B).

Proof of sub-step A: We make the ∇v 's into $X_* \text{div}$'s in (2.94) and replace the resulting equations into our Lemma hypothesis. We thus derive a new equation:

$$\begin{aligned}
& \sum_{\zeta \in Z_a} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1} + \sum_{\zeta \in Z_b^1} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1} + \\
& \sum_{\zeta \in Z_b^2} a_\zeta X_* \text{div}_{i_1} \dots X_* \text{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in J} a_j C_g^j = 0,
\end{aligned} \tag{2.98}$$

where we now have the tensor fields indexed in Z_a have a free index among the factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla Y$ (or $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla\chi_1, \nabla\chi_2$), and furthermore the factor ∇Y (or the factors $\nabla\omega_1, \nabla\omega_2$) are not contracting against a bad index in the chosen factor T . The tensor fields indexed in Z_b^1 have a free index that *does* not belong to one of the factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla Y$ (or $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla\chi_1, \nabla\chi_2$), and furthermore *if* the factor ∇Y (or one of the factors $\nabla\omega_1, \nabla\omega_2$) is contracting against a bad index in the chosen factor T , then T does not contain the free index i_1 . Finally the tensor fields indexed in Z_b^2 each have rank $a \geq 2$ and all free indices belong to the factors $\nabla\psi_1, \dots, \nabla\psi_\tau, \nabla Y, (\nabla\omega_1, \nabla\omega_2)$. We may then re-write our equation (2.98) in the form:

$$\begin{aligned}
& \sum_{\zeta \in Z_a} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1} + \sum_{\zeta \in Z_b^1} a_\zeta X_* \text{div}_{i_1} C_g^{\zeta, i_1} + \\
& \sum_{\zeta \in Z_b^{2'}} a_\zeta X_* \text{div}_{i_1} \dots X_* \text{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in J} a_j C_g^j = 0,
\end{aligned} \tag{2.99}$$

where now for the tensor fields indexed in $Z_b^{2'}$, each $a \geq 1$ and the factor $\nabla\psi_1$ does not contain a free index for any of the tensor fields for which ∇Y (or one of $\nabla\omega_1, \nabla\omega_2$) is contracting against a bad index in the chosen factor.

We will denote by $Z_{b, \#}^1 \subset Z_b^1$ and $Z_{b, \#}^{2'} \subset Z_b^{2'}$ the index sets of tensor fields where ∇Y (or one of $\nabla\omega_1, \nabla\omega_2$) is contracting against a bad index in the chosen factor T .

From (2.99) we derive an equation:

$$\begin{aligned}
& \sum_{\zeta \in Z_{b, \#}^1} a_\zeta X_{**} \text{div}_{i_1} C_g^{\zeta, i_1} + \\
& \sum_{\zeta \in Z_{b, \#}^{2'}} a_\zeta X_{**} \text{div}_{i_1} \dots X_{**} \text{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in J} a_j C_g^j = 0,
\end{aligned} \tag{2.100}$$

where $X_{**} \text{div}_i$ stands for the sublinear combination in $X_* \text{div}_i$ for which ∇_i is in addition no allowed to hit the chosen factor T .

Then, applying operation Op as in Step 3 and the the inductive assumption of the general claim we are proving,⁶³ and then using the operation $Op^{-1}[\dots]$ as in the proof of Step 3, we derive a new equation:

⁶³The resulting equation falls under the inductive assumption, as in Step 3.

$$\begin{aligned}
& \sum_{\zeta \in Z_{b,\sharp}^1} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1} + \sum_{\zeta \in Z_{b,\sharp}^2} a_\zeta X_* \operatorname{div}_{i_1} \dots X_* \operatorname{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} = \\
& \sum_{\zeta \in Z_{OK}} a_\zeta X_* \operatorname{div}_{i_1} \dots X_* \operatorname{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in J} a_j C_g^j = 0,
\end{aligned} \tag{2.101}$$

where the tensor fields indexed in Z_{OK} have rank $a \geq 1$ (all free indices *not* belonging to factors $\nabla\psi_1, \dots, \nabla Y$ or $\nabla\psi_1, \dots, \nabla\chi_2$) and furthermore have the property that the one index in ∇Y or $\nabla\omega_1$ is not contracting against a bad index in the chosen factor (and it is also not free). Thus, replacing the above back into (2.99), we derive:

$$\begin{aligned}
& \sum_{\zeta \in Z_a} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1} + \sum_{\zeta \in Z_b^1} a_\zeta X_* \operatorname{div}_{i_1} C_g^{\zeta, i_1} + \\
& \sum_{\zeta \in Z_b^{2''}} a_\zeta X_* \operatorname{div}_{i_1} \dots X_* \operatorname{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in J} a_j C_g^j = 0,
\end{aligned} \tag{2.102}$$

where the tensor fields indexed in $Z_b^1, Z_b^{2''}$ have the additional restriction that if the factor ∇Y (or $\nabla\omega_1, \nabla\omega_2$) is contracting against the chosen factor T then it is not contracting against a bad index in T .

We are now in a position to derive sub-step A from the above: To see this claim, we just apply $\operatorname{Erase}_{\nabla Y}$ or $\operatorname{Erase}_{\nabla\omega_1}$ to (2.102) and multiply the resulting equation by $\nabla_{i_1} Y \nabla^{i_1} v$.

Sub-step B: Now, we are reduced to showing our claim when $Z_{a, \text{spec}} = \emptyset$. In that setting, we denote by $Z_{a,s} \subset Z_a$ the index set of vector fields in Z_a for which the free index i_1 belongs to the factor $\nabla\psi_s$; we prove our claim separately for each of the sublinear combinations $\sum_{\zeta \in Z_{a,s}} \dots$. This claim is proven by picking out the sublinear combinations in (2.66), (2.67) where the factors $\nabla\psi_s$ and ∇Y (or $\nabla\chi_1$) are contracting against the same factor; we then apply the eraser to $\nabla\psi_s$ (this is well-defined and produces a true equation), and multiply by $\nabla_{i_1} \psi_s \nabla^{i_1} v$. The resulting equation is precisely our claim for the sublinear combination $\sum_{\zeta \in Z_{a,s}} \dots$.

(Sketch of the) Proof of Steps 1 and 2 (i.e. of (2.92) and (2.93)): We will sketch the proof of these claims for the sublinear combinations in $Z_a^{\text{spec}} \cup Z_b^{\text{spec}} \cup \overline{Z}_{\text{spec}}^a$ where one of the special indices in C^{ζ, i_1} is an index k or l that belongs to a factor $S_* \nabla^{(\nu)} R_{ijkl}$. The remaining case (where the special indices belong to factors $\nabla^{(a)} \Omega_h$) can be seen by a similar (simpler) argument.⁶⁴

For each $\zeta \in Z_a^{\text{spec}} \cup Z_b^{\text{spec}} \cup \overline{Z}_{\text{spec}}^a$ We denote by $\overline{C}_g^{\zeta, i_1}, \overline{C}_g^{\zeta, i_1 \dots i_\gamma}$ the tensor fields that arise from $C^{\zeta, i_1} C_g^{\zeta, i_1 \dots i_\gamma}$ in (2.66), (2.68) by replacing the expressions

⁶⁴The only extra feature in this setting is that one must prove the claim by a separate induction on the *number* of factors $\nabla\psi_z$ that are contracting against $\nabla^{(a)} \Omega_h$.

$S_* \nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^k Y$, $S_* \nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^k \chi_2$ by a factor $\nabla_{r_1 \dots r_\nu, j l}^{(\nu+2)} \Omega_{b+1}$. We denote by $\tilde{\kappa}_{simp}$ the resulting simple character. We derive an equation:

$$\begin{aligned} & \sum_{\zeta \in Z_a^{spec} \cup Z_b^{spec}} a_\zeta X_* \text{div}_{i_1} \overline{C}_g^{\zeta, i_1} + \\ & \sum_{\zeta \in \overline{Z}_a^{spec}} a_\zeta X_* \text{div}_{i_1} \dots X_* \text{div}_{i_\tau} \overline{C}_g^{\zeta, i_1} + \sum_{j \in J} a_j \overline{C}_g^j = 0. \end{aligned} \quad (2.103)$$

Now, again applying the inductive assumption of our general statement, we derive that there is a linear combination of tensor fields (indexed in W below) with a free index i_1 belonging to one of the factors $\nabla \psi_1, \dots, \nabla \psi_\tau$ or $\nabla \psi_1, \dots, \nabla \psi_\tau, \nabla \chi_1$ so that:

$$\sum_{\zeta \in Z_a^{spec}} a_\zeta \overline{C}_g^{\zeta, i_1} \nabla_{i_1} v - X_* \text{div}_{i_2} \sum_{w \in W} a_w C_g^{w, i_1 i_2} \nabla_{i_1} v = \sum_{j \in J} a_j \overline{C}_g^j. \quad (2.104)$$

Now, applying an operation Op^* to the above which formally replaces the factor $\nabla_{r_1 \dots r_A}^{(A)} \Omega_x$ by a factor $S_* \nabla_{r_1 \dots r_{A-2}}^{(A-2)} R_{i r_{A-1} k r_A} \nabla^i \tilde{\phi}_1 \nabla^k Y$ or $S_* \nabla_{r_1 \dots r_{A-2}}^{(A-2)} R_{i r_{A-1} k r_A} \nabla^i \tilde{\phi}_1 \nabla^k \chi_2$, we derive (2.92) (since we can repeat the permutations by which (2.104) is made to hold formally, modulo introducing correction terms that allowed in the RHS of (2.92)).

We will now prove (2.93) by repeating the induction performed in the ‘‘Consequence’’ we derived above (where we showed that inductively assuming (2.84), (2.85) we can derive (2.78), (2.79)):

We will show the claim of Step 2 in pieces: First consider the tensor fields indexed in $\overline{Z}_{a, \textcircled{a}}$ of minimum rank 2 (denote the corresponding index set is $\overline{Z}_{a, \textcircled{a}}^2$); we then show that we can write:

$$\begin{aligned} & \sum_{\zeta \in \overline{Z}_{a, \textcircled{a}}^2} a_\zeta X \text{div}_{i_1} X \text{div}_{i_2} C_g^{\zeta, i_1 i_2} = \\ & \sum_{\zeta \in \overline{Z}_{a, \textcircled{a}}^3} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_3} C_g^{\zeta, i_1 \dots i_3} + \sum_{\zeta \in Z_{b, \textcircled{a}}} a_\zeta X \text{div}_{i_1} C_g^{\zeta, i_1} + \\ & \sum_{\zeta \in Z_{OK}} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in J} a_j C^j. \end{aligned} \quad (2.105)$$

The tensor fields indexed in $\overline{Z}_{a, \textcircled{a}}^3, Z_{b, \textcircled{a}}$ in the RHS are generic linear combinations in those forms (the first with rank 3). The tensor fields indexed in Z_{OK} are generic linear combinations as allowed in the RHS of our Step 2. Assuming we can prove (2.105), we are then reduced to showing our claim when the minimum rank among the tensor fields indexed in $Z_{a, \textcircled{a}}$ is 3. We may then ‘‘forget’’ about

any $X \text{div}_{i_h}$ where i_h belongs to the factor $\nabla\psi_1$. Therefore, we are reduced to showing our claim when the minimum rank is 2 and the factor $\nabla\psi_1$ does not contain a free index. We then show our claim by an induction (for the rest of this derivation, all tensor fields will *not* have a free index in the factor $\nabla\psi_1$): Assume that the minimum rank of the tensor fields indexed in $\overline{Z}_{a,\otimes}$ is k , and they are indexed in $\overline{Z}_{a,\otimes}^k$. We then show that we can write:

$$\begin{aligned}
& \sum_{\zeta \in \overline{Z}_{a,\otimes}^k} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_k} C_g^{\zeta, i_1 \dots i_k} = \\
& \sum_{\zeta \in \overline{Z}_{a,\otimes}^{k+1}} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_{k+1}} C_g^{\zeta, i_1 \dots i_{k+1}} + \sum_{\zeta \in Z_{b,\otimes}} a_\zeta X \text{div}_{i_1} C_g^{\zeta, i_1} + \quad (2.106) \\
& \sum_{\zeta \in Z_{OK}} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in} a_j C^j.
\end{aligned}$$

The tensor fields indexed in $\overline{Z}_{a,\otimes}^3, Z_{b,\otimes}$ in the the RHS are generic linear combinations in those forms (the first with rank $k+1$). The tensor fields indexed in Z_{OK} are generic linear combinations as allowed in the RHS of our Step 2.

Iteratively repeating this step we are reduced to showing our Step 2 when $Z_{a,\otimes} = \emptyset$.

In that case we then assume that the tensor fields indexed in $Z_{b,\otimes}$ have minimum rank k (and the corresponding index set is $Z_{b,\otimes}^k$) and we show that we can write:

$$\begin{aligned}
& \sum_{\zeta \in Z_{b,\otimes}^k} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_k} C_g^{\zeta, i_1 \dots i_k} = \\
& \sum_{\zeta \in Z_{b,\otimes}^{k+1}} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_{k+1}} C_g^{\zeta, i_1 \dots i_{k+1}} + \quad (2.107) \\
& \sum_{\zeta \in Z_{OK}} a_\zeta X \text{div}_{i_1} \dots X \text{div}_{i_a} C_g^{\zeta, i_1 \dots i_a} + \sum_{j \in} a_j C^j,
\end{aligned}$$

(with the same conventions as in the above equation).

If we can prove (2.105) and (2.107) we will have shown our step 2.

Proof of (2.105), (2.106), (2.107): We start with a small remark: If the chosen factor is of the form $S_* \nabla^{(\nu)} R_{ijkl}$, we replace our assumption by a more convenient equation: Consider the tensor fields $C_g^{\zeta, i_1 \dots i_a}$, $\zeta \in \overline{Z}_{a,\otimes} \cup Z_{b,\otimes}$; we denote by $\tilde{C}_g^{\zeta, i_1 \dots i_a}$ the tensor fields that arise from $C_g^{\zeta, i_1 \dots i_a}$ by replacing the expression $\nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^k Y$ (or $\nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_1 \nabla^k \chi_2$) by a factor $\nabla_{r_1 \dots r_\nu j l}^{(\nu+2)} \Omega_{p+1}$. We then derive an equation:

$$\begin{aligned}
& \sum_{\zeta \in \bar{Z}_a \cup Z_b} a_\zeta X_* \text{div}_{i_1} \dots X_* \text{div}_{i_a} \tilde{C}_g^{\zeta, i_1 \dots i_a} (\Omega_1, \dots, \Omega_{p+1}, \phi_2, \dots, \phi_u, (\chi_1), \psi_1, \dots, \psi_\tau) \\
& + \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_{p+1}, \phi_2, \dots, \phi_u, (\chi_1), \psi_1, \dots, \psi_\tau).
\end{aligned} \tag{2.108}$$

Now we can derive our claims:

Proof of (2.106): We divide the index set $Z_{\bar{Z}_a, \textcircled{a}}$ according to the two factors which contain the two free indices and we show our claim for each of those tensor fields separately. The proof goes as follows: We pick out the sublinear combination in our hypothesis (or in (2.108)) where the factors $\nabla\psi_h, \nabla\psi_{h'}$ (or $\nabla\psi_h, \nabla\chi_2$) are contracting against the same factor. Clearly, this sublinear combination, X_g , vanishes separately. We then formally erase the factor $\nabla\psi_h$. Then, we may apply the inductive assumption of our general claim to the resulting equation (the minimum rank of the tensor fields will be 1), and (in case our assumption is (2.108) we also apply an operation Op^{-1} which replaces the factor $\nabla_{r_1 \dots r_y}^{(y)} \Omega_{p+1}$ by $S_* \nabla_{r_1 \dots r_{y-2}}^{(y-2)} R_{ir_{y-1}kr_y} \nabla^i \tilde{\phi}_1 \nabla^k Y(\nabla^k \chi_1)$). This is our desired conclusion.

Proof of (2.105), (2.107): Now, we show (2.105) for the subset $Z_{\bar{Z}_a, \textcircled{a}}^{k, \alpha}$ (which indexes the k -tensor fields for which the free indices i_1, \dots, i_k belong to a chosen subset of the factors $\nabla\psi_1, \dots, \nabla\psi_\tau, (\nabla\chi_1)$ (hence the label α designates the chosen subset). To prove this equation, we pick out the sublinear combination in the equation (2.108) where the factors $\nabla\psi_2, \dots, \nabla\psi_\tau, (\nabla\chi_1)$ (indexed in α) are contracting against the same factor as $\nabla\psi_1$. Then we apply the eraser to these factors and the indices they contract against. This is our desired conclusion. To show (2.107), we only have to treat the factors $\nabla\psi_h$ as factors $\nabla\phi_h$. The claim then follows by applying Corollary 1 in [6] and making the factors ∇v into $X \text{div}$'s.⁶⁵ \square

Proof of the base case ($v + b = 2$) of the general claim: We firstly prove our claim when our hypothesis is (2.67) (as opposed to (2.66)).

Proof of the base case under the hypothesis (2.67): We observe that the weight $-K$ in our assumption must satisfy $K \geq 2\tau + 8$ if $v > 0$ and $K \geq 2\tau + 6$ if $v = 0$.

First consider the case where we have the strict inequalities $K > 2\tau + 8$ if $v > 0$ and $K > 2\tau + 6$ if $v = 0$. In that case our first claim of the base case can be proven straightforwardly, by picking out a removable index in each $C_g^{\zeta, i_a}, \zeta \in Z_a$ and treating it as an $X_* \text{div}$ (which can be done when we only have

⁶⁵Observe that by virtue of the factor $\nabla\psi_1$, we must have at least one non-simple factor $S_* \nabla^{(\nu)} R_{ijkl}$ or $\nabla^{(B)} \Omega_h$ in (2.108)—hence (2.108) does not fall under any of the “forbidden cases” of Corollary 1 in [6], by inspection.

two real factors). Thus, in this setting we only have to show our second claims (2.70), (2.71).

In this setting, by using the “manual” constructions as in [5], we can construct explicit tensor fields which satisfy all the assumptions of our claim in the base case (each with rank ≥ 2), so that:

$$\begin{aligned}
& X_+ \operatorname{div}_{i_1} \sum_{\zeta \in Z'_a \cup Z_b} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{p \in P} a_p X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_{c+1}} C_g^{p, i_1 \dots i_{c+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \\
& \psi_1, \dots, \psi_\tau) + \sum_{j \in J} a_j C_g^j
\end{aligned} \tag{2.109}$$

Here the tensor field $C_g^{p, i_1 \dots i_{c+1}}$ will be in one of three forms:

If $v = 2$ then each $C_g^{p, i_1 \dots i_{c+1}}$ will be in the form:

$$\begin{aligned}
& p \operatorname{contr}(S_* \nabla^{(\nu_1)}{}_{i_1 \dots i_{c-1}}{}^{f_{b_1} \dots f_{b_h}} R_{x_1 j i_c l} \otimes S_* \nabla^{(\nu_2)}{}^{f_{d_1} \dots f_{d_y}} R_{x_v}{}^{j' i_{c+1} l} \otimes \\
& \otimes [\nabla^j \chi_1 \otimes \nabla_{j'} \chi_2] \otimes \nabla_{f_1} \psi_1 \dots \otimes \nabla_{f_\tau} \psi_\tau \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \nabla^{x_2} \tilde{\phi}_2),
\end{aligned} \tag{2.110}$$

where $\{b_1, \dots, b_h, d_1, \dots, d_y\} = \{1, \dots, \tau\}$.

If $v = 1$ then $\sum_{p \in P} \dots = 0$ (this can be arranged because of the two anti-symmetric indices k, l in the one factor $S_* \nabla^{(\nu)} R_{ijkl}$).

If $v = 0$ then each $C_g^{p, i_1 \dots i_{c+1}}$ will be in the form:

$$\begin{aligned}
& p \operatorname{contr}(\nabla^{(A_1)}{}_{i_1 \dots i_{c-1} j i_c}{}^{f_{b_1} \dots f_{b_h}} \Omega_1 \otimes \nabla^{(A_2)}{}^{f_{d_1} \dots f_{d_y} j' i_{c+1}} \Omega_2 \\
& \otimes [\nabla^j \chi_1 \otimes \nabla_{j'} \chi_2] \otimes \nabla_{f_1} \psi_1 \dots \otimes \nabla_{f_\tau} \psi_\tau \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \nabla^{x_2} \tilde{\phi}_2),
\end{aligned} \tag{2.111}$$

where $\{b_1, \dots, b_h, d_1, \dots, d_y\} = \{1, \dots, \tau\}$.

Then, picking out the sublinear combination in (2.120), (2.121) with factors $\nabla \psi_1, \dots, \nabla \psi_\tau, \nabla \chi_1, \nabla \chi_2$ we derive that $\sum_{p \in P} \dots = 0$. This is precisely our desired conclusion in this case.

Now, the case where we have the equalities in our Lemma hypothesis, $K = 2\tau + 8$ if $v > 0$ and $K = 2\tau + 6$ if $v = 0$. In this case we note that in our hypothesis $Z_b = \emptyset$ if $v \neq 1$, while $Z_a = \bar{Z}_a = \emptyset$ if $v = 1$.

Then, if $v \neq 1$, by the “manual” constructions as in [5], it follows that we can construct tensor fields (as required in the claim of our “general claim”), so that:

$$\begin{aligned}
& \sum_{\zeta \in Z_a} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v \\
& X_* \operatorname{div}_{i_2} a_\zeta C_g^{\zeta, i_1 i_2}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v = \\
& a_* C_g^{*, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v + \\
& \sum_{j \in J} a_j C_g^{j, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \nabla_{i_1} v,
\end{aligned} \tag{2.112}$$

where the tensor field C_g^{*, i_1} is in the form:

$$\begin{aligned}
& p\operatorname{contr}(S_* \nabla^{(\nu_1)^{f_1 \dots f_{\tau-1}}} R_{x_1}{}^{f_\tau}{}_{kl} \otimes R_{x_2}{}^{j'kl} \otimes \otimes \\
& [\nabla_{i_1} \chi_1 \otimes \nabla_{j'} \chi_2] \otimes \nabla_{f_1} \psi_1 \cdots \otimes \nabla_{f_\tau} \psi_\tau \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \nabla^{x_2} \tilde{\phi}_2),
\end{aligned} \tag{2.113}$$

if $v = 2$, and in the form:

$$\begin{aligned}
& p\operatorname{contr}(\nabla^{(\tau+1)^{f_1 \dots f_\tau}} \Omega_1 \otimes \nabla^{j's} \Omega_2 \otimes \\
& [\nabla_{i_1} \chi_1 \otimes \nabla_{j'} \chi_2] \otimes \nabla_{f_1} \psi_1 \cdots \otimes \nabla_{f_\tau} \psi_\tau \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \nabla^{x_2} \tilde{\phi}_2),
\end{aligned} \tag{2.114}$$

if $v = 0$.

Thus, we are reduced to the case where Z_a only consists of the vector field (2.113) or (2.114), and all other tensor fields in our Lemma hypothesis have rank ≥ 2 (we have denoted their index set by Z'_a). We then show that we can write:

$$\begin{aligned}
& X_+ \operatorname{div}_{i_1} \sum_{\zeta \in Z'_a} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \operatorname{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{p \in P} a_p X_+ \operatorname{div}_{i_1} \dots X_+ \operatorname{div}_{i_{c+1}} C_g^{p, i_1 \dots i_{c+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \\
& + \sum_{j \in J} a_j C_g^j,
\end{aligned} \tag{2.115}$$

where the tensor fields indexed in P here each have rank ≥ 2 and are all in the form:

$$\begin{aligned}
& p\operatorname{contr}(S_* \nabla^{(\nu_1)^{f_1 \dots f_{\tau-1}}} R_{x_1}{}^{f_\tau}{}_{ikl} \otimes S_* R_{x_v}{}^{j'kl} \otimes \\
& [\nabla_{i_1} \chi_1 \otimes \nabla_{j'} \chi_2] \otimes \nabla_{y_1} \psi_1 \cdots \otimes \nabla_{y_\tau} \psi_\tau \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \nabla^{x_2} \tilde{\phi}_2),
\end{aligned} \tag{2.116}$$

$$\begin{aligned}
& p\text{contr}(\nabla^{(\nu_1)^{f_1} \dots f_\tau} \Omega_1 \otimes \nabla^{j's} \Omega_2 \otimes \\
& [\nabla_{i_1} \chi_1 \otimes \nabla_{j'} \chi_2] \otimes \nabla_{y_1} \psi_1 \cdots \otimes \nabla_{y_\tau} \psi_\tau),
\end{aligned} \tag{2.117}$$

where each of the indices f_h is contracting against one of the indices y_q . The indices y_q that are *not* contracting against an index f_h are free indices.

Then, replacing the above into our Lemma hypothesis (and making all the ∇v 's into $X_+ \text{div}$'s), we derive that $a_p = 0$ for every $p \in P$ and $a_* = 0$. This concludes the proof of the base case when $v + b = 2$, $v \neq 1$. In the case $v = 1$ we show our claim by just observing that we can write:

$$\begin{aligned}
& X_+ \text{div}_{i_1} \sum_{\zeta \in Z_b} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \\
& + \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, \phi_1, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau);
\end{aligned} \tag{2.118}$$

this concludes the proof of the base case, when the tensor fields in our Lemma hypothesis are in the form (2.67).

Now, we consider the setting where our hypothesis is (2.66). We again observe that if $v = 0$ then the weight $-K$ in our hypothesis must satisfy $K \geq 2\tau + 4$. If $v > 0$ it must satisfy $K \geq 2\tau + 6$. We then again first consider the case where we have the strict inequalities in the hypothesis of our general claim.

In this case (where we have the strict inequalities $K > 2\tau + 4$ if $v = 0$ and $K > 2\tau + 6$ if $v \neq 0$) our first claim follows straightforwardly (as above, we just pick out one removable index in each C_g^{ζ, i_1} , $\zeta \in Z_a$ and treat it as an $X_* \text{div}$). To show the second claim we proceed much as before:

We can “manually” construct tensor fields in order to write:

$$\begin{aligned}
& X_+ \text{div}_{i_1} \sum_{\zeta \in Z'_a \cup Z_b} a_\zeta C_g^{\zeta, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, Y, \psi_1, \dots, \psi_\tau) = \\
& \sum_{q \in Q} a_q X_+ \text{div}_{i_1} C_g^{q, i_1}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) + \\
& \sum_{p \in P} a_p X_+ \text{div}_{i_1} \dots X_+ \text{div}_{i_{c+1}} C_g^{p, i_1 \dots i_{c+1}}(\Omega_1, \dots, \Omega_b, \phi_1, \dots, \phi_v, [\chi_1, \chi_2], \psi_1, \dots, \psi_\tau) \\
& + \sum_{j \in J} a_j C_g^j.
\end{aligned} \tag{2.119}$$

Here the tensor field $C_g^{p, i_1 \dots i_{c+1}}$ will be in one of three forms:

If $v = 2$ then each $C_g^{p, i_1 \dots i_{c+1}}$ will be:

$$pcontr(S_* \nabla^{(\nu_1)} \overset{f_{b_1} \dots f_{b_h}}{i_1 \dots i_{c-1}} R_{x_1} \overset{f_{b_{h+1}}}{i_{c,l}} \otimes S_* \nabla^{(\nu_2)} \overset{f_{d_1} \dots f_{d_y}}{i_{c+1}} R_{x_v} \overset{f_{d_{y+1}}}{i_{c+1}} \overset{l}{\otimes} \quad (2.120)$$

$$\nabla_{f_{\tau+1}} Y \otimes \nabla_{f_1} \psi_1 \dots \otimes \nabla_{f_\tau} \psi_\tau \otimes \nabla^{x_1} \tilde{\phi}_1 \otimes \nabla^{x_2} \tilde{\phi}_2),$$

where $\{b_1, \dots, b_{h+1}, d_1, \dots, d_{y+1}\} = \{1, \dots, \tau + 1\}$.

If $v = 1$ then $\sum_{p \in P} \dots = 0$ (this is because of the two antisymmetric indices k, l in the one factor $S_* \nabla^{(\nu)} R_{ijkl}$).

If $v = 0$ then each $C_g^{p, i_1 \dots i_{c+1}}$ will be in the form:

$$pcontr(\nabla^{(A_1)} \overset{f_{b_1} \dots f_{b_h}}{i_1 \dots i_{c-1} i_c} \Omega_1 \otimes \nabla^{(A_2)} \overset{f_{d_1} \dots f_{d_y}}{i_{c+1}} \Omega_2 \quad (2.121)$$

$$\otimes \nabla_{f_{\tau+1}} Y \otimes \nabla_{f_1} \psi_1 \dots \otimes \nabla_{f_\tau} \psi_\tau),$$

where $\{b_1, \dots, b_h, d_1, \dots, d_y\} = \{1, \dots, \tau + 1\}$.

Then, picking out the sublinear combination in (2.120), (2.121) with factors $\nabla \psi_1, \dots, \nabla \psi_\tau, \nabla Y$ we derive that $\sum_{p \in P} \dots = 0$. This is precisely our desired conclusion in this case.

Finally, we prove our claim when we have the equalities $K = 2\tau + 4$ if $v < 2$ and $K = 2\tau + 6$ if $v = 2$) in the hypothesis of our general claim.

In this case by “manually” constructing $X_+ div$'s so that we can write:

$$\sum_{\zeta \in Z'_a \cup Z_b \cup \bar{Z}_a} a_\zeta X_+ div_{i_1} \dots X_+ div_{i_a} C_g^{\zeta, i_1 \dots i_a}(\Omega_1, \dots, \Omega_b, Y, \psi_1, \dots, \psi_\tau) =$$

$$\sum_{q \in Q} a_q X_+ div_{i_1} \dots X_+ div_{i_a} C_g^{q, i_1 \dots i_a}(\Omega_1, \dots, \Omega_b, Y, \psi_1, \dots, \psi_\tau) +$$

$$\sum_{p \in P} a_p X_+ div_{i_1} \dots X_+ div_{i_a} C_g^{p, i_1 \dots i_a}(\Omega_1, \dots, \Omega_b, Y, \psi_1, \dots, \psi_\tau)$$

$$+ \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_b, Y, \psi_1, \dots, \psi_\tau). \quad (2.122)$$

Here the tensor fields indexed in P are in the following form:

If $v = 0$ then they will either be in the form:

$$pcontr(\nabla_{i_*} Y \otimes \nabla^{(A)} \overset{f_{x_1} \dots f_{x_a}}{s} \Omega_1 \otimes \nabla^{(B)} \overset{f_{x_{a+1}} \dots f_{x_\tau}}{s} \Omega_2 \otimes \nabla_{f_1} \psi_1 \otimes \dots \otimes \nabla_{f_\tau} \psi_\tau), \quad (2.123)$$

(where $\{x_1, \dots, x_\tau\} = \{1, \dots, \tau\}$), or in the form:

$$pcontr(\nabla_q Y \otimes \nabla^{(A)} \overset{f_{x_1} \dots f_{x_a}}{i_*} \Omega_1 \otimes \nabla^{(B)} \overset{f_{x_{a+1}} \dots f_{x_\tau}}{q} \Omega_2 \otimes \nabla_{f_1} \psi_1 \otimes \dots \otimes \nabla_{f_\tau} \psi_\tau), \quad (2.124)$$

(where $\{x_1, \dots, x_\tau\} = \{1, \dots, \tau\}$).

If $v = 2$ they will be in the form:

$$\begin{aligned} & p\text{contr}(\nabla_{i_*} Y \otimes \nabla^{(A) f_{x_1} \dots f_{x_{a-1}}} S_* R^{i f_{x_a} k l} \otimes \nabla^{(B) f_{x_{a+1}} \dots f_{x_{\tau-1}}} R^{i' f_{x_\tau} k l} \\ & \otimes \nabla_{f_1} \psi_1 \otimes \dots \otimes \nabla_{f_\tau} \phi_\tau \nabla_i \tilde{\phi}_1 \otimes \nabla_{i'} \tilde{\phi}_2), \end{aligned} \quad (2.125)$$

(where $\{x_1, \dots, x_\tau\} = \{1, \dots, \tau\}$), or in the form:

$$\begin{aligned} & p\text{contr}(\nabla_q Y \otimes \nabla^{(A) f_{x_1} \dots f_{x_{a-1}}} S_* R^{i f_{x_a} q l} \otimes \nabla^{(B) f_{x_{a+1}} \dots f_{x_{\tau-1}}} R^{i' f_{x_\tau} i_* l} \\ & \otimes \nabla_{y_1} \psi_1 \otimes \dots \otimes \nabla_{y_\tau} \phi_\tau \nabla_i \tilde{\phi}_1 \otimes \nabla_{i'} \tilde{\phi}_2). \end{aligned} \quad (2.126)$$

If $v = 1$ the equation (2.122) will hold with $P = \emptyset$:

Then, picking out the sublinear combination in (2.122) which consists of terms with a factor ∇Y and replacing into our hypothesis, we derive that the coefficient of each of the tensor fields indexed in P must be zero. This completes the proof of our claim. \square

2.5 Proof of Lemmas 2.2, 2.4:

Proof of Lemma 2.2:

The first claim follows immediately, since each tensor field has a removable index (thus each tensor field separately can be written as an $X_* \text{div}$).

The proof of the second claim essentially follows the “manual” construction of divergences, as in [5]. By “manually” constructing explicit divergences out of each $C_g^{h, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$, $h \in H_2$, we derive that we can write:

$$\begin{aligned} & \sum_{h \in H_2} a_h X \text{div}_{i_1} \dots X \text{div}_{i_\alpha} C_g^{h, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) = \\ & (Const)_1 X \text{div}_{i_1} \dots X \text{div}_{i_\epsilon} C_g^{1, i_1 \dots i_\epsilon}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & (Const)_2 X \text{div}_{i_1} \dots X \text{div}_{i_\zeta} C_g^{2, i_1 \dots i_\zeta}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & \sum_{q \in Q} a_q X \text{div}_{i_1} \dots X \text{div}_{i_\alpha} C_g^{q, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u), \end{aligned} \quad (2.127)$$

where the tensor fields indexed in Q are as required by our Lemma hypothesis, while the tensor fields C^1, C^2 are explicit tensor fields which we will write out below (they depend on the values p, σ_1, σ_2).⁶⁶

⁶⁶In some cases there will be no tensor fields C^1, C^2 (in which case we will just say that in (2.127) we have $(Const)_1 = 0, (Const)_2 = 0$).

We will then show that in (2.127) we will have $(Const)_1 = (Const)_2 = 0$. That will complete the proof of Lemma 2.2. We distinguish cases based on the value of p : Either $p = 2$ or $p = 1$ or $p = 0$.

The case $p = 2$: With no loss of generality we assume that the factor $\nabla^{(A)}\Omega_1$ is contracting against the factors $\nabla\phi_1, \dots, \nabla\phi_x$ and $\nabla^{(B)}\Omega_2$ is contracting against $\nabla\phi_{x+1}, \dots, \nabla\phi_{x+t}$; we may also assume wlog that $x \leq t$. By manually constructing divergences, it follows that we can derive (2.127), where each of the tensor fields C^1, C^2 will be in the forms, respectively:

$$pcontr(\nabla_{i_*} Y \otimes \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(A)} \Omega_1 \otimes \nabla_{y_1 \dots y_t i_{\gamma+1} \dots i_{\gamma+\delta}}^{(B)} \Omega_2 \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.128)$$

(where if $t \geq 2$ then $\delta = 0$, otherwise $t + \delta = 2$), or:

$$pcontr(\nabla_q Y \otimes \nabla^q \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(A)} \Omega_1 \otimes \nabla_{y_1 \dots y_t i_{\gamma+1} \dots i_{\gamma+\delta}}^{(B)} \Omega_2 \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.129)$$

(where if $t \geq 2$ then $\delta = 0$, otherwise $t + \delta = 2$).

The case $p = 1$: We “manually” construct divergences to derive (2.127), where if $\sigma_1 = 1$ then there are no tensor fields C^1, C^2 (and hence (2.127) is our desired conclusion); if $\sigma_1 = 0, \sigma_2 = 1$ then there is only the tensor field C^1 in (2.127) and it is in the form:

$$pcontr(\nabla^q Y \otimes S_* \nabla_{v_2 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} q} \otimes \nabla_{y_1 \dots y_t i_{\gamma+1} \dots i_{\gamma+\delta}}^{(B)} \Omega_2 \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{v_1} \phi_2 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.130)$$

where if $t \geq 2$ then $\delta = 0$, otherwise $\delta = 2 - t$.

The case $p = 0$: We have three subcases: Firstly $\sigma_2 = 2$, secondly ($\sigma_2 = 1, \sigma_1 = 1$), and thirdly $\sigma_1 = 2$.

In the case $\sigma_2 = 2$, the tensor fields C^1, C^2 must be in the forms, respectively:

$$pcontr(\nabla_{i_*} Y \otimes S_* \nabla_{v_2 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes S_* \nabla_{y_1 \dots y_t}^{(t-1)} R_{i' i_{\gamma+3} i_{\gamma+4}}{}^l \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{i'} \tilde{\phi}_2 \otimes \nabla^{v_1} \phi_3 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.131)$$

$$pcontr(\nabla^q Y \otimes S_* \nabla_{qv_2 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes S_* \nabla_{y_1 \dots y_t}^{(t-1)} R_{i' i_{\gamma+3} i_{\gamma+4}}{}^l \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{i'} \tilde{\phi}_2 \otimes \nabla^{v_1} \phi_3 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.132)$$

(if $x = t = 0$ then the tensor field C^1 above will not be present).

In the case $\sigma_1 = 2$, the tensor fields C^1, C^2 must be in one of the two forms:

$$p\text{contr}(\nabla_{i_*} Y \otimes \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(m_1)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes \nabla_{y_1 \dots y_t}^{(t-1)} R_{i_{\gamma+3} i_{\gamma+4}}^i \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.133)$$

$$p\text{contr}(\nabla_q Y \otimes \nabla^q \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(m_1)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes \nabla_{y_1 \dots y_t}^{(t-1)} R_{i_{\gamma+3} i_{\gamma+4}}^i \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u). \quad (2.134)$$

In the case $\sigma_1 = 1, \sigma_2 = 1$, there will be only one tensor field C^1 , in the form:

$$p\text{contr}(\nabla^q Y \otimes S_* \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(m_1)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes \nabla_{y_1 \dots y_t}^{(t-1)} R_{q i_{\gamma+3} i_{\gamma+4}}^i \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{v_1} \phi_2 \otimes \dots \otimes \nabla^{y_t} \phi_u). \quad (2.135)$$

We then derive that $(Const)_1 = (Const)_2 = 0$ as in [5] (by picking out the sublinear combination in (2.127) that consists of complete contractions with a factor ∇Y -differentiated only once).

Proof of Lemma 2.4:

We again “manually” construct explicit $X \text{div}'$ to write:

$$\begin{aligned} & \sum_{h \in H_2} a_h X \text{div}_{i_1} \dots X \text{div}_{i_\alpha} C_g^{h, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) = \\ & (Const)_1 X \text{div}_{i_1} \dots X \text{div}_{i_\xi} C_g^{1, i_1 \dots i_\xi}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & (Const)_2 X \text{div}_{i_1} \dots X \text{div}_{i_\zeta} C_g^{2, i_1 \dots i_\zeta}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & \sum_{q \in Q} a_q X \text{div}_{i_1} \dots X \text{div}_{i_\alpha} C_g^{q, i_1 \dots i_\alpha}(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u) + \\ & \sum_{j \in J} a_j C_g^j(\Omega_1, \dots, \Omega_p, Y, \phi_1, \dots, \phi_u), \end{aligned} \quad (2.136)$$

where the tensor fields indexed in Q are as required by our Lemma hypothesis, while the tensor fields C^1, C^2 are explicit tensor fields which we will write out below (they depend on the values p, σ_1, σ_2). In some cases there will be no tensor fields C^1, C^2 (in which case we will just say that in (2.127) we have $(Const)_1 = 0, (Const)_2 = 0$).

The case $p = 2$: With no loss of generality we assume that the factor $\nabla^{(A)} \Omega_1$ is contracting against the factors $\nabla \phi_1, \dots, \nabla \phi_x$ and $\nabla^{(B)} \Omega_2$ is contracting against $\nabla \phi_{x+1}, \dots, \nabla \phi_{x+t}$; we may also assume wlog that $x \leq t$. By manual construction of divergences, it follows that we can derive (2.127), where there is only the tensor field C^1 and it is in the form:

$$p\text{contr}(\nabla_{[i_*} \chi_1 \otimes \nabla^q] \chi_2 \otimes \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(A)} \Omega_1 \otimes \nabla_{qy_1 \dots y_t i_{\gamma+1} \dots i_{\gamma+\delta}}^{(B)} \Omega_2 \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.137)$$

(where if $t \geq 1$ then $\delta = 0$, otherwise $\delta = 1$).

The case $p = 1$: We “manually” construct divergences to derive (2.136), where: if $\sigma_1 = 1$ then there are no tensor fields C^1, C^2 in the RHS of (2.136) (and this is our desired conclusion); if $\sigma_1 = 0, \sigma_2 = 1$ then there is only the tensor field C^1 in the RHS of (2.136) and it is of the form:

$$p\text{contr}(\nabla_{[i_*} \omega_1 \otimes \nabla^q] \omega_2 \otimes S_* \nabla_{v_2 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} q} \otimes \nabla_{y_1 \dots y_t i_{\gamma+1} \dots i_{\gamma+\delta}}^{(B)} \Omega_2 \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{v_1} \phi_2 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.138)$$

where if $t \geq 2$ then $\delta = 0$, otherwise $\delta = 2 - t$.

The case $p = 0$: We have three subcases: Firstly $\sigma_2 = 2$, secondly ($\sigma_2 = 1, \sigma_1 = 1$), and thirdly $\sigma_1 = 2$.

In the case $\sigma_2 = 2$, the tensor fields C^1, C^2 in the RHS of (2.127) will be in the two forms, respectively:

$$p\text{contr}(\nabla_{[i_*} \omega_1 \otimes \nabla^q] \omega_2 \otimes S_* \nabla_{qv_2 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes S_* \nabla_{y_1 \dots y_t}^{(t-1)} R_{i' i_{\gamma+3} i_{\gamma+4}}{}^l \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{i'} \tilde{\phi}_2 \otimes \nabla^{v_1} \phi_3 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.139)$$

$$p\text{contr}(\nabla_{[p} \omega_1 \otimes \nabla^q] \omega_2 \otimes S_* \nabla_{qv_2 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} p} \otimes S_* \nabla_{y_1 \dots y_t}^{(t-1)} R_{i' i_{\gamma+3} i_{\gamma+4} q} \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{i'} \tilde{\phi}_2 \otimes \nabla^{v_1} \phi_3 \otimes \dots \otimes \nabla^{y_t} \phi_u). \quad (2.140)$$

In the case $\sigma_1 = 2$, the tensor fields C^1, C^2 will be the forms, respectively:

$$p\text{contr}(\nabla_{[i_*} \omega_1 \otimes \nabla^q] \otimes \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(m_1)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes \nabla_{qy_1 \dots y_t}^{(t-1)} R^i{}_{i_{\gamma+3} i_{\gamma+4}}{}^l \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.141)$$

$$p\text{contr}(\nabla_{[p} \omega_1 \otimes \nabla^q] \omega_2 \otimes \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(m_1)} R_{ii_{\gamma+1} i_{\gamma+2} p} \otimes \nabla_{y_1 \dots y_t}^{(t-1)} R^i{}_{i_{\gamma+3} i_{\gamma+4} q} \otimes \nabla^{v_1} \phi_1 \otimes \dots \otimes \nabla^{y_t} \phi_u), \quad (2.142)$$

(if at least one of the two factors $\nabla^{(m)} R_{ijkl}$ is contracting against a factor $\nabla \phi_h$. Otherwise, we can prove (2.136) with no tensor fields C^1, C^2 on the RHS).

In the case $\sigma_1 = 1, \sigma_2 = 1$, the tensor fields C^1, C^2 must be in the forms, respectively:

$$\begin{aligned}
& p\text{contr}(\nabla_{[i_*} \omega_1 \otimes \nabla^q] \omega_2 \otimes S_* \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(\nu)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes \nabla_{y_1 \dots y_t}^{(t-1)} R_{qi_{\gamma+3} i_{\gamma+4}}{}^l \\
& \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{v_1} \phi_2 \otimes \dots \otimes \nabla^{y_t} \phi_u),
\end{aligned} \tag{2.143}$$

$$\begin{aligned}
& p\text{contr}(\nabla_{[p} \omega_1 \otimes \nabla^q] \omega_2 \otimes S_* \nabla_{v_1 \dots v_x i_1 \dots i_\gamma}^{(m_1)} R_{ii_{\gamma+1} i_{\gamma+2} l} \otimes \nabla_{y_1 \dots y_t}^{(t-1)} R_{pq i_{\gamma+3}}{}^l \\
& \otimes \nabla^i \tilde{\phi}_1 \otimes \nabla^{v_1} \phi_2 \otimes \dots \otimes \nabla^{y_t} \phi_u).
\end{aligned} \tag{2.144}$$

We then derive that $(Const)_1 = (Const)_2 = 0$ by picking out the sublinear combination in (2.136) that consists of complete contractions with two factors $\nabla Y, \nabla \omega_2$ —each factor differentiated only once). \square

3 The proof of Proposition 1.1 in the special cases:

3.1 The direct proof of Proposition 1.1 (in case II) in the “special cases”.

We now prove Proposition 1.1 directly in the special subcases of case II. We recall the setting of the special subcases of Proposition 1.1 in case II are as follows: In subcase IIA for each μ -tensor field of maximal refined double character, $C_g^{l, i_1 \dots i_\mu}$ there is a unique factor in the form $T = \nabla^{(m)} R_{ijkl}$ for which two internal indices are free, and each derivative index is either free or contracting against a factor $\nabla \phi_h$. For subcase IIB there is a unique factor in the form $T = \nabla^{(m)} R_{ijkl}$ for which one internal index is free, and each derivative index is either free or contracting against a factor $\nabla \phi_h$. In both subcases IIA, IIB there is at least one free derivative index in the factor T .

Moreover, both in subcases IIA, IIB, all *other* real factors in $C_g^{l, i_1 \dots i_\mu}$ are either in the form $S_* R_{ijkl}$ or $\nabla^{(2)} \Omega_h$, or they are in the form $\nabla^{(m)} R_{ijkl}$, where all the m derivative indices contract against factors $\nabla \phi_h$.⁶⁷

In order to prove Proposition 1.1 directly in the special subcases of subcases IIA, IIB we will rely on a new Lemma:

Our new Lemma deals with two different settings, which we will label setting A and setting B below.

In setting A, we let

$$\sum_{l \in L} a_l C_g^{l, i_1 \dots i_\mu} (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$$

⁶⁷For the rest of this subsection, we will slightly abuse notation and *not* write out the derivative indices that contract against factors $\nabla \phi_h$ —we will thus refer to factors R_{ijkl} , setting $m = 0$.

stand for a linear combination of μ -tensor fields with one factor $\nabla^{(m)}R_{ijkl}$ containing $\alpha \geq 2$ free indices, distributed according to the pattern $\nabla_{(free)\dots(free)}^{(m)}R_{(free)j(free)l}$, and all other other factors being all in one of the forms $R_{ijkl}, S_*R_{ijkl}, \nabla^{(2)}\Omega_h$. (I.e. they have no removable indices).

In setting B we let

$$\sum_{l \in \bar{L}} a_l C_g^{l, i_1 \dots i_\mu} (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$$

stand for a linear combination of μ -tensor fields with one factor $\nabla^{(m)}R_{ijkl}$ containing $\alpha \geq 2$ free indices, distributed according to the pattern $\nabla_{(free)\dots(free)}^{(m)}R_{(free)j(free)l}$, and all *but one* of the other factors being in one of the forms $R_{ijkl}, S_*R_{ijkl}, \nabla^{(2)}\Omega_h$; one of the other factors (which we label T') will be in the form $\nabla R_{ijkl}, S_*\nabla R_{ijkl}, \nabla^{(3)}\Omega_h$. We will call this other factor “the factor with the extra derivative”. Moreover, in setting B we impose the additional restriction that if both the indices j, l in the factor $\nabla_{(free)\dots(free)}^{(m)}R_{(free)j(free)l}$ contract against the same other factor T' , then either T' is *not* the factor with the extra derivative, or if it is, then T' is in the form $\nabla_s R_{abcd}$, and furthermore the indices j, l contract against the indices b, c *and* we assume that the indices s, a, c are symmetrized over.⁶⁸

Lemma 3.1 *Let $\sum_{l \in \bar{L}} a_l C_g^{l, i_1 \dots i_\mu}$ be a linear combination of μ -tensor fields as described above. We assume the following special case of (1.7):*

$$\begin{aligned} & \sum_{l \in \bar{L} \cup L'} a_l X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_\mu} C_g^{l, i_1 \dots i_\mu} (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) + \\ & \sum_{h \in H} a_h X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_\mu} C_g^{h, i_1 \dots i_\beta} (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u) \\ & + \sum_{j \in J} a_j C_g^j (\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u); \end{aligned} \quad (3.1)$$

here, in both cases A and B the terms indexed in \bar{L} will be as described above; the μ -tensor fields indexed in L' will have fewer than α free indices in any given factor of the form $\nabla^{(m)}R_{ijkl}$. The tensor fields indexed in H each have rank $> \mu$ and also each of them has fewer than α free indices in any given factor of the form $\nabla^{(m)}R_{ijkl}$. Finally, the terms indexed in J are simply subsequent to \vec{K}_{simp} .

We claim that:

$$\sum_{l \in \bar{L}} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v = 0. \quad (3.2)$$

⁶⁸In other words, in that case the factors T, T' contract according to the pattern: $\nabla_{(free)\dots(free)}^{(m)}R_{(free)j(free)l} \nabla_{(s} R_{a^j k}{}^d)$, where the indices s, a, d are symmetrized over.

We will prove this Lemma shortly. Let us now, however, note how the above Lemma directly implies Proposition 1.1 in the special subcases IIA (directly) and IIB (after some manipulation).

Lemma 3.1 implies Proposition 1.1 in the special subcases of case II:

We first start with subcase IIA: Consider the sublinear combination of μ -tensor fields of maximal refined double character in (1.7). Denote their index set by $L_{Max} \subset L$. Recall that since we are considering the subcase where (1.7) falls under the special case of Proposition 1.1 in case IIA, it follows that for each $C_g^{l, i_1 \dots i_\mu}$ there is a unique factor in the form $\nabla^{(m)} R_{ijkl}$ for which two internal indices are free, and each derivative index is either free or contracting against a factor $\nabla \phi_h$; denote by $M + 2$ the number of free indices in that factor.⁶⁹

Now, by weight considerations (since we are in a special subcase of Proposition 1.1 in case IIA), any tensor field of rank $> \mu$ in (1.7) must have strictly fewer than $M + 2$ free indices in any given factor $\nabla^{(m)} R_{ijkl}$. Therefore in subcase IA, (1.7) is of the form (3.1), with $L_{Max} \subset \bar{L}$. Therefore, we apply Lemma 3.1 to (1.7) and pick out the sublinear combination of terms with a refined double character $Doub(\bar{L}^z), z \in Z'_{Max}$ ⁷⁰ we thus obtain a *new* true equation, since (3.2) holds formally, and the double character is invariant under the formal permutations of indices that make (3.2) formally zero. This proves our claim in subcase IIA.

Now we deal with subcase IIB:

We consider the μ -tensor fields of maximal refined double character in (1.7). By definition (since we now fall under a special case), they will each have a factor in the form $\nabla_{(free) \dots (free)}^{(m)} R_{(free)ijkl}$, with a total of $M + 1 > 1$ free indices.⁷¹ Each of the other factors will be in the form R_{ijkl} or be simple factors in the form $S_* R_{ijkl}$, or in the form $\nabla^{(2)} \Omega_h$.

We denote by $\bar{L} \subset L$ the index set of μ -tensor fields with $M + 1$ free indices in a factor $\nabla^{(m)} R_{ijkl}$. It follows by weight considerations that the factor in question will be unique for each $C_g^{l, i_1 \dots i_\mu}, l \in \bar{L}$. We then start out with some explicit manipulation of the terms indexed in \bar{L} :

We will prove that there exists a linear combination of $\mu + 1$ -tensor fields, $\sum_{h \in H} a_h C_g^{h, i_1 \dots i_{\mu+1}}$, as allowed in the statement of Proposition 1.1, so that:

$$\begin{aligned} \sum_{l \in \bar{L}} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v &= \sum_{h \in H} a_h X div_{i_{\mu+1}} C_g^{h, i_1 \dots i_{\mu+1}} \nabla_{i_1} v \dots \nabla_{i_\mu} v \\ &+ \sum_{l \in \bar{L}_{new}} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v \sum_{j \in J} a_j C_g^{j, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v. \end{aligned} \quad (3.3)$$

⁶⁹So we set $\alpha = M + 2$.

⁷⁰Recall that $\bar{L}^z, z \in Z'_{Max}$ is the collection of maximal refined double characters that Proposition 1.1 deals with.

⁷¹So, we set $\alpha = M + 1$.

Here the μ -tensor fields indexed in \bar{L}_{new} have a factor

$T = \nabla_{(free)\dots(free)}^{(M-1)} R_{(free)j(free)l}$, and *one other factor* T' has an extra derivative (meaning that T' is either in the form ∇R_{ijkl} or $S_* \nabla R_{ijkl}$, or $\nabla^{(3)} \Omega_h$). Moreover if both indices j, l in T contract against indices j, l in the same factor T'' and at least one of j, l is removable, then $T' \neq T''$. Clearly, (3.3) in conjunction with Lemma 3.1 implies Proposition 1.1 in the “special cases” of case II. So, matters are reduced to showing (3.3) (and then deriving Lemma 3.1).

Proof of (3.3): We first apply the second Bianchi identity to the factor T to move one of the derivative free indices into the position kl in the factor $\nabla_{(free)\dots(free)}^{(M-1)} R_{(free)j(free)l}$. Thus, we derive that modulo terms of length $\geq \sigma + u + 1$:

$$C_g^{l, i_1 \dots i_\mu} = -C_g^{l, i_1 \dots i_\mu} + C_g^{l, 2, i_1 \dots i_\mu},$$

where the partial contractions $C_g^{l, 1, i_1 \dots i_\mu}, C_g^{l, 2, i_1 \dots i_\mu}$ have the factor T replaced by a factor in the form: $\nabla_{k(free)\dots(free)}^{(m)} R_{(free)j(free)l}, \nabla_{l(free)\dots(free)}^{(m)} R_{(free)jk(free)}$, respectively. We then erase the indices k, l in these two factors (thus creating a new tensor field $C_g^{l, 1, i_1 \dots, i_\mu, i_{\mu+1}}, C_g^{l, 2, i_1 \dots, i_\mu, i_{\mu+1}}$) by creating a free index $i_{\mu+1}$, and subtract the $Xdiv_{i_{\mu+1}}[\dots]$ of the corresponding $(\mu + 1)$ -tensor field. We then derive an equation:

$$C_g^{l, i_1 \dots i_\mu} = Xdiv_{i_{\mu+1}} C_g^{l, i_1 \dots i_{\mu+1}} + \sum_{l \in L_{new}} C_g^{l, i_1 \dots i_\mu}, \quad (3.4)$$

where all the tensor fields indexed in L_{new} satisfy the required property of Lemma 3.1, except for the fact that one could have both indices j, l in the factor $\nabla_{(free)\dots(free)}^{(M-1)} R_{(free)j(free)l}$ contracting against indices j, l in a factor T' which has an additional derivative index. If $C_g^{l, i_1 \dots i_\mu}, l \in L_{new}$ is not in the form allowed in the claim of Lemma 3.1, then (after possibly applying the second Bianchi identity and possibly introducing simply subsequent complete contractions) we may arrange that one of the indices j, l is a derivative index.

In that case we construct another $(\mu + 1)$ -tensor field by erasing the derivative index j or l and making the index j or l in a free index $i_{\mu+1}$. Then, subtracting the corresponding $Xdiv_{i_{\mu+1}}$ of this new $(\mu + 1)$ -tensor field, we derive our claim. \square

Therefore, matters are reduced to proving Lemma 3.1.

Proof of Lemma 3.1:

Let us start with some notational conventions

Recall the first variation law of the curvature tensor under variations by a symmetric 2-tensor by v_{ij} : For any complete or partial contraction $T(g_{ij})$ (which is a function of the metric g_{ij}), we define: $Image_{v_{ij}}^1 = \frac{d}{dt}|_{t=0} [T(g_{ij} + tv_{ij})]$. (We

write $Image_{v_{ij}}^1[\dots]$ or $Image_{v_{ab}}^1[\dots]$ below to stress that we are varying by a 2-tensor, rather than just by a scalar).

We consider the equation $Image_{v_{ij}}^1[L_g] = 0$ (which corresponds to the first *metric* variation of our Lemma hypothesis (i.e. of (1.7)). This equation holds modulo complete contractions with at least $\sigma + u + 1$ factors.

Thus, we derive a new local equation:

$$\begin{aligned} & \sum_{l \in L_\mu} a_l X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_\mu} Image_{v_{ab}}^1[C_g^{l, i_1 \dots i_\mu}] \\ & + \sum_{l \in L \setminus L_\mu} a_l X \operatorname{div}_{i_1} \dots X \operatorname{div}_{i_a} Image_{v_{ab}}^1[C_g^{l, i_1 \dots i_a}] \\ & = \sum_{j \in J} a_j Image_{v_{ab}}^1[C_g^j], \end{aligned} \quad (3.5)$$

which holds modulo terms of length $\geq \sigma + u + 1$.

Now, we wish to pass from the local equation above to an integral equation, and then to apply the *silly divergence formula* from [1] to that integral equation (thus deriving a *new* local equation).

In order to do this, we start by introducing some more notation: Let us write out:

$$Image_{v_{ab}}^1[C_g^{l, i_1 \dots i_\mu}] = \sum_{t \in T^l} a_t C_g^{t, i_1 \dots i_a}$$

where each $C_g^{t, i_1 \dots i_a}$ is in the form:

$$\begin{aligned} & p \operatorname{contr}(\nabla_{r_1 \dots r_{A+2}}^{(A+2)} v_{ab} \otimes \nabla^{(m_1)} R_{ijkl} \otimes \dots \otimes \nabla^{(m_{\sigma-1})} R \otimes \nabla^{(b_1)} \Omega_1 \otimes \dots \otimes \nabla^{(b_p)} \Omega_p \\ & \otimes \nabla \phi_1 \otimes \dots \otimes \nabla \phi_u). \end{aligned} \quad (3.6)$$

For our next technical tool we introduce some notation: For each tensor field $C_g^{l, i_1 \dots i_a}$ in the form above, we denote by C_g^l the complete contraction that arises by hitting each factor T_i ($i = 1, 2, 3$) by m derivative indices $\nabla^{u_1 \dots u_m}$, where u_1, \dots, u_m are the free indices that belong to T_i in $C_g^{l, i_1 \dots i_a}$ (thus we obtain a factor with m internal contraction, each involving a derivative index). Notice there is a one-to-one correspondence between the tensor fields and the complete contractions we are constructing. We can then easily observe that there are two linear combinations $\sum_{r \in R_1} a_r C_g^r(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$, $\sum_{r \in R_2} a_r C_g^r(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u)$ where each C_g^r , $r \in R_1$ has at least $\sigma + u + 1$ factors, while each C_g^r , $r \in R_2$ has $\sigma + u$ factors but at least one factor $\nabla^{(p)} \phi_h \neq \Delta \phi_h$ with $p \geq 2$, so that for any compact orientable (M, g) :

$$\int_M \sum_{l \in L} a_l \sum_{t \in T^l} a_t C_g^{t, *}(v_{ab}) + \sum_{r \in R_1} a_r C_g^r(v_{ab}) + \sum_{r \in R_2} a_r C_g^r(v_{ab}) dV_g = 0 \quad (3.7)$$

(denote the integrand of the above by $Z_g(v_{ab})$). Here again each C_g^j has $\sigma + u$ factors and all factors $\nabla\phi_h$ have only one derivative but its simple character is subsequent to $\vec{\kappa}$. We call this technique (of going from the local equation (3.5) to the integral equation (3.7)) the “inverse integration by parts”.

Now, we derive a “silly divergence formula” from the above by performing integrations by parts with respect to the factor $\nabla^{(B)}v_{ab}$ (until we are left with a factor v_{ab} —without derivatives). This produces a new local equation which we denote by $silly[Z_g(v_{ab})] = 0$. We will be using this equation in our derivation of Lemma 3.1.

Now, for each $C_g^{l,i_1\dots i_\mu}$, $l \in \bar{L}$, we consider the factor $T = \nabla_{(free)\dots(free)}^{(M)} R_{(free)j(free)l}$ with the $M+2$ free indices. We define T^j to be the factor in $C_g^{l,i_1\dots i_\mu}$ that contracts against the index j in T and by T^l to be the factor in $C_g^{l,i_1\dots i_\mu}$ that contracts against the index l in T . We define $\bar{L}_{same} \subset \bar{L}$ to be the index set of tensor fields for which $T^j = T^l$; we define $\bar{L}_{not.same} \subset \bar{L}$ to be the index set of tensor fields for which $T^j \neq T^l$. We will then prove (3.2) separately for the two sublinear combinations indexed in $\bar{L}_{same}, \bar{L}_{not.same}$.

Proof of (3.2) for the index set \bar{L}_{same} :

We first prove our claim for $\sigma > 3$ and then note how to prove it when $\sigma = 3$.

Consider $silly[L_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = 0$. Pick out the sublinear combination $silly_+[L_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = 0$ with $\mu - M - 2$ internal contractions, and with the indices in the factor v_{ab} contracting against a factor T' which either has *no* extra derivative indices, or if it does, then the contraction is according to the pattern $v^{ab} \otimes \nabla_s R_{ajbl}$; we also require that the two factors T'', T''' with an extra $M+2$ extra derivatives each. This sublinear combination must vanish separately, hence we derive:

$$silly_+[Z_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = 0. \quad (3.8)$$

We also observe that this sublinear combination can only arise (in the process of passing from the equation $L_g = 0$ to deriving $silly_+[Z_g(v_{ab})] = 0$) by replacing the factor $\nabla_{(free)\dots(free)}^{(M)} R_{(free)j(free)l}$ by $\nabla_{(free)\dots(free)}^{(M)} v_{jl}$ and then (in the inverse integration by parts) replacing all μ free indices by internal contractions,⁷² and finally integrating by parts the $M+2$ pairs of derivative indices (∇^a, ∇_a) and forcing all upper indices hit a factor $T'' \neq T'$ and the lower indices to hit a factor $T''' \neq T', T''' \neq T''$.⁷³

Thus, we can prove our claim by starting from the equation (3.8) and applying Sub_v $\mu - M - 2$ times,⁷⁴ just applying the eraser to the extra $M+2$ pairs of contracting derivatives,⁷⁵ and then replacing the factor v_{ab} by

⁷²(Thus the factor $\nabla_{(free)\dots(free)}^{(M)} v_{jl}$ gets replaced by $\Delta^{M+2} v_{ij}$).

⁷³The fact that $\sigma > 3$ ensures the existence of two such factors.

⁷⁴See the Appendix in [3] and just set $\omega = v$.

⁷⁵This can be done by just repeating the proof of the “Eraser” Lemma in the Appendix of [3].

$\nabla_{r_1 \dots r_M}^{(M)} R_{iajb} \nabla^{r_1} v \dots \nabla^{r_M} v \nabla^a v \nabla^b v$. Finally we just divide by the combinatorial constant $\binom{\sigma-3}{2}$.

Let us now consider the case $\sigma = 3$: In those case the terms of maximal refined double character can only arise in the subcase IIA,⁷⁶ and can only be in one of the forms: $\nabla_{(free) \dots (free)}^{(M)} R_{(free)j(free)l} \otimes R^{ijkl} \otimes \nabla_{ik}^{(2)} \Omega_1$, $\nabla_{(free) \dots (free)}^{(M)} R_{(free)j(free)l} \otimes R^{ijkl} \otimes R_{(free)}^j \binom{l}{(free)}$. Thus, in that case we define $silly_+[Z_g(v_{ab})]$ to stand for the terms $(v_{jl} \otimes \nabla_{t_1 \dots t_{M+2}}^{(M+2)} R^{ijkl} \otimes \nabla_{t_1 \dots t_{M+4} ik}^{(M+4)} \Omega_1)$, $(v_{jl} \otimes \nabla_{t_1 \dots t_{M+2}}^{(M+2)} R^{ijkl} (\nabla^{(M+2)})^{t_1 \dots t_{M+2}} \otimes R_{(free)}^j \binom{l}{(free)})$ respectively, and then repeat the argument above.

Proof of (3.2) for the index set $\bar{L}_{not.same}$:

We prove our claim in steps: We first denote by $\bar{L}_{not.same}^{**} \subset \bar{L}_{not.same}$ the index set of tensor fields in $\bar{L}_{not.same}$ for which both indices j, l in the factor $T = \nabla_{(free) \dots (free)}^{(M)} R_{(free)j(free)l}$ contract against special indices in factors T^j, T^l of the form $S_* R_{ijkl}$. We will firstly prove that:

$$\sum_{l \in \bar{L}_{not.same}^{**}} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v = \sum_{l \in L'} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v. \quad (3.9)$$

Here the terms in the RHS have all the features of the terms in $\bar{L}_{not.same}$, but in addition at most one of the indices in the factor $T = \nabla_{(free) \dots (free)}^{(M)} R_{(free)j(free)l}$ contract against a special index in a factor of the form $S_* R_{ijkl}$. Thus, if we can prove (3.9), we are reduced to proving our claim under the additional assumption that $\bar{L}_{not.same}^{**} = \emptyset$.

For our next claim, we denote by $\bar{L}_{not.same}^* \subset \bar{L}_{not.same}$ the index set of tensor fields in $\bar{L}_{not.same}$ for which one of the indices j, l in the factor $T = \nabla_{(free) \dots (free)}^{(M)} R_{(free)j(free)l}$ contracts against a special index in factors T^j, T^l of the form $S_* R_{ijkl}$.

We will then prove that:

$$\sum_{l \in \bar{L}_{not.same}^*} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v = \sum_{l \in L''} a_l C_g^{l, i_1 \dots i_\mu} \nabla_{i_1} v \dots \nabla_{i_\mu} v. \quad (3.10)$$

Here the terms in the RHS have all the features of the terms in $\bar{L}_{not.same}$, but in addition none of the indices in the factor $T = \nabla_{(free) \dots (free)}^{(M)} R_{(free)j(free)l}$ contracting against a special index in a factors of the form $S_* R_{ijkl}$. Thus, if we can prove (3.9), we are reduced to proving our claim under the additional assumption that for eac $C_g^{l, i_1 \dots i_\mu}$, $l \in \bar{L}$ the two indices j, l in the factor $T =$

⁷⁶This follows by virtue of the symmetry of the indices s, a, d in any factor $\nabla_s R_{abcd}$ as discussed above.

$\nabla_{(free)\dots(free)}^{(M)} R_{(free)j(free)l}$ contract against two different factors and none of the indices j, l are special indices in a factor of the form $S_* R_{ijkl}$.

In our third step, we prove (3.2) under this additonnal assumption.

We now present our proof of the third step. We will indicate in the end how this proof can be easily modified to derive the first two steps.

For each $l \in \bar{L}_{not.same}$, let us denote by $link(l)$ the number of particular contractions between the factors T^j, T^l in the tensor fields $C_g^{l, i_1 \dots i_\mu}$. (Note that by weight considerations $0 \leq link(l) \leq 3$). Let B be the maximum value of $link(l), l \in \bar{L}_{not.same}$, and by $\bar{L}_{not.same}^B \subset \bar{L}_{not.same}$ the corresponding index set. We will then prove our claim for the tensor fields indexed in $\bar{L}_{not.same}^B$. By repeating this step at most four times, we will derive our third claim.

Consider $silly[L_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = 0$. Pick out the sublinear combination $silly_*[L_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = 0$ with $\mu - M - 2$ internal contractions, and with an extra $M + 2$ derivatives on the factors T^j, T^l against which the two indices of the factor v_{ab} contract, *and* with $M + 2 + B$ partical contractions between the factors T^j, T^l . This sublinear combination must vanish separately:

$$silly_*[L_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = 0.$$

Moreover, we observe by following the ‘‘inverse integration by parts’’ and the silly divergence formula obtained from $\int_{M^n} Z_g(v_{ab}) dV_g = 0$, that the LHS of the above can be described as follows:

For each $C_g^{l, i_1 \dots i_\mu}, l \in \bar{L}_{not.same}^B$, we denote by $\tilde{C}_g^l(v_{ab})$ the complete contraction that arises by replacing the factor $T = \nabla_{(free)\dots(free)}^{(M)} R_{(free)j(free)l}$ by $\nabla_{(free)\dots(free)}^{(M+2)} v_{jl}$, and then replacing each free index that *does not* belong to the factor T by an internal contraction. We then denote by $\hat{C}_g^l(v_{ab})$ the complete contraction that rises from $\tilde{C}_g^l(v_{ab})$ by hitting the factor T^j (against which the index j in v_{jl} contracts) by $(M + 2)$ derivative indices $\nabla_{t_1}, \dots, \nabla_{t_{M+2}}$ and hitting the factor T^l (against which the index l in v_{jl} contracts) by derivatives $\nabla^{t_1}, \dots, \nabla^{t_{M+2}}$.⁷⁷ It follows that:

$$(0 =) silly_*[L_g(\Omega_1, \dots, \Omega_p, \phi_1, \dots, \phi_u, v_{ab})] = \sum_{\bar{L}_{not.same}^B} a_l 2^{M+1} [\hat{C}_g^l(v_{ab})].$$

Now, to derive our claim, we introduce a formal operation $Op[\dots]$ which acts on the terms above by applying Sub_v to each of the $\mu - M - 2$ internal contractions,⁷⁸ erasing $M + 2$ particular contractions between the factors T^j, T^l and then replacing the factor v_{jl} by $\nabla_{r_1 \dots r_M}^{(m)} R_{ijkl} \nabla^{r_1} v \dots \nabla^{r_M} v \nabla^i v \nabla^k v$. This operation produces a new true equation; after we divide this new true equation by 2^{M+1} , we derive our claim.

⁷⁷These derivatives contract against the indices $\nabla_{t_1}, \dots, \nabla_{t_{M+2}}$ that have hit T^j .

⁷⁸See the Appendix of [3] for the definition of this operation.

Note on the derivation of (3.9), (3.10): The equations can be derived by a straightforward modification of the ideas above: The only extra feature we must add is that in the silly divergence formula we must pick out the terms for which (both/one of the) indices j, l in v_{jl} contract against a special index in a factor $S_* \nabla^{(M+2)} R_{abcd} \nabla^a \tilde{\phi}_h$. This linear combination will vanish, modulo terms where one/none of the indices j, l in v_{jl} contract against a special index in the factor $S_* R_{ijkl}$: This follows by the same argument that is used in [6] to derive that Lemma 3.1 in [6] implies Proposition 1.1 in case I: We firstly replace the factor v_{jl} by an expression $y_{(j} y_{l)}$. We then just replace both/one of the expressions $\nabla_i \tilde{\phi}_h, y_j$ by g_{ij} and apply *Ricto* Ω twice/once.⁷⁹ The only terms that survive this true equation are the ones indexed in $\bar{L}_{not.same}$, for which the expression(s) $S_* \nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_h \nabla^k y$ are replaced by $\nabla_{r_1 \dots r_\nu}^{(\nu+2)} Y_f$. We then proceed as above, deriving that the sublinear combination of terms indexed in $\bar{L}_{not.same}$ must vanish, *after* we replace two/one expressions $S_* \nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_h \nabla^k y$ by $\nabla_{r_1 \dots r_\nu}^{(\nu+2)} Y_f$. Then, repeating the permutations applied to any factors $\nabla_{r_1 \dots r_\nu}^{(\nu+2)} Y_f$, to $S_* \nabla_{r_1 \dots r_\nu}^{(\nu)} R_{ijkl} \nabla^i \tilde{\phi}_h \nabla^k y$ we derive our claim. \square

3.2 The remaining cases of Proposition 1.1 in case III:

We recall that there are remaining cases only when $\sigma = 3$. In that case we have the remaining cases when $p = 3$ and $n - 2u - 2\mu \leq 2$, or when $p = 2, \sigma_2 = 1$ and $n = 2u + 2\mu$.

The case $p = 3$: Let us start with the subcase $n - 2u - 2\mu = 0$. In this case, all tensor fields in (1.7) will be in the form:

$$\begin{aligned}
& p \text{contr}(\nabla_{i_1 \dots i_a j_1 \dots j_b}^{(A)} \Omega_1 \otimes \nabla_{i_{a+1} \dots i_{a+a'} j_{b+1} \dots j_{b+b'}}^{(B)} \Omega_2 \otimes \\
& \nabla_{i_{a+a'+1} \dots i_{a+a'+a''} j_{b+b'+1} \dots j_{b+b'+b''}}^{(C)} \Omega_3 \otimes \nabla^{j_{x_1}} \phi_1 \dots \otimes \nabla^{x_{j+j'+j''}} \phi_u)
\end{aligned} \tag{3.11}$$

where we are making the following conventions: Each of the indices i_f is free; also, each of the indices j_f is contracting against some factor $\nabla \phi_h$, and also $A, B, C \geq 2$.

Thus, we observe that in this subcase μ is also the *maximum* rank among the tensor fields appearing in (1.7). Now, assume that the μ -tensor fields in (1.7) of *maximal refined double character* have $a = \alpha, a' = \alpha', a'' = \alpha''$. With no loss of generality (only up to renaming the factors $\Omega_1, \Omega_2, \Omega_3, \phi_1, \dots, \phi_u$) we may assume that $\alpha \geq \alpha' \geq \alpha''$ and that only the functions $\nabla \phi_1, \dots, \nabla \phi_{u_1}$ contract against $\nabla^{(A)} \Omega_1$ in $\vec{\mathcal{F}}_{simp}$. We will then show that the coefficient $a_{\alpha, \alpha', \alpha''}$ of this tensor field must be zero. This will prove Proposition 1.1 in this subcase.

We prove that $a_{\alpha, \alpha', \alpha''} = 0$ by considering the global equation $\int Z_g dV_g = 0$ and considering the silly divergence formula $silly[Z_g] = 0$. We then consider

⁷⁹Recall that this operation has been defined in the Appendix of [3] and produces a true equation.

the sublinear combination $silly_+[Z_g]$ consisting of terms with α', α'' internal contractions in the factors $\nabla^{(D)}\Omega_2, \nabla^{(E)}\Omega_3$, with α particular contractions between those factors and with all factors $\nabla\phi_h$ that contracted against $\nabla^{(A)}\Omega_1$ in $\vec{\kappa}_{simp}$ being replaced by $\Delta\phi_h$, while all factors $\nabla\phi_h$ that contracted against $\nabla^{(B)}\Omega_2, \nabla^{(C)}\Omega_3$ still do so. We easily observe that $silly_+[Z_g] = 0$, and furthermore $silly_+[Z_g]$ consists of the complete contraction:

$$\begin{aligned} & \text{contr}(\Omega_1 \otimes \nabla^{f_1 \dots f_\alpha}_{j_{b+1} \dots j_{b+b'}} \Delta^{\alpha'} \Omega_2 \otimes \nabla_{f_1 \dots f_\alpha j_{b+b'+1} \dots j_{b+b'+b''}} \Delta^{\alpha''} \Omega_3 \otimes \Delta\phi_1 \\ & \dots \nabla^{j_{x_{b+b'+b''}}} \phi_u) \end{aligned} \quad (3.12)$$

times the constant $(-1)^{u_1} 2^\alpha a_{\alpha, \alpha', \alpha''}$. Thus, we derive that $a_{\alpha, \alpha', \alpha''} = 0$.

The second subcase: We now consider the setting where $\sigma = p = 3$, $n - 2u - 2\mu = 2$. In this setting, the maximum rank of the tensor fields appearing in (1.7) is $\mu + 1$. In this case, all $(\mu + 1)$ -tensor fields in (1.7) will be in the form (3.11) (with $\alpha + \alpha' + \alpha'' = \mu + 1$, while all the μ -tensor fields will be in the form (3.11) but with $\alpha + \alpha' + \alpha'' = \mu$, and with one particular contraction c, c between two of the factors $\nabla^{(A)}\Omega_1, \nabla^{(B)}\Omega_2, \nabla^{(C)}\Omega_3$.

Now, if both the indices c, c described above are removable, we can explicitly express $C_g^{l, i_1 \dots i_\mu}$ as an $Xdiv$ of an acceptable $(\mu + 1)$ -tensor field. Therefore, we are reduced to showing our claim in this setting where for each μ -tensor field in (1.7) at least one of the indices c, c is not removable. Now, let $z \in Z_{Max}$ stand for one of the index sets for which the sublinear combination $\sum_{l \in L^z} a_l C_g^{l, i_1 \dots i_\mu}$ in (1.7) indexes tensor fields of maximal refined double character. We assume with no loss of generality that for each $l \in L^z$ the factors $\nabla^{(A)}\Omega_1, \nabla^{(B)}\Omega_2, \nabla^{(C)}\Omega_3$ have $\alpha \geq \alpha' \geq \alpha''$ free indices respectively.⁸⁰ Therefore, the tensor fields indexed in L^z can be in one of the following two forms:

$$\begin{aligned} & p\text{contr}(\nabla^c \nabla_{i_1 \dots i_\alpha j_1 \dots j_b}^{(A)} \Omega_1 \otimes \nabla_{i_{\alpha+1} \dots i_{\alpha+\alpha'} j_{b+1} \dots j_{b+b'}}^{(B)} \Omega_2 \otimes \\ & \nabla_{c i_{\alpha+\alpha'+1} \dots i_{\alpha+\alpha'+\alpha''} j_{b+b'+1} \dots j_{b+b'+b''}}^{(2)} \Omega_3 \otimes \nabla^{j_{x_1}} \phi_1 \dots \otimes \nabla^{x_{j+j'+j''}} \phi_u), \end{aligned} \quad (3.13)$$

$$\begin{aligned} & p\text{contr}(\nabla_{i_1 \dots i_\alpha j_1 \dots j_b}^{(A)} \Omega_1 \otimes \nabla^c \nabla_{i_{\alpha+1} \dots i_{\alpha+\alpha'} j_{b+1} \dots j_{b+b'}}^{(B)} \Omega_2 \otimes \\ & \nabla_{c i_{\alpha+\alpha'+1} \dots i_{\alpha+\alpha'+\alpha''} j_{b+b'+1} \dots j_{b+b'+b''}}^{(2)} \Omega_3 \otimes \nabla^{j_{x_1}} \phi_1 \dots \otimes \nabla^{x_{j+j'+j''}} \phi_u), \end{aligned} \quad (3.14)$$

(where $A, B \geq 3$).

Now, by “manually subtracting” $Xdiv$'s from these μ -tensor fields, we can assume wlog that the tensor fields indexed in our chosen L^z are in the form (3.14).

⁸⁰Recall that by our hypothesis $\alpha' \geq 2$.

With that extra assumption, we can show that the coefficient of the tensor field (3.14) is zero. We see this by considering the (global) equation $\int_M Z_g dV_g = 0$ and using the silly divergence formula $silly[Z_g] = 0$ (which arises by integrations by parts w.r.t. to the factor $\nabla^{(A)}\Omega_1$). Picking out the sublinear combination $silly_+[Z_g]$ which consists of the complete contraction:

$$\begin{aligned} & \text{contr}(\Omega_1 \otimes \nabla^c \nabla^{f_1 \dots f_\alpha}_{j_{b+1} \dots j_{b+b'}} \Delta^{\alpha'} \Omega_2 \otimes \nabla_c \nabla_{f_1 \dots f_\alpha j_{b+b'+1} \dots j_{b+b'+b''}} \Delta^{\alpha''} \Omega_3 \otimes \\ & \Delta \phi_1 \dots \nabla^{j_{b+b'+b''}} \phi_u) \end{aligned} \quad (3.15)$$

(notice that $silly_+[Z_g] = 0$), we derive that the coefficient of (3.14) must vanish. Thus, we have shown our claim in this second subcase also. \square

The case $p = 2, \sigma_2 = 1$: Recall that in this case we fall under the special case when $n = 2u + 2\mu$. In this setting, we will have that in each index set $L^z, z \in Z'_{Max}$ (see the statement of Lemma 3.5 in [6]) there is a *unique* μ -tensor field of maximal refined double character in (1.7), where the two indices k, l in the factor $S_* \nabla^{(\nu)} R_{ijkl}$ will be contracting against one of the factors $\nabla^{(A)}\Omega_1, \nabla^{(B)}\Omega_2$ (wlog we may assume that they are contracting against *different factors*). But now, recall that since we are considering case *A* of Lemma 3.5 in [6], one of the factors $\nabla^{(A)}\Omega_1, \nabla^{(B)}\Omega_2$ will have at least two free indices. Hence, in at least one of the factors $\nabla^{(A)}\Omega_1, \nabla^{(B)}\Omega_2$, the index k, l is removable (meaning that it can be erased, and we will be left with an acceptable tensor field). We denote by $C_g^{l, i_1 \dots i_\mu i_{\mu+1}}$ the tensor field that arises from $C_g^{l, i_1 \dots i_\mu}$ by erasing the aforementioned k, l and making k or l into a free index, we then observe that:

$$C_g^{l, i_1 \dots i_\mu} - X \text{div}_{i_{\mu+1}} C_g^{l, i_1 \dots i_\mu i_{\mu+1}} = 0 \quad (3.16)$$

(modulo complete contractions of length $\geq \sigma + u + 1$). This completes the proof of our claim. \square

References

- [1] S. Alexakis *On the decomposition of Global Conformal Invariants I*, Ann. of Math. **170** (2009), no. 3, 1241–1306.
- [2] S. Alexakis *On the decomposition of Global Conformal Invariants II*, Adv. in Math. **206** (2006), 466–502.
- [3] S. Alexakis *The decomposition of Global Conformal Invariants: A Conjecture of Deser an Schwimmer I*, arXiv.
- [4] S. Alexakis *The decomposition of Global Conformal Invariants: A Conjecture of Deser an Schwimmer II*, arXiv.
- [5] S. Alexakis *The decomposition of Global Conformal Invariants: A Conjecture of Deser an Schwimmer III*, arXiv.

- [6] S. Alexakis *The decomposition of Global Conformal Invariants: A Conjecture of Deser an Schwimmer IV*, arXiv.
- [7] S. Alexakis *The decomposition of Global Conformal Invariants: A Conjecture of Deser an Schwimmer V*, arXiv.
- [8] S. Alexakis *The decomposition of Global Conformal Invariants: A Conjecture of Deser an Schwimmer VI*, arXiv.
- [9] M. Atiyah, R. Bott, V. K. Patodi, *On the heat equation and the index theorem* Invent. Math. **19** (1973), 279–330.
- [10] T. N. Bailey, M. G. Eastwood, A. R. Gover *Thomas’s structure bundle for conformal, projective and related structures* Rocky Mountain J. Math. **24** (1994), no. 4, 1191–1217.
- [11] T. N. Bailey, M. G. Eastwood, C. R. Graham *Invariant Theory for Conformal and CR Geometry* Ann. of Math. (2), **139** (1994), 491-552.
- [12] N. Berline, E. Getzler, M. Vergne *Heat kernels and Dirac operators* Grundlehren Text Editions. Springer-Verlag, Berlin, 2004.
- [13] N. Boulager, J. Erdmenger *A Classification of Local Weyl Invariants in $D=8$* , Class. Quantum Gravity **21** (2004), 4305-4316.
- [14] T. Branson, P. Gilkey, J. Pohjanpelto *Invariants of locally conformally flat manifolds* Trans. Amer. Math. Soc. **347** (1995), no. 3, 939–953.
- [15] A. Ćap, A.R. Gover *Tractor calculi for parabolic geometries* Trans. Amer. Math. Soc. **354** (2002), no. 4, 1511–1548.
- [16] A. Ćap, A.R. Gover *Standard tractors and the conformal ambient metric construction* Ann. Global Anal. Geom. **24** (2003), no. 3, 231–259.
- [17] É. Cartan *Sur la réduction à sa forme canonique de la structure d’un groupe de transformations fini et continu*, Oeuvres Complètes **1**, Part 1, 293-355, Gauthier-Villars, Paris, 1952.
- [18] S. Deser, A. Schwimmer *Geometric classification of conformal anomalies in arbitrary dimensions*, Phys. Lett. B309 (1993) 279-284.
- [19] L.P. Eisenhart *Riemannian Geometry* Princeton University Press (1925)
- [20] C. Fefferman *Monge-Ampère equations, the Bergman kernel and geometry of pseudo-convex domains*, Ann. of Math. **103** (1976), 395-416; Erratum **104** (1976), 393-394.
- [21] C. Fefferman, C. R. Graham *Conformal Invariants* Élie Cartan et les mathématiques d’aujourd’hui, Astérisque numero hors serie, 1985, 95-116.
- [22] C. Fefferman, C. R. Graham *The ambient metric*, arXiv:0710.0919.

- [23] P. Gilkey *Local invariants of an embedded Riemannian manifold* Ann. of Math. (2) **102** (1975), no. 2, 187–203.
- [24] C. R. Graham, K. Hirachi *Inhomogeneous ambient metrics*, Symmetries and overdetermined systems of partial differential equations, 403–420, IMA Vol. Math. Appl., **144**, Springer, New York, 2008.
- [25] K. Hirachi *Construction of Boundary Invariants and the Logarithmic Singularity of the Bergman Kernel* Ann. of Math. (2) **151** (2000), no. 2 151–191.
- [26] K. Hirachi *Logarithmic singularity of the Szegő kernel and a global invariant of strictly pseudoconvex domains* Ann. of Math. (2) **163** (2006), no. 2, 499–515.
- [27] T. Y. Thomas *The differential invariants of generalized spaces*, Cambridge University Press, Cambridge 1934.
- [28] H. Weyl *The classical groups*, Princeton University press.
- [29] S. Zelditch *Szegő kernels and a theorem of Tian*, Internat. Math. Res. Notices 1998, no. 6, 317–331.