

PROPERTIES OF SERIES

March 6th, 2019

For next week

For Monday (Mar 11), watch the videos:

- Integral and comparison tests: 13.10, 13.11, 13.12

For Wednesday (Mar 13), watch the videos:

- Alternating series: 13.13, 13.14
- Absolute convergence: 13.15, (13.16), (13.17)

Comment: I wrote some notes about slide 7 from last lecture.

<http://www.math.toronto.edu/campesato/ens/1819/lec39-notes.pdf>

What is wrong with this calculation? Fix it

Claim:

$$\sum_{n=2}^{\infty} \ln \frac{n}{n+1} = \ln 2$$

“Justification”

$$\begin{aligned} \sum_{n=2}^{\infty} \ln \frac{n}{n+1} &= \sum_{n=2}^{\infty} [\ln n - \ln(n+1)] \\ &= \sum_{n=2}^{\infty} \ln(n) - \sum_{n=2}^{\infty} \ln(n+1) \\ &= (\ln 2 + \ln 3 + \ln 4 + \dots) - (\ln 3 + \ln 4 + \dots) \\ &= \ln 2 \end{aligned}$$

Geometric sums

Prove the following claim.

Let $x \in \mathbb{R} \setminus \{1\}$ then

$$\sum_{n=p}^q x^n = \frac{1 - x^{q-p+1}}{1 - x} x^p$$

$$\sum_{n=p}^q x^n = \frac{1 - x^{q-p+1}}{1 - x} x^p$$

First term of the sum

Number of elements in the sum

Compute the value of the following series:

$$1 \quad S_1 = 1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \frac{1}{81} + \dots$$

$$2 \quad S_2 = \frac{1}{2} - \frac{1}{4} + \frac{1}{8} - \frac{1}{16} + \frac{1}{32} - \dots$$

$$3 \quad S_3 = \frac{3}{2} - \frac{9}{4} + \frac{27}{8} - \frac{81}{16} + \dots$$

$$4 \quad S_4 = 1 + \frac{1}{2^{0.5}} + \frac{1}{2} + \frac{1}{2^{1.5}} + \frac{1}{2^2} + \frac{1}{2^{2.5}} + \dots$$

$$5 \quad S_5 = \sum_{n=1}^{\infty} (-1)^n \frac{3^n}{2^{2n+1}} \quad 6 \quad S_6 = \sum_{n=r}^{\infty} x^n$$

$$1 \quad \sum_{n=0}^{\infty} \frac{1}{2^n}$$

$$2 \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{2^n}$$

$$3 \quad \sum_{n=1}^{\infty} \frac{1}{2^{n/2}}$$

$$4 \quad \sum_{n=5}^{\infty} \frac{3^n}{2^{2n+1}}$$

$$5 \quad \sum_{n=3}^{\infty} \frac{3^n}{1000 \cdot 2^{n+2}}$$

$$6 \quad \sum_{n=0}^{\infty} (-1)^n$$

True or False – Series

Let $\sum_{n=0}^{\infty} a_n$ be a series. Let $\{S_n\}_{n=0}^{\infty}$ be its partial-sum sequence.

1 IF the series $\sum_{n=0}^{\infty} a_n$ is divergent, THEN $\exists n \in \mathbb{N}$ such that $a_n > 100$

2 IF the series $\sum_{n=0}^{\infty} a_n$ is divergent, THEN $\exists n \in \mathbb{N}$ such that $S_n > 100$

3 IF the series $\sum_{n=0}^{\infty} a_n$ converges

THEN the series $\sum_{n=100}^{\infty} a_n$ converges to a smaller number.

4 IF the series $\sum_{n=0}^{\infty} a_n$ converges

THEN the sequence $\{S_n\}_{n=0}^{\infty}$ is eventually monotonic.

True or False – Series

Let $\sum_{n=0}^{\infty} a_n$ be a series. Let $\{S_n\}_{n=0}^{\infty}$ be its partial-sum sequence.

5 IF the sequence $\{S_n\}_{n=0}^{\infty}$ is bounded and eventually monotonic, THEN the series $\sum_{n=0}^{\infty} a_n$ is convergent.

6 IF the sequence $\{S_n\}_{n=0}^{\infty}$ is increasing, THEN $\forall n > 0, a_n > 0$.

7 IF the sequence $\{S_n\}_{n=0}^{\infty}$ is increasing, THEN $\forall n \geq 0, a_n > 0$.

- 8 IF $\lim_{n \rightarrow \infty} a_n = 0$, THEN $\sum_n a_n$ is convergent.
- 9 IF $\lim_{n \rightarrow \infty} a_n \neq 0$, THEN $\sum_n a_n$ is divergent.
- 10 IF $\sum_n a_n$ is convergent THEN $\lim_{n \rightarrow \infty} a_n = 0$.
- 11 IF $\sum_n a_n$ is divergent THEN $\lim_{n \rightarrow \infty} a_n \neq 0$.

Is $0.999999\dots = 1$?

First, we need to fix the meaning of $0.999999\dots$

- 1 Write the number $0.999999\dots$ as a series
Hint: $427 = 400 + 20 + 7$.
- 2 Compute the first few partial sums
- 3 Add up the series.
Hint: it is geometric.

Let $\sum_{n=0}^{\infty} a_n$ be a series and $c \in \mathbb{R}$.

Prove that

IF $\sum_{n=0}^{\infty} a_n$ is convergent,

THEN $\sum_{n=0}^{\infty} (ca_n)$ is also convergent and $\sum_{n=0}^{\infty} (ca_n) = c \left(\sum_{n=0}^{\infty} a_n \right)$.

Are all decimal expansions well-defined?

The decimal expansion of a number is a series:

$$0.a_1a_2a_3a_4a_5\dots = \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{1000} + \dots$$

for any “digits” a_1, a_2, a_3, \dots

Are such series always convergent, no matter which infinite string of digits we choose?

Yes! Prove it.

(Hint: all the terms in the series are positive.)

Challenge

We want to calculate the value of

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)3^n}$$

Hints:

- 1 Compute $\sum_{n=0}^{\infty} (-1)^n x^{2n}$
- 2 Compute $\frac{d}{dx} [\arctan x]$
- 3 Pretend you can take derivatives and antiderivatives of series the way you can take them of sums. Which series adds up to $\arctan x$?
- 4 Now attempt the original problem.

A more difficult challenge - part 1

Homework, not mandatory

Let $x \in (0, 1)$. Then, there exists a unique sequence $(a_n)_{n \geq 1}$ of digits not eventually constant to 9 such that $x = 0.a_1 a_2 a_3 \dots$. We want to prove that x is rational if and only if the sequence $(a_n)_n$ is eventually periodic, i.e.

$$\exists n_0 \geq 1, \exists p \geq 1, \forall n \geq 1, (n \geq n_0 \implies a_{n+p} = a_n)$$

- 1 Prove that if the sequence $(a_n)_n$ is eventually periodic then

$$x = r + \frac{y}{10^{n_0}} \sum_{l=0}^{+\infty} 10^{-lp}$$

where $y = a_{n_0+1} + \frac{a_{n_0+1}}{10} + \dots + \frac{a_{n_0+p-1}}{10^{p-1}}$ and $r \in \mathbb{R}$.

Conclude that $x \in \mathbb{Q}$.

A more difficult challenge - part 2

Homework, not mandatory

- 2 Conversely, we now assume that $x = \frac{a}{b}$.
 - 1 Prove there exist $0 < s < t$ such that the divisions of $10^s a$ and $10^t a$ by b have the same remainder.
 - 2 We denote by $(b_n)_{n \geq 1}$ and $(c_n)_{n \geq 1}$ the digits of the fractional parts of $\frac{10^s a}{b}$ and $\frac{10^t a}{b}$. Express b_n and c_n in terms of a_n .
 - 3 Prove that $\frac{10^s a}{b} - \frac{10^t a}{b} \in \mathbb{Z}$.
 - 4 Using the above question, find a relation between b_n and c_n .
 - 5 Conclude that $(a_n)_n$ is eventually periodic.