

## RATIO TEST

March 18<sup>th</sup>, 2019

## For next lecture

For Wednesday (Mar 20), watch the videos:

- Power series: 14.1, 14.2
- Taylor polynomials: 14.3, 14.4

## Ratio test: Convergent or divergent?

Use Ratio test to decide which series are convergent:

$$1 \quad \sum_{n=1}^{\infty} \frac{3^n}{n!}$$

$$3 \quad \sum_{n=1}^{\infty} \frac{n!(2n)!}{(3n)!}$$

$$2 \quad \sum_{n=1}^{\infty} \frac{(2n)!}{n!^2 3^{n+1}}$$

$$4 \quad \sum_{n=2}^{\infty} \frac{n!}{n^n}$$

## Ratio test

Try to apply the ratio test to :

$$1 \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$$

$$2 \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$$

$$3 \quad \sum_{n=1}^{\infty} \frac{1}{n}$$

## Review: Convergent or Divergent?

$$1 \quad \sum_{n=2}^{+\infty} \frac{\cos(n\pi)}{\ln n}$$

$$2 \quad \sum_{n=2}^{+\infty} \frac{\sin(n\pi)}{\ln n}$$

Conclusion: Think first, then compute!

## Convergent alternating series failing the AST *Homework*

$$\text{Consider } S = \sum_{n=2}^{+\infty} \frac{(-1)^n}{n + (-1)^n}.$$

1 Why does  $S$  fail the AST?

*Hint: compare  $b_{2n}$  and  $b_{2n+1}$ .*

2 Prove that  $S$  is nevertheless convergent!

*Hint:  $\frac{(-1)^n}{n+(-1)^n} = \frac{(-1)^n}{n} + v_n$*

## Review: Bertrand series

For which  $a, b \in \mathbb{R}$  is the following series convergent:

$$\sum_{n=2}^{+\infty} \frac{1}{n^a \ln(n)^b}$$

## AST – Homework


Construct a series of the form  $\sum_{n=1}^{\infty} (-1)^n b_n$  s.t.

•  $\forall n \geq 1, b_n > 0$

•  $\lim_{n \rightarrow \infty} b_n = 0$

• the series  $\sum_{n=1}^{\infty} (-1)^n b_n$  is divergent.

## Example of decision tree to study the convergence of $\sum a_n$



## The two reference series

Quite often we apply the comparison tests with these two series that you MUST know!

### Riemann series

$$\sum_{n=k}^{+\infty} \frac{1}{n^p} \text{ is convergent if and only if } p > 1.$$

Proof: use the  $\sum - \int$  comparison.

### Geometric series

$$\sum_{n=k}^{+\infty} x^n \text{ is convergent if and only if } |x| < 1,$$

$$\text{then } \sum_{n=k}^{+\infty} x^n = \frac{x^k}{1-x}.$$

Proof: use the geometric sum formula from last week.

## Abel transformation (or Summation by parts)

Let  $(a_n)_{n \geq p}$  and  $(b_n)_{n \geq p}$  be two sequences.

We set, for  $n \geq p$ ,  $B_n = \sum_{k=p}^n b_k$ .

- Express  $b_n$  in terms of  $B_n$ .  
(you have to distinguish  $n = p$  and  $n > p$ )

- Prove that  $\sum_{n=p}^q a_n b_n = a_p B_p + \sum_{n=p+1}^q a_n (B_n - B_{n-1})$ .

- Prove that  $\sum_{n=p}^q a_n b_n = a_q B_q + \sum_{n=p}^{q-1} B_n (a_n - a_{n+1})$ .

This last equality is called “Abel transformation” or “Summation by parts”.

## Dirichlet's test - Homework

- Use an Abel transformation to prove the Dirichlet's test:  
Let  $(a_n)_{n \geq p}$  and  $(b_n)_{n \geq p}$  be two sequences.

IF

- $(a_n)_n$  is monotonic,
- $\lim_{n \rightarrow +\infty} a_n = 0$ , and,
- the sequence of partial sums  $\left( \sum_{k=p}^n b_k \right)_n$  is bounded

THEN  $\sum_{n=p}^{+\infty} a_n b_n$  is convergent.

- Hint:  $\lim_{q \rightarrow +\infty} a_q B_q = 0$  and the series  $\sum_{n=p}^{+\infty} B_n (a_n - a_{n+1})$  is absolutely convergent (hint: it is almost telescopic).
- Application: use this test to obtain a new proof of the AST.

We want to prove that  $\sum_{n=1}^{+\infty} \frac{\sin n}{n}$  is convergent.

1 Prove that

$$2 \sin(n) \sin\left(\frac{1}{2}\right) = \cos\left(n - \frac{1}{2}\right) - \cos\left(n + \frac{1}{2}\right)$$

2 Use the above equality to compute

$$\sum_{n=1}^k \sin(n)$$

3 Conclude using the Dirichlet's test.