

MOTIVIC CHERN CLASSES AND IWAHORI INVARIANTS OF PRINCIPAL SERIES

CHANGJIAN SU

ABSTRACT. In this expository note, we review the proof [AMSS19] of conjectures of Bump, Nakasuji and Naruse about principal series representations of p -adic groups. The ingredients of the proof involve Maulik–Okounkov K-theoretic stable basis for the Springer resolution, and motivic Chern classes of Schubert cells for the Langlands dual group

CONTENTS

1. Introduction	1
Notations	3
2. p -adic side	3
2.1. Iwahori invariants of the principal series representations	3
2.2. Two bases in the Iwahori invariants of the principal series representations	4
2.3. Conjectures of Bump, Nakasuji and Naruse	5
3. Complex dual side	7
3.1. Definition of Motivic Chern classes	7
3.2. Two bases in the localized equivariant K theory of the flag variety	8
3.3. Hecke algebra action	8
3.4. Smoothness criterion	10
4. The proof	10
References	11

1. INTRODUCTION

The Iwahori–Hecke algebra of a p -adic reductive group is the convolution algebra on the compactly supported functions on the rational points of the group, which are left and right invariant under the Iwahori subgroup. It contains the finite Hecke algebra and the group algebra of the character lattice of the complex dual maximal torus, which is called the lattice part of the Iwahori–Hecke algebra. The Iwahori invariant subspace of any unramified principal series representation is a module over the Iwahori–Hecke algebra via convolution. In the invariant subspace, there is a natural basis induced from the cell decomposition of the group. I.e., the elements in this basis are just characteristic functions of the cells. If the unramified character is regular, then the invariant subspace is a regular representation of the finite Hecke algebra, and this basis is the standard basis. However, this basis does not behave well under the intertwiners, making it difficult to compute the corresponding Iwahori–Whittaker functions.

In Casselman’s study of unramified principal series [Cas80], he introduced another basis, which is called the Casselman basis now, in the Iwahori invariant subspace. The Casselman

Date: September 10, 2019.

basis enjoys many good properties. For example, this is a eigenbasis for the lattice part in the Iwahori–Hecke algebra, and it behaves nicely with the intertwiners. Using this new basis, Casselman can reprove the Macdonald’s formula for the spherical function [Mac68], and he also established the formula for the spherical Whittaker function with Shalika [CS80], which is called the Casselman–Shalika formula. Moreover, there are simple formulas for the Whittaker function for the Casselman basis elements [Ree93].

Therefore, to compute the Iwahori–Whittaker functions, we only need to know the transition matrix between these two bases. Special entries of this matrix can be deduced from the Gindikin–Karpelevich formula proved by Langlands [Lan71] in the p -adic setting. Motivated by this formula, Bump and Nakasuji [BN11, BN19] make two conjectures about certain matrix coefficients assuming the Dynkin diagram of the group is simply laced. One of them is about the factorization property, while the other one is about the analyticity property of the coefficients. On the other hand, Nakasuji and Naruse [NN16] use Yang–Baxter basis to give a combinatorial formula for every entry of this matrix. However, the formula is too complicated so that we do not know how to use it to prove the conjectures of Bump and Nakasuji.

The difficulty in the factorization conjecture lies in the fact that the condition in the conjecture is about the triviality of certain Kazhdan–Lusztig polynomials [KL79], which is quite hard to work with combinatorially. However, if the Dynkin diagram is simply laced, it is well known that the triviality of a Kazhdan–Lusztig polynomial is equivalent to the condition that the Schubert variety in the dual flag manifold is smooth at certain torus fixed point [BL00]. Based on this observation, Naruse refined the factorization conjecture, so that the refined conjecture can be thought of as a smoothness criterion for the Schubert varieties using p -adic representations. In Naruse’s refinement, the Dynkin diagram of the group is no longer assumed to be simply laced, and it is an if and only if statement. One of the directions was claimed by Naruse [Nar14].

In this refined factorization conjecture, one side is about p -adic representations, while the other side is about Schubert varieties for the complex dual groups. The connection between these two sides are well studied in geometric representation theory. For example, Kazhdan and Lusztig used equivariant K theory to prove the Deligne–Langlands conjecture [KL87]. Recall the Iwahori–Hecke algebra is defined as certain functions on the p -adic group. Kazhdan and Lusztig gave another geometric realization of the Iwahori–Hecke algebra via the equivariant K theory of the Steinberg variety, which is a convolution algebra, see also [CG09]. Moreover, these are categorified by R. Bezrukavnikov in [Bez16].

The proof in [AMSS19] can be thought of a shadow of these two geometric realizations of the Iwahori–Hecke algebra. To be more specific, the strategy is to consider two geometric realizations of the regular representation of the finite Hecke algebra. On one hand, we have the Iwahori invariants of a regular unramified principal series representation. On the other hand, we have the (specialized) equivariant K theory of the flag variety for the complex dual group, on which the lattice part subalgebra in the Iwahori–Hecke algebra acts by tensoring by line bundles on the flag variety. By the localization theorem in equivariant K theory, there is a natural basis in the localized equivariant K theory of the flag variety—the structure sheaf of the fixed points, and it is immediate to see that they form a eigenbasis for the lattice part of the Iwahori–Hecke algebra. Thus, they correspond to the Casselman basis.

The other basis in the equivariant K theory corresponding to the standard basis in the p -adic side is called the (dual) motivic Chern classes of the Schubert cells. To generalize the Chern–Schwartz–Macpherson classes for singular varieties in homology [Mac74, Sch65a, Sch65b], Brasselet, Schürmann and Yokura [BSY10] introduce motivic Chern classes in K

theory. See [AMSS19, FRW18] for the equivariant case. The motivic Chern classes of the Schubert cells form another basis in the equivariant K theory of the flag variety. Under the above two geometric realizations of the regular representation of the finite Hecke algebra, the standard basis corresponds to the (dual) motivic Chern classes.

Via these correspondences, we can use the equivariant K theory of the flag variety to give a geometric interpretation of the transition matrix coefficients on the p -adic side. I.e., the coefficients are related to the localizations of the motivic Chern classes of the Schubert cells¹. Finally, the factorization conjecture follows since the motivic Chern classes of Schubert cells contain information about the singularity of the Schubert varieties by its very definition. To be more specific, Kumar [Kum96] has a smoothness criterion for the Schubert varieties using the localization of the Schubert classes in the equivariant cohomology of the flag variety. We generalize it using the localization of the motivic Chern classes.

The analyticity conjecture is proved by proving an analogue of it on the flag variety side, which follows from a GKM type argument.

This note is structured as follows. In Section 2, we introduce the objects in the p -adic side: the Iwahori invariant subspace, the two bases, and conjectures of Bump, Nakasuji and Naruse. In Section 3, we introduce the motivic Chern classes in the complex dual side. Lastly, we sketch the proof of the conjectures in Section 4.

Acknowledgments. The author thanks the organizers of the the 8th International Congress of Chinese Mathematicians for the invitation to speak. He also thanks his advisor Prof. Andrei Okounkov for guidance, and Paolo Aluffi, Leonardo C. Mihalea, Jörg Schürmann, Gufang Zhao and Changlong Zhong for collaborations.

Notations. Let F be a non-Archimedean local field, with ring of integers \mathcal{O}_F , uniformizer $\varpi \in \mathcal{O}_F$, and residue field \mathbb{F}_q . Examples are finite extensions of the field of p -adic numbers \mathbb{Q}_p , or of the field of Laurent series over \mathbb{F}_p . Let G be a split, adjoint, reductive group defined over \mathcal{O}_F . Let $B = TN$ be a Borel subgroup containing a maximal torus T and its unipotent radical N . Let R^+ denote the roots in B . We will use α, β (resp. α^\vee, β^\vee) to denote roots (resp. coroots), and use $\alpha > 0$ to denote $\alpha \in R^+$. Let W be the Weyl group, and \leq denote the Burhat order on it. Let w_0 be the longest element in the Weyl group W .

Let $T^\vee \subset B^\vee \subset G^\vee$ be the complex Langlands dual groups, and $B^{\vee,-}$ be the opposite Borel subgroup. Since G is adjoint, G^\vee is simply connected. Let \mathcal{B} denote the flag variety G^\vee/B^\vee . For any $w \in W$, let $X(w)^\circ := B^\vee w B^\vee / B^\vee \subset G^\vee/B^\vee$ and $Y(w)^\circ := B^{\vee,-} w B^\vee / B^\vee \subset G^\vee/B^\vee$ be the (opposite) Schubert cells, with closures denoted by $X(w)$ and $Y(w)$, respectively. Let $X^*(T^\vee) = X_*(T)$ be the group of characters of T^\vee (=the group of cocharacters of T), and $X_*(T^\vee) = X^*(T)$ be the group of cocharacters of T^\vee (= the group of characters of T). For any $\lambda \in X^*(T^\vee)$, let $\mathcal{L}_\lambda := G^\vee \times_{B^\vee} \mathbb{C}_\lambda$ denote the line bundle over the flag variety \mathcal{B} .

2. p -ADIC SIDE

In this section, we introduce two bases in the Iwahori invariants of the principal series representations of a split reductive p -adic group G , and state the conjectures of Bump, Nakasuji and Naruse about the transition matrix coefficients between these two bases.

2.1. Iwahori invariants of the principal series representations. In this section, we introduce the unramified principal series representations.

¹The full proof of this fact also involves Maulik and Okounkov's stable basis [Oko17, SZZ17, SZ19], which is quite lengthy. The interested reader can refer to the paper [AMSS19].

Let $G(F)$ be the F -points of G , and similarly for the maximal torus $T(F)$ and Borel subgroup $B(F) = T(F)N(F)$. Let I be an Iwahori subgroup, i.e., the inverse image of $B(\mathbb{F}_q)$ under the evaluation map $G(\mathcal{O}_F) \rightarrow G(\mathbb{F}_q)$.

Let $\mathbb{H} = \mathbb{C}_c[I \backslash G(F) / I]$ be the Iwahori–Hecke algebra, consisting of compactly supported functions on $G(F)$ which are bi-invariant under I . As a vector space, $\mathbb{H} = \Theta \otimes_{\mathbb{C}} H_W(q)$, where Θ is a commutative subalgebra isomorphic to the coordinate ring $\mathbb{C}[T^\vee]$ of the complex dual torus $T^\vee := \mathbb{C}^* \otimes X^*(T)$, and where $H_W(q)$ is the finite Hecke (sub)algebra with parameter q associated to the finite Weyl group W . The finite Hecke algebra $H_W(q)$ is also a subalgebra of \mathbb{H} , and it is generated by elements T_w ($w \in W$) such that the following relations hold: $T_u T_v = T_{uv}$ if $\ell(uv) = \ell(u) + \ell(v)$, and $(T_{s_i} + 1)(T_{s_i} - q) = 0$ for a simple reflection s_i in W .

For any character τ of T , and α^\vee a coroot define e^{α^\vee} by $e^{\alpha^\vee}(\tau) = \tau(h_{\alpha^\vee}(\varpi))$, where $h_{\alpha^\vee} : F^\times \rightarrow T(F)$ is the one parameter subgroup. There is a pairing

$$\langle \cdot, \cdot \rangle : T(F) / T(\mathcal{O}_F) \times T^\vee \rightarrow \mathbb{C}^*$$

given by $\langle a, z \otimes \lambda \rangle = z^{\text{val}(\lambda(a))}$. This induces an isomorphism between $T(F) / T(\mathcal{O}_F)$ and the group $X^*(T^\vee)$ of rational characters of T^\vee . It also induces an identification between T^\vee and unramified characters of $T(F)$, i.e., characters which are trivial on $T(\mathcal{O})$.

From now on we take τ to be an unramified character of $T(F)$ such that $e^\alpha(\tau) \neq 1$ for all coroots α , and for which the stabilizer $W_\tau = 1$. The *principal series representation* is the induced representation $I(\tau) := \text{Ind}_{B(F)}^{G(F)}(\tau)$. As a \mathbb{C} -vector space, $I(\tau)$ consists of locally constant functions f on $G(F)$ such that $f(bg) = \tau(b)\delta^{\frac{1}{2}}(b)f(g)$ for any $b \in B(F)$, where $\delta(b) := \prod_{\alpha > 0} |\alpha^\vee(b)|_F$ is the modulus function on the Borel subgroup. The group $G(F)$ acts by the formula $(\pi(g)f)(h) = f(hg)$, where $g, h \in G(F)$ and $f \in I(\tau)$. The Iwahori–Hecke algebra \mathbb{H} acts through convolution from the right on the Iwahori invariant subspace $I(\tau)^I$, so that the restriction of this action to $H_W(q)$ is a regular representation. One can pass back and forth between left and right \mathbb{H} -modules by using the standard anti-involution ι on \mathbb{H} given by $\iota(h)(x) = h(x^{-1})$, where $h \in \mathbb{H}$ and $x \in G(F)$. For any $w \in W$, $\iota(T_w) = T_{w^{-1}}$ and $\iota(q) = q$, see [HKP10, Section 3.2]. We use π to denote this left action of \mathbb{H} on $I(\tau)^I$.

2.2. Two bases in the Iwahori invariants of the principal series representations.

In this section, we recall the definitions and properties of the two bases in $I(\tau)^I$.

From the decomposition $G(F) = \bigsqcup_{w \in W} B(F)wI$, one obtains the basis of the *characteristic functions* on the orbits, denoted by $\{\varphi_w \mid w \in W\}$ ². For $w \in W$, the element φ_w is characterized by the two conditions: [Ree92, pg. 319]:

- (1) φ_w is supported on $B(F)wI$;
- (2) $\varphi_w(bwg) = \tau(b)\delta^{\frac{1}{2}}(b)$ for any $b \in B(F)$ and $g \in I$.

The left action of \mathbb{H} on $I(\tau)^I$ was calculated by Casselman in [Cas80, Thm. 3.4] as follows. For any simple coroot α_i^\vee :

$$(1) \quad \pi(T_{s_i})(\varphi_w) = \begin{cases} q\varphi_{ws_i} + (q-1)\varphi_w & \text{if } ws_i < w; \\ \varphi_{ws_i} & \text{if } ws_i > w. \end{cases}$$

The second basis, called *Casselman's basis*, and denoted by $\{f_w \mid w \in W\}$, was defined by Casselman [Cas80, §3] by duality using certain intertwiner operators. For any character τ and $x \in W$, define $x\tau \in X^*(T)$ by the formula $x\tau(a) := \tau(x^{-1}ax)$ for any $a \in T$. Since τ that is unramified and it has trivial stabilizer under the Weyl group action, the space

²Our φ_w is equal to $\phi_{w^{-1}}$ in [BN11, BN19]

$\mathrm{Hom}_{G(F)}(I(\tau), I(x^{-1}\tau))$ is known to be one dimensional, spanned by an operator $\mathcal{A}_x = \mathcal{A}_x^\tau$ ³ defined by

$$\mathcal{A}_x(\varphi)(g) := \int_{N_x} \varphi(\dot{x}ng)dn,$$

where \dot{x} is a representative of $x \in W$, $N_x = N(F) \cap \dot{x}^{-1}N^-(F)\dot{x}$ where N^- is the unipotent radical of the opposite Borel subgroup B^- ; the measure on N_x is normalized by the condition that $\mathrm{Vol}(N_x \cap G(\mathcal{O}_F)) = 1$ [Ree92]. If $x, y \in W$ satisfy $\ell(x) + \ell(y) = \ell(xy)$, then $\mathcal{A}_y^{x^{-1}\tau} \mathcal{A}_x^\tau = \mathcal{A}_{xy}^\tau$. Then there exist unique functions $f_w \in I(\tau)^I$ such that

$$(2) \quad \mathcal{A}_x^\tau(f_w)(1) = \delta_{x,w}.$$

(Again under our conventions f_w equals the element denoted $f_{w^{-1}}$ in [BN11].) For the longest element w_0 in the Weyl group, Casselman showed in [Cas80, Prop. 3.7] that

$$\varphi_{w_0} = f_{w_0}.$$

Reeder [Ree92] calculated the action of \mathbb{H} on the functions f_w : he showed in [Ree92, Lemma 4.1] that the functions f_w are Θ -eigenvectors, and he calculated in [Ree92, Prop. 4.9] the action of $H_W(q)$. To describe the latter, for any coroot α^\vee , let

$$c_{\alpha^\vee} = \frac{1 - q^{-1}e^{\alpha^\vee}(\tau)}{1 - e^{\alpha^\vee}(\tau)}.$$

For any simple coroot α_i^\vee and $w \in W$, write

$$J_{i,w} = \begin{cases} c_{w(\alpha_i^\vee)} c_{-w(\alpha_i^\vee)} & \text{if } ws_i > w; \\ 1 & \text{if } ws_i < w. \end{cases}$$

Then, we have

$$(3) \quad \pi(T_{s_i})(f_w) = q(1 - c_{w(\alpha_i^\vee)})f_w + qJ_{i,w}f_{ws_i}.$$

2.3. Conjectures of Bump, Nakasuji and Naruse. In this section we state conjectures of Bump, Nakasuji and Naruse, regarding factorization and analyticity properties of the transition matrix coefficients between the bases $\{\varphi_w\}$ and $\{f_w\}$.

We follow mainly [BN11], and we recall that we use opposite notations from those in *loc.cit*: our φ_w and f_w are $\phi_{w^{-1}}$ and $f_{w^{-1}}$ respectively in [BN11]. Let

$$\phi_u := \sum_{u \leq w} \varphi_w \in I(\tau)^I,$$

and consider the expansion in terms of the Casselman's basis:

$$\phi_u = \sum_w m_{u,w} f_w.$$

Then by the definition of f_w , $m_{u,w} = \mathcal{A}_w(\phi_u)(1)$. It is also easy to see that $m_{u,w} = 0$ unless $u \leq w$, see [BN11, Theorem 3.5].

When $u = id$, ϕ_{id} is the spherical vector in $I(\tau)$, i.e. the vector fixed by the maximal compact subgroup $G(\mathcal{O}_F)$, and

$$(4) \quad \mathcal{A}_w(\phi_{id})(1) = m_{id,w} = \prod_{\alpha > 0, s_\alpha w < w} \frac{1 - q^{-1}e^{\alpha^\vee}(\tau)}{1 - e^{\alpha^\vee}(\tau)}.$$

This is the *Gindikin–Karpelevich formula*, which in the non-Archimedean setting was actually proved by Langlands [Lan71] after Gindikin and Karpelevich proved a similar statement

³The intertwiner \mathcal{A}_x is related to M_x from [Cas80, BN11] by the formula $\mathcal{A}_x = M_{x^{-1}}$.

for real groups. Casselman gave another proof using his basis f_w , and this plays an crucial role in his computation of Macdonald formula and the spherical Whittaker functions, see [Cas80, CS80]. See also [SZZ17] for an approach using the stable basis and the equivariant K-theory of the cotangent bundle $T^*(G^\vee/B^\vee)$.

2.3.1. *The factorization conjecture.* For any $u \leq w \in W$, let

$$S(u, w) := \{\beta \in R^+ | u \leq s_\beta w < w\}.$$

Based on Equation (4), Bump and Nakasuji conjectured that [BN11, BN19]

Theorem 2.1 (Bump–Nakasuji conjecture). *Assume the Dynkin diagram of G is simply laced. For any $u \leq w \in W$, then*

$$m_{u,w} = \prod_{\alpha \in S(u,w)} \frac{1 - q^{-1}e^{\alpha^\vee}(\tau)}{1 - e^{\alpha^\vee}(\tau)},$$

if the Kazhdan–Lusztig polynomial

$$P_{w_0w^{-1}, w_0u^{-1}}(q) = 1.$$

For any finite simple root system (not necessarily simply laced), it is easy to see that $P_{w_0w^{-1}, w_0u^{-1}}(q) = 1$ if and only if $P_{w_0w, w_0u}(q) = 1$ (see [AMSS19, Corollary 8.6]). If the Dynkin diagram of G is simply laced, then $P_{w_0w, w_0u}(q) = 1$ is equivalent to the condition that the opposite Schubert variety $Y(u) := \overline{B^{\vee, -u}B^\vee/B^\vee}$ in the dual complex flag manifold G^\vee/B^\vee is smooth at the torus fixed point $wB^\vee \in G^\vee/B^\vee$ [BL00]. Based on this observation, Naruse [Nar14] refined the Bump–Nakasuji conjecture as follows.

Theorem 2.2 (Bump–Nakasuji–Naruse factorization Conjecture). *For any $u \leq w \in W$,*

$$m_{u,w} = \prod_{\alpha \in S(u,w)} \frac{1 - q^{-1}e^{\alpha^\vee}(\tau)}{1 - e^{\alpha^\vee}(\tau)},$$

if and only if the opposite Schubert variety $Y(u)$ in the dual complex flag manifold G^\vee/B^\vee is smooth at the torus fixed point wB^\vee .

Thus, in this refined conjecture, the Dynkin diagram of G is not necessarily simply laced, and it is an if and only if statement. The original conjecture follows from this refined one. The “if” direction was claimed by Naruse in [Nar14].

2.3.2. *The analyticity conjecture.* Consider the expansion

$$\varphi_u := \sum_w r_{u,w} f_w \in I(\tau)^I.$$

Then by definition,

$$(5) \quad r_{u,w} := \sum_{u \leq x \leq w} (-1)^{\ell(x) - \ell(u)} m_{x,w}.$$

These two sets of matrix coefficients $r_{u,w}$ and $m_{u,w}$ depend on the unramified character τ , which can be thought of as a point in the complex dual torus T^\vee . Thus, $r_{u,w}$ and $m_{u,w}$ can be regarded as functions on T^\vee , and it is easy to see that they are rational functions on T^\vee . Theorem 2.2 implies that

$$\prod_{\alpha \in S(u,w)} (1 - e^{\alpha^\vee}) m_{u,w}$$

are analytic functions on T^\vee (i.e., having no poles) if the condition in the Theorem is satisfied.

In general, Bump and Nakasuji conjectured the following, see [BN19, Conjecture 1]:

Theorem 2.3 (Analyticity conjecture). *Let $u \leq w$ be two Weyl group elements. Then the functions*

$$\prod_{\alpha \in S(u,w)} (1 - e^{\alpha^\vee}) r_{u,w} \quad , \quad \prod_{\alpha \in S(u,w)} (1 - e^{\alpha^\vee}) m_{u,w}$$

are analytic on the complex dual torus T^\vee .

The goal of this short note is to sketch the proofs of Theorem 2.2 and Theorem 2.3.

3. COMPLEX DUAL SIDE

The idea of the proof of the above conjectures is to give a geometric meaning, in the Langlands dual side, of the matrix coefficients $m_{u,w}$. In this section, we work over the complex numbers \mathbb{C} , and introduce the motivic Chern classes. We give a smoothness criterion for the Schubert varieties using the motivic Chern classes.

3.1. Definition of Motivic Chern classes. In this section, we give the definition of the motivic Chern classes, which is a K-theoretic generalization of the Chern–Schwartz–Macpherson classes in homology [Mac74, Sch65a, Sch65b].

Let X be a quasi-projective, non-singular, complex algebraic variety, with an action of the dual torus T^\vee . Let $K_{T^\vee}(X) := K^0(\text{Coh}_{T^\vee}(X))$ be the T^\vee -equivariant K theory of X , where $\text{Coh}_{T^\vee}(X)$ is the Abelian category of T^\vee -equivariant sheaves on X (see [CG09] for a good reference for equivariant K theory). By definition, the equivariant K group of a point is

$$K_{T^\vee}(\text{pt}) = K^0(\text{Coh}_{T^\vee}(\text{pt})) = K^0(\text{Rep}(T^\vee)) = \mathbb{Z}[T^\vee],$$

where $\text{Rep}(T^\vee)$ is the Abelian category of finite dimensional representation category of T^\vee and $\mathbb{Z}[T^\vee]$ is the regular functions on T^\vee . Since we can tensor any element in $\text{Coh}_T^\vee(X)$ by a finite dimensional T^\vee representation, $K_{T^\vee}(X)$ is a module over $K_{T^\vee}(\text{pt}) = \mathbb{Z}[T^\vee]$. Let $K_{T^\vee}(X)_{\text{loc}} := K_{T^\vee}(X) \otimes_{K_{T^\vee}(\text{pt})} \text{Frac } K_{T^\vee}(\text{pt})$ denote the localized equivariant K theory of X , where $\text{Frac } K_{T^\vee}(\text{pt})$ denotes the fraction field of $K_{T^\vee}(\text{pt})$.

Recall the (relative) motivic Grothendieck group $G_0^{T^\vee}(\text{Var}/X)$ of varieties over X is the free Abelian group generated by isomorphism classes $[f : Z \rightarrow X]$ where Z is a quasi-projective T^\vee -variety and $f : Z \rightarrow X$ is a T^\vee -equivariant morphism modulo the usual additivity relations

$$[f : Z \rightarrow X] = [f : U \rightarrow X] + [f : Z \setminus U \rightarrow X]$$

for $U \subset Z$ an open invariant subvariety.

Let y be a formal variable. The following theorem is proved by Brasselet, Schürmann and Yokura [BSY10, Thm. 2.1] in the non-equivariant case. The equivariant case is proved in [AMSS19, FRW18].

Theorem 3.1. *There exists a unique natural transformation*

$$MC_y : G_0^{T^\vee}(\text{Var}/X) \rightarrow K_{T^\vee}(X)[y]$$

satisfying the following properties:

- (1) *It is functorial with respect to T^\vee -equivariant proper morphisms of non-singular, quasi-projective varieties.*

(2) *It satisfies the following normalization condition*

$$MC_y[id_X : X \rightarrow X] = \lambda_y(T_X^*) := \sum y^i [\wedge^i T_X^*] \in K_{T^\vee}(X)[y].$$

3.2. Two bases in the localized equivariant K theory of the flag variety. In this section, we introduce two bases in the localized equivariant K group of the flag variety $\mathcal{B} = G^\vee/B^\vee$.

The flag variety \mathcal{B} has a natural action of the maximal torus T^\vee , and the fixed points are in one-to-one correspondence with the Weyl group. For any $w \in W$, wB^\vee is the corresponding fixed point, and let $\iota_w : wB^\vee \hookrightarrow \mathcal{B}$ denote the inclusion of the fixed point. For any $\mathcal{F} \in K_{T^\vee}(\mathcal{B})$, let $\mathcal{F}|_w := \iota_w^*(\mathcal{F}) \in K_{T^\vee}(wB^\vee) = K_{T^\vee}(\text{pt}) = \mathbb{Z}[T^\vee]$ denote the restriction of \mathcal{F} to the fixed point wB^\vee . Let $T_w\mathcal{B}$ (resp. $T_w^*\mathcal{B}$) denote the tangent (resp. cotangent) space to \mathcal{B} at the fixed point wB . By the localization theorem [CG09, Chapter 5], the localized equivariant K group $K_{T^\vee}(\mathcal{B})_{loc}$ has a basis $\{[\mathcal{O}_{wB^\vee}] | w \in W\}$, where \mathcal{O}_{wB^\vee} denotes the structure sheaf of the fixed point wB^\vee .

Recall $X(w)^\circ$ (resp. $Y(w)^\circ$) denotes Schubert cells $B^\vee wB^\vee/B^\vee$ (resp. $B^\vee \cdot wB^\vee/B^\vee$). Hence we have $[X(w)^\circ \hookrightarrow \mathcal{B}], [Y(w)^\circ \hookrightarrow \mathcal{B}] \in G_0^{T^\vee}(\text{Var}/\mathcal{B})$. Applying the motivic Chern class transformation MC_y to them, we get

$$MC_y(X(w)^\circ) := MC_y([X(w)^\circ \hookrightarrow \mathcal{B}]) \in K_{T^\vee}(\mathcal{B})[y],$$

and

$$MC_y(Y(w)^\circ) := MC_y([Y(w)^\circ \hookrightarrow \mathcal{B}]) \in K_{T^\vee}(\mathcal{B})[y].$$

Since $X(id)^\circ = B^\vee$ is a point, $MC_y(X(id)^\circ) = [\mathcal{O}_{B^\vee}]$.

By definition, the localization $MC_y(X(w)^\circ)|_w = \lambda_y(T_w^*X(w)) \frac{\lambda_{-1}(T_w^*\mathcal{B})}{\lambda_{-1}(T_w^*X(w))} \in K_{T^\vee}(\text{pt})[y]$, where $T_w^*X(w)$ denotes the tangent space to $X(w)$ at its smooth point wB^\vee , and $\lambda_{-1}(V) := \sum (-1)^i [\wedge^i V] \in K_{T^\vee}(\text{pt})$ for any T^\vee representation V . On the other hand, the $MC_y(X(w)^\circ)$ is supported on $X(w) \subset \mathcal{B}$. Therefore, the transition matrix between $\{MC_y(X(w)^\circ)|_w \in W\}$ and the fixed point basis is upper triangular and the diagonal terms are non-zero. Thus, $\{MC_y(X(w)^\circ)|_w \in W\}$ forms another basis of the localized equivariant K group $K_{T^\vee}(\mathcal{B})_{loc}$.

Here is the example of \mathbb{P}^1 .

Example 3.2. *Let $G^\vee = \text{SL}(2, \mathbb{C})$. Then $\mathcal{B} = \mathbb{P}^1$, $X(id)^\circ = 0$ and $X(s_\alpha)^\circ = \mathbb{P}^1 \setminus \{0\}$. By definition, $MC_y(X(id)^\circ) = [\mathcal{O}_0]$, and $MC_y(X(s_\alpha)^\circ) = MC_y(\mathbb{P}^1) - MC_y(X(id)^\circ) = \lambda_y(T_{\mathbb{P}^1}^*) - [\mathcal{O}_0]$.*

Remark 3.3. *There is another natural basis, the Schubert basis, in the non-localized equivariant K group $K_{T^\vee}(\mathcal{B})$, i.e., the structure sheaves of the Schubert varieties $X(w)$'s. The transition matrix coefficients between the motivic Chern classes and the Schubert classes are conjectured to enjoy some (sign) positivity properties, see [AMSS19]. The cohomological analogue was conjectured by Aluffi and Mihalcea in [AM09, AM16]. The Grassmannian case is proved by J. Huh [Huh16]. The general non-equivariant case is proved in [AMSS17], the proof of which also involves the chomological stable basis of the cotangent bundle [Su17, MO19].*

3.3. Hecke algebra action. In this section, we recall the construction of the Hecke algebra action on $K_{T^\vee}(\mathcal{B})$ due to Lusztig [Lus85], and describe this action on the above two bases.

Recall the finite Hecke algebra $H_W(q)$ is generated over $\mathbb{Z}[q, q^{-1}]$ by elements T_w ($w \in W$) such that $T_u T_v = T_{uv}$ if $\ell(uv) = \ell(u) + \ell(v)$, and $(T_{s_i} + 1)(T_{s_i} - q) = 0$ for a simple reflection s_i in W . For any simple root α_i^\vee of G^\vee , recall $\mathcal{L}_{\alpha_i^\vee} := G^\vee \times_{B^\vee} \mathbb{C}_{\alpha_i^\vee}$. Let P_i^\vee be the corresponding minimal parabolic subgroup containing B^\vee , and let π_i denote the projection

$G^\vee/B^\vee \rightarrow G^\vee/P_i^\vee$. The BGG operator ∂_i is $\pi_i^* \circ \pi_{i,*}$. Form the following two sets of operators in $\text{End}_{K_{T^\vee}(\text{pt})[y]} K_{T^\vee}(\mathcal{B})[y]$

$$\mathcal{T}_i := (1 + y\mathcal{L}_{\alpha_i^\vee})\partial_i - \text{id}, \quad \text{and} \quad \mathcal{T}_i^\vee := \partial_i(1 + y\mathcal{L}_{\alpha_i^\vee}) - \text{id}.$$

On $K_{T^\vee}(\mathcal{B})$, we have a non-degenerate pairing $\langle -, - \rangle$ defined by

$$\langle \mathcal{F}, \mathcal{G} \rangle := \chi_{T^\vee}(\mathcal{B}, \mathcal{F} \otimes \mathcal{G}) \in K_{T^\vee}(\text{pt}),$$

where $\chi_{T^\vee}(-)$ denotes the T^\vee character on the virtual T^\vee -representation $\sum_i H^i(\mathcal{B}, -)$.

Then we have [AMSS19, Lus85]

Theorem 3.4. (1) *The operators \mathcal{T}_i and \mathcal{T}_i^\vee are adjoint to each other, i.e., for any $\mathcal{F}, \mathcal{G} \in K_{T^\vee}(\mathcal{B})$, $\langle \mathcal{T}_i(\mathcal{F}), \mathcal{G} \rangle = \langle \mathcal{F}, \mathcal{T}_i^\vee(\mathcal{G}) \rangle$.*
 (2) *Sending $\mathcal{T}_i \in H_W(q)$ to \mathcal{T}_i (or \mathcal{T}_i^\vee), and q to $-y$ defines an action of $H_W(q)$ on $K_{T^\vee}(\mathcal{B})[y]$.*

The second part of the theorem says that the operators \mathcal{T}_i (resp. \mathcal{T}_i^\vee) satisfy the relations in the Hecke algebra. Therefore, we can form \mathcal{T}_w and \mathcal{T}_w^\vee for any $w \in W$. The motivic Chern classes of the Schubert cells are described by these operators in the following nice formula.

Theorem 3.5. [AMSS19] *For any $w \in W$ and simple root α_i^\vee such that $ws_i > w$, we have*

$$\mathcal{T}_i(MC_y(X(w)^\circ)) = MC_y(X(ws_i)^\circ).$$

In particular,

$$MC_y(X(w)^\circ) = \mathcal{T}_{w^{-1}}([\mathcal{O}_{B^\vee}]).$$

The cohomological statement is proved in [AM16]. Motivated by this and Theorem 3.4(1), we make the following definition.

Definition 3.6. Let $w \in W$. The *dual motivic Chern class* is defined by

$$MC_y^\vee(Y(w)^\circ) := (\mathcal{T}_{w_0w}^\vee)^{-1}(MC_y(Y(w_0))) = (\mathcal{T}_{w_0w}^\vee)^{-1}(\mathcal{O}_{w_0B^\vee}) \in K_T(\mathcal{B})[y, y^{-1}].$$

The name of this class is explained by the following theorem, which is the K-theoretic analogue of [AMSS17, Theorem 5.7].

Theorem 3.7. [AMSS19] *For any $u, v \in W$,*

$$\langle MC_y(X(u)^\circ), MC_y^\vee(Y(v)^\circ) \rangle = \delta_{u,v}(-y)^{\ell(u) - \dim \mathcal{B}} \prod_{\alpha > 0} (1 + ye^{-\alpha^\vee}).$$

Let \mathcal{D} be the Serre duality functor on $K_{T^\vee}(\mathcal{B})$. I.e., for any $\mathcal{F} \in K_{T^\vee}(\mathcal{B})$, let $\mathcal{D}(\mathcal{F}) := \text{RHom}_{\mathcal{O}_{\mathcal{B}}}(\mathcal{F}, \omega_{\mathcal{B}}^\bullet)$, where $\omega_{\mathcal{B}}^\bullet = (-1)^{\dim \mathcal{B}} \mathcal{L}_{2\rho^\vee}$ is the dualizing complex of \mathcal{B} , and $2\rho^\vee$ is the sum of all the positive roots in G^\vee . Extend \mathcal{D} to $K_{T^\vee}(\mathcal{B})[y, y^{-1}]$ by sending y^i to y^{-i} .

Then the class $MC_y^\vee(Y(w)^\circ)$ is related to the original motivic Chern class $MC_y(X(w)^\circ)$ via the following lemma.

Lemma 3.8. *For any $w \in W$, we have*

$$MC_y^\vee(Y(w)^\circ) = \prod_{\alpha > 0} (1 + ye^{-\alpha^\vee}) \frac{\mathcal{D}(MC_y(X(w)^\circ))}{\lambda_y(T^*(\mathcal{B}))} \in K_{T^\vee}(\mathcal{B})_{\text{loc}}[y, y^{-1}].$$

This lemma is quite technical. Its proof in [AMSS19] uses the Maulik–Okounkov stable basis [MO19, Oko17] in the equivariant K theory of the cotangent bundle of the flag variety. To be more specific, we can produce another dual basis for the basis $\{MC_y(X(w)^\circ) | w \in W\}$

using the duality on the stable basis. Comparing this new dual basis with the Hecke dual basis $\{MC_y^\vee(Y(w)^\circ)|w \in W\}$, we arrive at the lemma.

Finally, the Hecke algebra action on the fixed point basis is given by the following lemma.

Lemma 3.9. [AMSS19] *The action of the operator \mathcal{T}_i^\vee on the fixed point basis $\{[\mathcal{O}_{wB^\vee}]|w \in W\}$ is given by*

$$\mathcal{T}_i^\vee([\mathcal{O}_{wB^\vee}]) = -\frac{1+y}{1-e^{-w\alpha_i^\vee}}[\mathcal{O}_{wB^\vee}] + \frac{1+ye^{w\alpha_i^\vee}}{1-e^{-w\alpha_i^\vee}}[\mathcal{O}_{ws_iB^\vee}].$$

3.4. Smoothness criterion. For the smoothness of Schubert varieties, Kumar has the following theorem.

Theorem 3.10 ([Kum96]). *For any $u \leq w \in W$. The opposite Schubert variety $Y(u) \subset \mathcal{B}$ is smooth at wB^\vee if and only if the localization at wB^\vee of the equivariant fundamental class $[Y(u)] \in H_{T^\vee}^*(\mathcal{B})$ is given by:*

$$[Y(u)]|_w = \left(\prod_{\beta > 0, u \not\leq s_\beta w} \beta^\vee \right) \in H_{T^\vee}^*(\text{pt}) = \mathbb{Z}[\text{Lie } T^\vee].$$

We can generalize it using the motivic Chern classes in equivariant K theory.

Theorem 3.11. [AMSS19] *For any $u \leq w \in W$. The opposite Schubert variety $Y(u)$ is smooth at the torus fixed point wB^\vee if and only if*

$$MC_y(Y(u))|_w = \prod_{\alpha > 0, ws_\alpha \geq u} (1 + ye^{w\alpha^\vee}) \prod_{\alpha > 0, u \not\leq ws_\alpha} (1 - e^{w\alpha^\vee}).$$

The if direction follows from the properties of the motivic Chern classes. For the other direction, we can specialize $y = 0$, take the Chern character map, and use Kumar's theorem.

4. THE PROOF

In this section, we outline the proof of Theorem 2.2 and Theorem 2.3. The bridge connecting the p -adic side and the complex side is a shadow of the following well known two geometric realizations of the affine Hecke algebra [CG09, Introduction]

$$K_{G^\vee \times \mathbb{C}^*}(\text{St}) \simeq \mathbb{H} \simeq \mathbb{C}_c[I \backslash G(F) / I],$$

where St is the Steinberg variety.

As before, τ is an unramified regular character of T , which determines a point in the dual torus T^\vee . Let \mathbb{C}_τ denote the evaluation representation at $\tau \in T^\vee$ of $K_{T^\vee}(\text{pt}) = \mathbb{Z}[T^\vee]$.

For any $w \in W$, define

$$b_w := (-1)^{\dim \mathcal{B} - \ell(w)} \prod_{\alpha > 0, w\alpha > 0} \frac{y^{-1} + e^{-w\alpha^\vee}}{1 - e^{w\alpha^\vee}} [\mathcal{O}_{wB^\vee}] \in K_{T^\vee}(\mathcal{B})[y, y^{-1}].$$

The coefficient are determined by the following identity

$$b_w|_w = MC_y^\vee(Y(w)^\circ)|_w.$$

In particular, for the longest element w_0 , $b_{w_0} = MC_y^\vee(Y(w_0)^\circ)$.

We are now ready state the main result which connects the equivariant K-theory of \mathcal{B} to the Iwahori invariants in $I(\tau)$. Endow $K_{T^\vee}(\mathcal{B})[y, y^{-1}]$ with the $H_W(q)$ action by using the operators \mathcal{T}_w^\vee . Then

Theorem 4.1. [AMSS19] *There is a unique left $H_W(q)$ -module homomorphism*

$$\Psi : K_{T^\vee}(G^\vee/B^\vee)[y, y^{-1}] \otimes_{K_{T^\vee}(\text{pt})[y, y^{-1}]} \mathbb{C}_\tau \xrightarrow{\sim} I(\tau)^I,$$

such that:

- $y \mapsto -q$;
- $MC_y^\vee(Y(w)^\circ) \otimes 1 \mapsto \varphi_w$;
- $b_w \otimes 1 \mapsto f_w$.

This type of relation between the equivariant K theory of the flag variety and representation theory of p -adic groups is also studied by Braverman, Kazhdan and Lusztig [BK99, Lus98]. The case of the cotangent bundle is also studied in [SZZ17]. In *loc. cit.*, the authors using this connection to give an equivariant K-theoretic interpretation of the Macdonald's formula for the spherical function [Mac68, Cas80] and the Casselman–Shalika formula for the spherical Whittaker function [CS80].

The proof of this theorem is quite simple. We can define the map Ψ using the first two properties. Then using Equation (1) and Definition 3.6, we check that Ψ is a $H_W(q)$ -module isomorphism. Finally, using the fact that $f_{w_0} = \varphi_{w_0}$, $b_{w_0} = MC_y^\vee(Y(w_0)^\circ)$, Equation (3), Lemma 3.9 and the fact that Ψ is a $H_W(q)$ -module isomorphism, we can check the last property.

Define the dual operator $(-)^{\vee}$ on $K_{T^\vee}(\text{pt})[y, y^{-1}] = \mathbb{Z}[X^*(T^\vee)][y, y^{-1}]$ by $(e^\lambda)^{\vee} = e^{-\lambda}$ for any $\lambda \in X^*(T^\vee)$ and $(y^i)^{\vee} = y^{-i}$ for any $i \in \mathbb{Z}$.

The conjectures of Bump, Nakasuji and Naruse are about the transition matrix coefficients $m_{u,w}$ between the two bases on the p -adic side, using Theorem 4.1 and Lemma 3.8, we get the following geometric interpretation of it.

Proposition 4.2. *For any $w \geq u \in W$, we have*

$$m_{u,w} = \left(\frac{MC_y(Y(u))|_w}{MC_y(Y(w)^\circ)|_w} \right)^{\vee} (\tau),$$

where (τ) means the evaluation at $\tau \in T^\vee$ of the rational function $\frac{MC_y(Y(u))|_w}{MC_y(Y(w)^\circ)|_w} \in \text{Frac}(K_{T^\vee}(\text{pt})[y])$.

Notice that $Y(w)$ is smooth at the torus fixed point wB^\vee . Hence, we have an explicit formula for the denominator $MC_y(Y(w)^\circ)|_w$ by Theorem 3.11. Finally, the factorization conjecture (Theorem 2.2) follows from this proposition and Theorem 3.11.

For the analyticity conjecture (Theorem 2.3), we only need to prove it for $m_{u,w}$ because of Equation (5) and the fact that $S(x, w) \subset S(u, w)$ for any $u \leq x \leq w$. Using Proposition 4.2, Theorem 2.3 is translated into the following

Proposition 4.3. *For any $w \leq u \in W$, the polynomial $MC_y(Y(w)^\circ)|_u \in K_{T^\vee}(\text{pt})[y]$ is divisible by*

$$\prod_{\alpha > 0, u\alpha > 0} (1 + ye^{u\alpha^\vee}) \prod_{\alpha > 0, w \not\leq u s_\alpha < u} (1 - e^{u\alpha^\vee}).$$

This geometric property about motivic Chern classes can be easily proved using an GKM type argument, see [AMSS19].

REFERENCES

- [AM09] Paolo Aluffi and Leonardo Constantin Mihalcea. Chern classes of Schubert cells and varieties. *J. Algebraic Geom.*, 18(1):63–100, 2009.
- [AM16] Paolo Aluffi and Leonardo Constantin Mihalcea. Chern–Schwartz–MacPherson classes for Schubert cells in flag manifolds. *Compos. Math.*, 152(12):2603–2625, 2016.

- [AMSS17] Paolo Aluffi, Leonardo C. Mihalcea, Jörg Schürmann, and Changjian Su. Shadows of characteristic cycles, Verma modules, and positivity of Chern-Schwartz-MacPherson classes of Schubert cells. *arXiv preprint arXiv:1709.08697*, 2017.
- [AMSS19] Paolo Aluffi, Leonardo C Mihalcea, Jörg Schürmann, and Changjian Su. Motivic Chern classes of Schubert cells, Hecke algebras, and applications to Casselman’s problem. *arXiv preprint arXiv:1902.10101*, 2019.
- [Bez16] Roman Bezrukavnikov. On two geometric realizations of an affine Hecke algebra. *Publications mathématiques de l’IHÉS*, 123(1):1–67, 2016.
- [BK99] Alexander Braverman and David Kazhdan. On the Schwartz space of the basic affine space. *Selecta Mathematica*, 5(1):1, 1999.
- [BL00] Sara Billey and Venkatramani Lakshmibai. *Singular loci of Schubert varieties*, volume 182. Springer Science & Business Media, 2000.
- [BN11] Daniel Bump and Maki Nakasuji. Casselman’s basis of Iwahori vectors and the Bruhat order. *Canad. J. Math.*, 63(6):1238–1253, 2011.
- [BN19] Daniel Bump and Maki Nakasuji. Casselmanns basis of Iwahori vectors and Kazhdan–Lusztig polynomials. *Canadian Journal of Mathematics*, pages 1–16, 2019.
- [BSY10] Jean-Paul Brasselet, Jörg Schürmann, and Shoji Yokura. Hirzebruch classes and motivic Chern classes for singular spaces. *J. Topol. Anal.*, 2(1):1–55, 2010.
- [Cas80] William Casselman. The unramified principal series of p -adic groups. I. the spherical function. *Compositio Math.*, 40(3):387–406, 1980.
- [CG09] Neil Chriss and Victor Ginzburg. *Representation theory and complex geometry*. Springer Science & Business Media, 2009.
- [CS80] William Casselman and Joseph Shalika. The unramified principal series of p -adic groups. II. the Whittaker function. *Compositio Mathematica*, 41(2):207–231, 1980.
- [FRW18] Lászlo M. Feher, Richárd Rimanyi, and Andrzej Weber. Motivic Chern classes and K-theoretic stable envelopes. *arXiv preprint arXiv:1802.01503*, 2018.
- [HKP10] Thomas J. Haines, Robert E. Kottwitz, and Amritanshu Prasad. Iwahori-Hecke algebras. *J. Ramanujan Math. Soc.*, 25(2):113–145, 2010.
- [Huh16] June Huh. Positivity of Chern classes of Schubert cells and varieties. *J. Algebraic Geom.*, 25(1):177–199, 2016.
- [KL79] David Kazhdan and George Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.*, 53(2):165–184, 1979.
- [KL87] David Kazhdan and George Lusztig. Proof of the Deligne–Langlands conjecture for Hecke algebras. *Inventiones mathematicae*, 87(1):153–215, 1987.
- [Kum96] Shrawan Kumar. The nil Hecke ring and singularity of Schubert varieties. *Inventiones mathematicae*, 123(3):471–506, 1996.
- [Lan71] Robert P. Langlands. *Euler products*. Yale University Press, New Haven, Conn.-London, 1971. A James K. Whittemore Lecture in Mathematics given at Yale University, 1967, Yale Mathematical Monographs, 1.
- [Lus85] George Lusztig. Equivariant K-theory and representations of Hecke algebras. *Proceedings of the American Mathematical Society*, 94(2):337–342, 1985.
- [Lus98] G Lusztig. Bases in equivariant K-theory. *Representation Theory of the American Mathematical Society*, 2(9):298–369, 1998.
- [Mac68] IG Macdonald. Spherical functions on a p -adic Chevalley group. *Bulletin of the American Mathematical Society*, 74(3):520–525, 1968.
- [Mac74] R. D. MacPherson. Chern classes for singular algebraic varieties. *Ann. of Math. (2)*, 100:423–432, 1974.
- [MO19] D. Maulik and A. Okounkov. Quantum groups and quantum cohomology. *Astérisque*, 408, 2019.
- [Nar14] Hiroshi Naruse. Schubert calculus and Hook formula. *Slides at 73rd Sém. Lothar. Combin., Strobl, Austria*, 2014.
- [NN16] Maki Nakasuji and Hiroshi Naruse. Yang–Baxter basis of Hecke algebra and Casselman’s problem (extended abstract). *DMTCS proc. BC*, pages 935–946, 2016.
- [Oko17] Andrei Okounkov. Lectures on K-theoretic computations in enumerative geometry. In *Geometry of Moduli Spaces and Representation Theory*, volume 24 of *IAS/Park City Mathematics Series*, pages 251–380. 2017.
- [Ree92] Mark Reeder. On certain Iwahori invariants in the unramified principal series. *Pacific journal of mathematics*, 153(2):313–342, 1992.

- [Ree93] Mark Reeder. p -adic Whittaker functions and vector bundles on flag manifolds. *Compositio Math*, 85(1):9–36, 1993.
- [Sch65a] Marie-Hélène Schwartz. Classes caractéristiques définies par une stratification d’une variété analytique complexe. I. *C. R. Acad. Sci. Paris*, 260:3262–3264, 1965.
- [Sch65b] Marie-Hélène Schwartz. Classes caractéristiques définies par une stratification d’une variété analytique complexe. II. *C. R. Acad. Sci. Paris*, 260:3535–3537, 1965.
- [Su17] Changjian Su. Restriction formula for stable basis of the Springer resolution. *Selecta Math. (N.S.)*, 23(1):497–518, 2017.
- [SZ19] Changjian Su and Changlong Zhong. Stable bases of the Springer resolution and representation theory. *arXiv preprint arXiv:1904.06613*, 2019.
- [SZZ17] Changjian Su, Gufang Zhao, and Changlong Zhong. On the K-theory stable bases of the Springer resolution. *arXiv preprint arXiv:1708.08013*, 2017.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ON, M5S 2E4, CANADA
E-mail address: changjiansu@gmail.com