

The Problem. Let $G = \langle g_1, \dots, g_\alpha \rangle$ be a subgroup of S_n , with $n = O(100)$. Before you die, understand G :

1. Compute $|G|$.
2. Given $\sigma \in S_n$, decide if $\sigma \in G$.
3. Write a $\sigma \in G$ in terms of g_1, \dots, g_α .
4. Produce *random* elements of G .

The Commutative Analog. Let $V = \text{span}(v_1, \dots, v_\alpha)$ be a subspace of \mathbb{R}^n . Before you die, understand V .

Solution: Gaussian Elimination. Prepare an empty table,

1	2	3	4	...	n-1	n
---	---	---	---	-----	-----	---

Space for a vector $u_4 \in V$, of the form $u_4 = (0, 0, 0, 1, *, \dots, *)$; 1 := "the pivot".

Feed v_1, \dots, v_α in order. To feed a non-zero v , find its pivotal position i .

1. If box i is empty, put v there.
2. If box i is occupied, find a combination v' of v and u_i that eliminates the pivot, and feed v' .

Non-Commutative Gaussian Elimination

Prepare a mostly-empty table,

(1,1)	I				
(1,2)	(2,2)	I			
(1,3)	(2,3)	(3,3)	I		
⋮				(i,j)	⋮
(1,n)	(2,n)	(3,n)	⋮		(n,n)
					I

Space for a $\sigma_{i,j} \in S_n$ of the form $(1, 2, \dots, i-2, i-1, j, *, *, \dots, *)$
So $\sigma_{i,j}$ fixes $1, \dots, i-1$, sends "the pivot" i to j and goes wild afterwards, and $\sigma_{i,j}^{-1}$ "does sticker j ".

Feed g_1, \dots, g_α in order. To feed a non-identity σ , find its pivotal position i and let $j := \sigma(i)$.

1. If box (i, j) is empty, put σ there.
2. If box (i, j) contains $\sigma_{i,j}$, feed $\sigma' := \sigma_{i,j}^{-1} \sigma$.

The Twist. When done, for every occupied (i, j) and (k, l) , feed $\sigma_{i,j} \sigma_{k,l}$. Repeat until the table stops changing.

Claim. The process stops in our lifetimes, after at most $O(n^6)$ operations. Call the resulting table T .

Claim. Anything fed in T is a monotone product in T :
 f was fed $\Rightarrow f \in M_1 := \{\sigma_{1,j_1} \sigma_{2,j_2} \dots \sigma_{n,j_n} : \forall i, j_i \geq i \text{ and } \sigma_{i,j_i} \in T\}$

Homework Problem 1.
Can you do cosets?



Homework Problem 2.
Can you do categories (groupoids)?

7	9	2	5
1	4	8	3
6	10	11	12
13	14	15	




The Results

```
In[3]:= (Feed[#]; Product[1 + Length[Select[Range[n], Head[s[i, #]] == P &]], {i, n}]) & /@ gs
Out[3]= {4, 16, 159993501696000, 21119142223872000, 43252003274489856000, 43252003274489856000}
```

<http://www.math.toronto.edu/~drorbn/Talks/Mathcamp-0907/> and links there

1	2	3						
4	5	6						
7	8	9						
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36
37	38	39						
40	41	42						
43	44	45						
46	47	48						
49	50	51						
52	53	54						

Based on an algorithm by



Otto Schreier Charles Sims
See also *Permutation Group Algorithms* by Akos Seress.

The Generators

```
In[1]:= gs = {
purple = P[18, 27, 36, 4, 5, 6, 7, 8, 9, 3, 11, 12, 13, 14, 15, 16, 17,
45, 2, 20, 21, 22, 23, 24, 25, 26, 44, 1, 29, 30, 31, 32, 33, 34, 35, 43,
37, 38, 39, 40, 41, 42, 10, 19, 28, 52, 49, 46, 53, 50, 47, 54, 51, 48],
white = P[1, 2, 3, 4, 5, 6, 16, 25, 34, 10, 11, 9, 15, 24, 33, 39, 17,
18, 19, 20, 8, 14, 23, 32, 38, 26, 27, 28, 29, 7, 13, 22, 31, 37, 35, 36,
12, 21, 30, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54],
green = P[1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18,
19, 20, 21, 22, 23, 24, 25, 26, 27, 31, 32, 33, 34, 35, 36, 48, 47, 46,
39, 42, 45, 38, 41, 44, 37, 40, 43, 30, 29, 28, 49, 50, 51, 52, 53, 54],
blue = P[3, 6, 9, 2, 5, 8, 1, 4, 7, 54, 53, 52, 10, 11, 12, 13, 14, 15,
19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36,
37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 18, 17, 16],
red = P[13, 2, 3, 22, 5, 6, 31, 8, 9, 12, 21, 30, 37, 14, 15, 16, 17,
18, 11, 20, 29, 40, 23, 24, 25, 26, 27, 10, 19, 28, 43, 32, 33, 34, 35,
36, 46, 38, 39, 49, 41, 42, 52, 44, 45, 1, 47, 48, 4, 50, 51, 7, 53, 54],
yellow = P[1, 2, 48, 4, 5, 51, 7, 8, 54, 10, 11, 12, 13, 14, 3, 18, 27,
36, 19, 20, 21, 22, 23, 6, 17, 26, 35, 28, 29, 30, 31, 32, 9, 16, 25, 34,
37, 38, 15, 40, 41, 24, 43, 44, 33, 46, 47, 39, 49, 50, 42, 52, 53, 45]
};
```

Theorem. $G = M_1$.

G^{-1} is more fun!

$G = M_1 := \{\sigma_{1,j_1} \sigma_{2,j_2} \dots \sigma_{n,j_n} : \forall i, j_i \geq i \text{ and } \sigma_{i,j_i} \in T\}$.

Proof. The inclusions $M_1 \subset G$ and $\{g_1, \dots, g_\alpha\} \subset M_1$ are obvious. The rest follows from the following

Lemma. M_1 is closed under multiplication.

Proof. By backwards induction. Let

$$M_k := \{\sigma_{k,j_k} \dots \sigma_{n,j_n} : \forall i \geq k, j_i \geq i \text{ and } \sigma_{i,j_i} \in T\}.$$

Clearly $M_n M_n \subset M_n$. Now assume that $M_5 M_5 \subset M_5$ and show that $M_4 M_4 \subset M_4$. Start with $\sigma_{8,j} M_4 \subset M_4$:

$$\begin{aligned} \sigma_{8,j}(\sigma_{4,j_4} M_5) &\stackrel{1}{=} (\sigma_{8,j} \sigma_{4,j_4}) M_5 \stackrel{2}{\subset} M_4 M_5 \\ &\stackrel{3}{=} \sigma_{4,j_4} (M_5 M_5) \stackrel{4}{\subset} \sigma_{4,j_4} M_5 \subset M_4 \end{aligned}$$

(1: associativity, 2: thank the twist, 3: associativity and tracing i_4 , 4: induction). Now the general case


$$(\sigma_{4,j'_4} \sigma_{5,j'_5} \dots)(\sigma_{4,j_4} \sigma_{5,j_5} \dots)$$

falls like a chain of dominos.

Problem Solved!

A Demo Program

```
1 In[2]:= ($RecursionLimit = 2^16;
2 n = 54;
3 P /: p_P ** P[a_] := p[[a]];
4 Inv[p_P] := P @@ Ordering[p];
5 Feed[P @@ Range[n]] := Null;
6 Feed[p_P] := Module[{i, j},
7 For[i = 1, p[[i]] == i, ++i];
8 j = p[[i]];
9 If[Head[s[i, j]] == P,
10 Feed[Inv[s[i, j]] ** p],
11 (* Else *) s[i, j] = p;
12 Do[If[Head[s[k, 1]] == P,
13 Feed[s[i, j] ** s[k, 1]];
14 Feed[s[k, 1] ** s[i, j]]
15 ], {k, n}, {1, n}]]];
16
```



Enter

that's cool!

Enter