

**COURSE INTRODUCTION:
WHAT HAPPENS TO A QUANTUM PARTICLE ON A PENDULUM $\frac{\pi}{2}$
SECONDS AFTER IT IS TOSSED IN?**

DROR BAR-NATAN

Follows a lecture given by the author in the “trivial notions” seminar in Harvard on April 29, 1989.

ABSTRACT. This subject is the best one-hour introduction I know for the mathematical techniques that appear in quantum mechanics — in one short lecture we start with a meaningful question, visit Schrödinger’s equation, operators and exponentiation of operators, Fourier analysis, path integrals, the least action principle, and Gaussian integration, and at the end we land with a meaningful and interesting answer.

CONTENTS

1. The Question	1
2. The Solution	2
3. The Lemmas	3
4. The Morals	5

1. THE QUESTION

Let the complex valued function $\psi = \psi(t, x)$ be a solution of the Schrödinger equation

$$\frac{\partial \psi}{\partial t} = -i \left(-\frac{1}{2} \Delta_x + \frac{1}{2} x^2 \right) \psi \quad \text{with} \quad \psi|_{t=0} = \psi_0.$$

What is $\psi|_{t=T=\frac{\pi}{2}}$?

In fact, the major part of our discussion will work just as well for the general Schrödinger equation,

$$\frac{\partial \psi}{\partial t} = -iH\psi, \quad H = -\frac{1}{2} \Delta_x + V(x), \quad \psi|_{t=0} = \psi_0, \quad \text{arbitrary } T,$$

where:

- ψ is the “wave function”, with $|\psi(t, x)|^2$ representing the probability of finding our particle at time t in position x .
- H is the “energy”, or the “Hamiltonian”.
- $-\frac{1}{2} \Delta_x$ is the “kinetic energy”.
- $V(x)$ is the “potential energy at x ”.

Date: This edition: March 27, 1999; First edition: April 29, 1989.

2. THE SOLUTION

The equation $\frac{\partial \psi}{\partial t} = -iH\psi$ with $\psi|_{t=0} = \psi_0$ formally implies

$$\psi(T, x) = (e^{-iTH}\psi_0)(x) = \left(e^{i\frac{T}{2}\Delta - iTV}\psi_0\right)(x).$$

By Lemma 3.1 with $n = 10^{58} + 17$ and setting $x_n = x$ we thus get:

$$\psi(T, x) = \left(e^{i\frac{T}{2n}\Delta} e^{-i\frac{T}{n}V} e^{i\frac{T}{2n}\Delta} e^{-i\frac{T}{n}V} \dots e^{i\frac{T}{2n}\Delta} e^{-i\frac{T}{n}V} \psi_0\right)(x_n).$$


Now using Lemmas 3.2 and 3.3 we find that this is: (c denotes the ever-changing universal fixed numerical constant)

$$c \int dx_{n-1} e^{i\frac{(x_n - x_{n-1})^2}{2T/n}} e^{-i\frac{T}{N}V(x_{n-1})} \dots \int dx_1 e^{i\frac{(x_2 - x_1)^2}{2T/n}} e^{-i\frac{T}{N}V(x_1)} \int dx_0 e^{i\frac{(x_1 - x_0)^2}{2T/n}} e^{-i\frac{T}{N}V(x_0)} \psi_0(x_0).$$

Repackaging, we get

$$\psi(T, x) = c \int dx_0 \dots dx_{n-1} \exp\left(i\frac{T}{2n} \sum_{k=1}^n \left(\frac{x_k - x_{k-1}}{T/n}\right)^2 - i\frac{T}{n} \sum_{k=0}^{n-1} V(x_k)\right) \psi_0(x_0).$$

Now comes the big novelty. keeping in mind the picture



and replacing Riemann sums by integrals, we can write

$$\psi(T, x) = c \int dx_0 \int_{W_{x_0 x_n}} \mathcal{D}x \exp\left(i \int_0^T dt \left(\frac{1}{2}\dot{x}^2(t) - V(x(t))\right)\right) \psi_0(x_0),$$

where $W_{x_0 x_n}$ denotes the space of paths that begin at x_0 and end at x_n ,

$$W_{x_0 x_n} = \{x : [0, T] \rightarrow \mathbb{R} : x(0) = x_0, x(T) = x_n\},$$

and $\mathcal{D}x$ is the formal “path integral measure”.

This is a good time to introduce the “action” \mathcal{L} :

$$\mathcal{L}(x) := \int_0^T dt \left(\frac{1}{2}\dot{x}^2(t) - V(x(t))\right).$$

With this notation,

$$\psi(T, x) = c \int dx_0 \psi_0(x_0) \int_{W_{x_0 x_n}} \mathcal{D}x e^{i\mathcal{L}(x)}.$$

Let x_c denote the path on which $\mathcal{L}(x)$ attains its minimum value, write $x = x_c + x_q$ with $x_q \in W_{00}$, and get

$$\psi(T, x) = c \int dx_0 \psi_0(x_0) \int_{W_{00}} \mathcal{D}x_q e^{i\mathcal{L}(x_c+x_q)}.$$

In our particular case \mathcal{L} is quadratic in x , and therefore $\mathcal{L}(x_c + x_q) = \mathcal{L}(x_c) + \mathcal{L}(x_q)$ (this uses the fact that x_c is an extremal of \mathcal{L} , of course). Plugging this into what we already have, we get:

$$\psi(T, x) = c \int dx_0 \psi_0(x_0) \int_{W_{00}} \mathcal{D}x_q e^{i\mathcal{L}(x_c)+i\mathcal{L}(x_q)} = c \int dx_0 \psi_0(x_0) e^{i\mathcal{L}(x_c)} \int_{W_{00}} \mathcal{D}x_q e^{i\mathcal{L}(x_q)}.$$

Now this is excellent news, because the remaining path integral over W_{00} does not depend on x_0 or x_n , and hence it is a constant! Allowing c to change its value from line to line, we get

$$\psi(T, x) = c \int dx_0 \psi_0(x_0) e^{i\mathcal{L}(x_c)}.$$

Lemma 3.4 now shows us that $x_c(t) = x_0 \cos t + x_n \sin t$. An easy explicit computation gives $\mathcal{L}(x_c) = -x_0 x_n$, and we arrive at our final result:

$$\psi\left(\frac{\pi}{2}, x\right) = c \int dx_0 \psi_0(x_0) e^{-ix_0 x_n}.$$

Notice that this is precisely the formula for the Fourier transform of ψ_0 ! That is, the answer to the question in the title of this document is “the particle gets Fourier transformed”, whatever that may mean.

3. THE LEMMAS

Lemma 3.1. *For any two matrices A and B ,*

$$e^{A+B} = \lim_{n \rightarrow \infty} (e^{A/n} e^{B/n})^n.$$

Proof. (sketch) Using Taylor expansions, we see that $e^{\frac{A+B}{n}}$ and $e^{A/n} e^{B/n}$ differ by terms at most proportional to c/n^2 . Raising to the n th power, the two sides differ by at most $O(1/n)$, and thus

$$e^{A+B} = \lim_{n \rightarrow \infty} \left(e^{\frac{A+B}{n}} \right)^n = \lim_{n \rightarrow \infty} (e^{A/n} e^{B/n})^n,$$

as required. □

Lemma 3.2.

$$(e^{itV} \psi_0)(x) = e^{itV(x)} \psi_0(x).$$

□

Lemma 3.3.

$$\left(e^{i\frac{t}{2}\Delta} \psi_0 \right)(x) = c \int dx' e^{i\frac{(x-x')^2}{2t}} \psi_0(x').$$

Proof. In fact, the left hand side of this equality is just a solution $\psi(t, x)$ of Schrödinger's equation with $V = 0$:

$$\frac{\partial \psi}{\partial t} = \frac{i}{2} \Delta_x \psi, \quad \psi|_{t=0} = \psi_0.$$

Taking the Fourier transform $\tilde{\psi}(t, p) = \frac{1}{\sqrt{2\pi}} \int e^{-ipx} \psi(t, x) dx$, we get the equation

$$\frac{\partial \tilde{\psi}}{\partial t} = -i \frac{p^2}{2} \tilde{\psi}, \quad \tilde{\psi}|_{t=0} = \tilde{\psi}_0.$$

For a fixed p , this is a simple first order linear differential equation with respect to t , and thus,

$$\tilde{\psi}(t, p) = e^{-i \frac{tp^2}{2}} \tilde{\psi}_0(p).$$

Taking the inverse Fourier transform, which takes products to convolutions and Gaussians to other Gaussians, we get what we wanted to prove. \square

Lemma 3.4. *With the notation of Section 2 and at the specific case of $V(x) = \frac{1}{2}x^2$ and $T = \frac{\pi}{2}$, we have*

$$x_c(t) = x_0 \cos t + x_n \sin t.$$

Proof. If x_c is a critical point of \mathcal{L} on $W_{x_0 x_n}$, then for any $x_q \in W_0$ there should be no term in $\mathcal{L}(x_c + \epsilon x_q)$ which is linear in ϵ . Now recall that

$$\mathcal{L}(x) = \int_0^T dt \left(\frac{1}{2} \dot{x}^2(t) - V(x(t)) \right),$$

so using $V(x_c + \epsilon x_q) \sim V(x_c) + \epsilon x_q V'(x_c)$ we find that the linear term in ϵ in $\mathcal{L}(x_c + \epsilon x_q)$ is

$$\int_0^T dt (\dot{x}_c \dot{x}_q - V'(x_c) x_q).$$

Integrating by parts and using $x_q(0) = x_q(T) = 0$, this becomes

$$\int_0^T dt (-\ddot{x}_c - V'(x_c)) x_q.$$

For this integral to vanish independently of x_q , we must have $-\ddot{x}_c - V'(x_c) \equiv 0$, or

$$\ddot{x}_c = -V'(x_c). \quad \left(\text{This is the famous } F = ma \text{ of Newton's, and we} \right. \\ \left. \text{have just rediscovered the principle of least action!} \right)$$

In our particular case this boils down to the equation

$$\ddot{x}_c = -x_c, \quad x_c(0) = x_0, \quad x_c(\pi/2) = x_n,$$

whose unique solution is displayed in the statement of this lemma. \square

4. THE MORALS

- Schrödinger's equation is related to some infinite dimensional "path integrals".
- These path integrals can sometime be evaluated, with interesting and useful results.
- The Fourier transform fits within some 1-parameter family of unitary operators, defined by $U_t = e^{-itH}$ for t other than $\frac{\pi}{2}$. The same techniques lead to explicit formulas for U_t for any t .
- We'd better work harder, to understand how all of this fits into some bigger coherent picture.

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY, GIV'AT-RAM, JERUSALEM 91904, ISRAEL
E-mail address: drorbn@math.huji.ac.il