

# ON THE VASSILIEV KNOT INVARIANTS

DROR BAR-NATAN

*Appeared in Topology* **34** (1995) 423–472.

ABSTRACT. The theory of knot invariants of finite type (Vassiliev invariants) is described. These invariants turn out to be at least as powerful as the Jones polynomial and its numerous generalizations coming from various quantum groups, and it is conjectured that these invariants are precisely as powerful as those polynomials. As invariants of finite type are much easier to define and manipulate than the quantum group invariants, it is likely that in attempting to classify knots, invariants of finite type will play a more fundamental role than the various knot polynomials.

## CONTENTS

1. Introduction	2
1.1. A sketchy introduction to the introduction	2
1.2. Acknowledgement	3
1.3. Weight systems and invariants of finite type	3
1.4. There are many invariants of finite type	5
1.5. The algebra $\mathcal{A}$ of diagrams	6
1.6. The primitive elements of $\mathcal{A}$	8
1.7. How big are $\mathcal{A}$ , $\mathcal{W}$ and $\mathcal{P}$ ?	9
1.8. Odds and ends	10
1.9. Summary of spaces and maps	10
2. The basic constructions	10
2.1. The classical knot polynomials	10
2.2. Constructing a weight system from a Vassiliev knot invariant	11
2.3. Framed links	12
2.4. Constructing invariant tensors from Lie algebraic information	13
3. The algebra $\mathcal{A}$ of diagrams	15
3.1. Proof of theorem 6	15
3.2. Proof of theorem 7	16
4. Kontsevich's Knizhnik-Zamolodchikov construction	21
4.1. Connections, curvature, and holonomy	21
4.2. The formal Knizhnik-Zamolodchikov connection	22
4.3. Kontsevich's integral invariants	23
4.4. Proof of theorem 1	27

---

*Date:* This edition: February 13, 2007; First edition: August 1992.

I wish to thank the Weizmann Institute of Science for their hospitality during the preparation of this paper, and the NSF for financial support.

This preprint is available electronically at <http://www.math.toronto.edu/~drorbn>.

5.	The primitive elements of $\mathcal{A}$	28
5.1.	Theorem 8 from the perspective of Lie algebras	28
5.2.	Proof of theorem 8	29
5.3.	Proof of part (2) of theorem 9	34
6.	How big are $\mathcal{A}$ , $\mathcal{W}$ and $\mathcal{P}$ ?	34
6.1.	Numerical results	34
6.2.	Marked surfaces	35
6.3.	The classical Lie algebras	37
6.4.	Marked surfaces and the classical Lie algebras	43
6.5.	Marked surfaces and the classical knot polynomials	48
6.6.	More on conjecture 2	48
7.	Odds and ends	49
7.1.	Some questions	49
7.2.	Some bad news	50
7.3.	Bibliographical remarks	50
	References	51

## 1. INTRODUCTION

1.1. **A sketchy introduction to the introduction.** Any numerical knot invariant  $V$  can be inductively extended to be an invariant  $V^{(m)}$  of immersed circles that have exactly  $m$  *transversal* self intersections using the formulas

$$V^{(0)} = V,$$

$$V^{(m)} \left( \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) = V^{(m-1)} \left( \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} \right) - V^{(m-1)} \left( \begin{array}{c} \diagdown \diagup \\ \diagdown \diagup \end{array} \right). \quad (1)$$

We know from multi-variable calculus that differences are cousins of derivatives, and so we wish to think of (1) as the definition of the  $m$ th partial derivative of a knot invariant in terms of its  $(m-1)$ st partial derivatives. (In a knot projection there can be many crossings, and so one can ‘differentiate’ with respect to many different ‘variables’. Hence we think of (this corner of) knot theory as *multi*-variable calculus).

An invariant  $V$  is called “(A *Vassiliev invariant*) of type  $m$ ” if its  $(m+1)$ st derivative vanishes identically<sup>1</sup>. Just like in calculus,  $V^{(m+1)} \equiv 0$  implies that  $V^{(m)}$  is a constant, in some sense. This ‘constant’ is actually a collection of constants corresponding to the different possible partial derivatives of  $V$ . From the special nature of the “function”  $V$  (it is a knot invariant), it follows that there are some relations among these constants. A collection of constants satisfying these relations is called a *weight system*. The purpose of this paper is to discuss the following questions:

- Which of the well-known knot invariants are Vassiliev invariants? (See theorems 2, 3, and 5).
- What are weight systems in a more precise language? What are the relations that a weight system has to satisfy? (See definitions 1.6, 1.7, 1.9, and theorem 6).

---

<sup>1</sup>Having the analogy with calculus in mind, a better name would have been “a polynomial invariant”. But unfortunately, this name is already used for something else.

- Is it always possible to ‘integrate’ a weight system  $m$  times and get a knot invariant? (See theorem 1).
- Knot invariants can be added and multiplied. Knots can be multiplied (the operation of ‘connected sum’) and therefore knot invariants can be co-multiplied. What are the operations that one can perform on a weight system? Is the space of weight systems a Hopf algebra? (See theorems 7, 8, and 9).
- Can one classify all weight systems? (See theorems 4, 10, 11, 12 and conjectures 1 and 2).

Quite unexpectedly, given a representation of a semi-simple Lie algebra there is a simple construction of a weight system. In fact, a closer look at weight systems reveals that for many purposes they are as good as Lie algebras, despite their a-priori different appearance. The following further questions therefore arise:

- Which part of the theory of Lie algebras can be translated to the language of weight systems? (See *e.g.* section 5).
- Do *all* weight systems come from Lie algebras? (See the statistics in section 6.1, theorem 11 and conjectures 1 and 2).

The following outstanding problem *will not* be discussed in this paper:

**Problem 1.1.** Is there an analog of Taylor’s theorem in our context — can an arbitrary knot invariant be approximated by Vassiliev invariants? Do Vassiliev invariants separate knots?

An affirmative answer to the above question will, of course, give tremendous further impetus to studying Vassiliev invariants.

1.2. **Acknowledgement.** I wish to thank the many people who taught me about Vassiliev invariants, and the many people who were patient enough to listen to my own ideas about the subject. Among others, these include: V. I. Arnold, A-B. Berger, J. Birman, R. Bott, S. Chmutov, S. Duzhin, P. Lee, X-S. Lin, G. Masbaum, G. Wetzel, S. Willerton, and E. Witten. I am particularly grateful to M. Kontsevich, for inventing a considerable part of the theory described in these pages, for our many conversations, and for his careful reading of an earlier version of this paper.

1.3. **Weight systems and invariants of finite type.** Let  $\mathbf{F}$  be a field of characteristic zero<sup>2</sup>. Any  $\mathbf{F}$ -valued invariant  $V$  of oriented knots in an oriented<sup>3</sup> three dimensional manifold  $M^3$  can be extended to be an invariant of immersed circles in  $M^3$ , which are allowed to have some *transversal* self intersections, using the following definition:

**Definition 1.2.** Suppressing the superfluous superscripts of (1), set

$$V \left( \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) = V \left( \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} \right) - V \left( \begin{array}{c} \diagup \diagdown \\ \diagdown \diagdown \end{array} \right). \quad (2)$$

As usual in knot theory and as will be the standard throughout the rest of this paper, when we write  $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$ ,  $\begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array}$  or  $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagdown \end{array}$ , we think of them as parts of bigger graphs which are identical outside of a small sphere, inside of which they look as in the figures. Notice also that in an

---

<sup>2</sup>In fact, a considerable fraction of the results proven here are true even with the field  $\mathbf{F}$  replaced by an arbitrary Abelian group; see problem 7.3.

<sup>3</sup>Przytycki [34] noticed that Vassiliev invariants can be defined on non-orientable manifolds as well.

oriented manifold the notion of an overcrossing (undercrossing) is well defined and does not depend on a choice of a planar projection. See for example [21, pp. 13].

**Definition 1.3.** Let  $m$  be a non-negative integer. An invariant  $V$  of oriented knots in an oriented three dimensional manifold  $M^3$  is called *an invariant of type  $m$* , if  $V$  vanishes on “knots” that have more than  $m$  self intersections:

$$\left( \begin{array}{c} \text{the} \\ \text{Birman-Lin} \\ \text{condition} \end{array} \right) \quad V \left( \underbrace{\begin{array}{c} \diagup \diagdown \quad \diagdown \diagup \quad \dots \quad \diagup \diagdown \\ \diagdown \diagup \quad \diagup \diagdown \quad \dots \quad \diagdown \diagup \end{array} }_{> m} \right) = 0.$$

An invariant  $V$  of oriented knots in an oriented three dimensional manifold  $M^3$  is called a *Vassiliev invariant* [42, 43], or *an invariant of finite type*, if it is of type  $m$  for some  $m \in \mathbf{N}$ . The space  $\mathcal{V}$  of all Vassiliev invariants is *filtered*, with  $\mathcal{V}_m = \{\text{invariants of type } m\}$ .

*Remark 1.4.* Vassiliev’s original approach is very different from ours. He derives his class of invariants from certain topological considerations that involve viewing the space of all embeddings of  $S^1$  into  $\mathbf{R}^3$  as the complement of the space of singular immersions in the space of all immersions. Our definition follows the ‘Birman-Lin axioms’ of [10] and can be extracted from Vassiliev’s more involved theory.

Throughout the rest of this paper we will consider only the case of  $M^3 = \mathbf{R}^3$ . Let us now turn to the study of the  $m$ th ‘derivative’ of a Vassiliev invariant:

**Definition 1.5.** A *chord diagram* (CD) is an oriented circle with finitely many chords marked on it, regarded up to orientation preserving diffeomorphisms of the circle. Denote the collection of all chord diagrams by  $\mathcal{D}^c$  ( $c$  for ‘chord’). This collection is naturally graded by the number of chords in such a diagram. Denote the piece of degree  $m$  of  $\mathcal{D}^c$  by  $\mathcal{G}_m \mathcal{D}^c$ .  $\mathcal{G}_m \mathcal{D}^c$  is simply the collection of all chord diagram having precisely  $m$  chords.

By convention, we will always orient the circle in a chord diagram counterclockwise, and always use dashed lines for the chords. For examples, the collection  $\mathcal{G}_3 \mathcal{D}^c$  of chord diagrams of degree 3 is  $\mathcal{G}_3 \mathcal{D}^c = \{ \text{⊕}, \text{⊗}, \text{⊗}, \text{⊕}, \text{⊗} \}$ .

**Definition 1.6.** An  $\mathbf{F}$ -valued *weight system* of degree  $m$  is a function  $W : \mathcal{G}_m \mathcal{D}^c \rightarrow \mathbf{F}$  having the following properties:

- (1) If  $D \in \mathcal{G}_m \mathcal{D}^c$  has an *isolated chord* — a chord that does not intersect any other chord in  $D$ , then  $W(D) = 0$ . This property is called *framing independence*.
- (2) Whenever four diagrams  $S$ ,  $E$ ,  $W$ , and  $N$  differ only as shown in figure 1, their weights satisfy


$$W(S) - W(E) = -W(W) + W(N). \tag{3}$$

This property is called the  $4T$  (4 Term) relation.

Let  $\mathcal{W}$  denote the graded vector space of all weight systems.

The first theorem of this paper says that over the real numbers the above two notions are essentially equivalent:

**Theorem 1.** *Over  $\mathbf{R}$ , the graded vector space associated with the filtered vector space  $\mathcal{V}$  of Vassiliev invariants is  $\mathcal{W}$ . More precisely:*



**Figure 1.** The diagrams  $S$ ,  $E$ ,  $W$ , and  $N$ . (The dotted arcs represent parts of the diagrams that are not shown in the figure. These parts are assumed to be the same in all four diagrams)

- (1) (Proof on page 11) For a given non-negative integer  $m$  there is a naturally defined map  $V \mapsto W_m(V)$  which to any given type  $m$   $\mathbf{F}$ -valued Vassiliev invariant  $V$  associates a degree  $m$   $\mathbf{F}$ -valued weight system  $W_m(V)$ .
- (2) (Kontsevich [25]) (Proof on page 27) Restricting to the case  $\mathbf{F} = \mathbf{R}$ , there is a naturally defined map  $W \mapsto V(W)$  which to any given  $\mathbf{R}$ -valued weight system of degree  $m$  associates a type  $m$   $\mathbf{R}$ -valued Vassiliev invariant  $V(W)$ .
- (3) (Proof on page 27) The above two maps are nearly each other's inverse — for any given  $W$  one has  $W = W_m(V(W))$ , and for any given  $V$  the invariants  $V$  and  $V(W_m(V))$  differ by a knot invariant of type  $m - 1$ .

#### 1.4. There are many invariants of finite type.

**Theorem 2.** (Bar-Natan [5, 6]) (Proof on page 10). Each coefficient of the Conway polynomial is an invariant of finite type.

**Theorem 3.** (Birman-Lin, [10]) (Proof on page 11) After a suitable change of variables, each coefficient in the Taylor expansion of the Jones [20], HOMFLY<sup>4</sup> [19], and Kauffman [22] polynomials is an invariant of finite type.

**Theorem 4.** (Bar-Natan [5, 6]) Let  $\mathfrak{g}$  be a finite dimensional Lie algebra over a field  $\mathbf{F}$ ,  $t$  an Ad-invariant symmetric non-degenerate bilinear form on  $\mathfrak{g}$ , and  $R$  a finite dimensional representation of  $\mathfrak{g}$ . Let  $m$  be a non-negative integer.

- (1) (Proof on page 14) Given this information, there is a natural construction of an associated functional<sup>5</sup>  $W_{\mathfrak{g},R,m} : \mathcal{G}_m \mathcal{D}^c \rightarrow \mathbf{F}$  satisfying the 4T identity (3).
- (2) (Proof on page 20) There is a canonical way to ‘renormalize’  $W_{\mathfrak{g},R,m}$  to a weight system  $\hat{W}_{\mathfrak{g},R,m}$ .

**Theorem 5.** (Lin, [28]) After a suitable change of variables, each coefficient in the Taylor expansion of the Reshetikhin-Turaev ‘quantum-group’ invariants [35, 36, 37, 41] corresponding to a quantization of a triple  $\{\mathfrak{g}, t, R\}$  as in theorem 4 is an invariant of finite type. Furthermore (Piunikhin, [33]), the weight systems underlying those invariants are precisely those constructed in theorem 4. (See remarks 2.2, 2.3, and 4.8 and problem 4.9).

<sup>4</sup>The HOMFLY polynomial is named after the initials of 6 of its discoverers, Hoste, Ocneanu, Millet, Freyd, Lickorish and Yetter. In fact, it was also discovered simultaneously by Przytycki and Traczyk and therefore, following L. Rudolph, it should more accurately be called “the LYMPH-TOFU polynomial”, with the last U standing for the Unknown further discoverers.


<sup>5</sup> $t$  is normally taken to be given by matrix trace in the defining representation, and is always suppressed from the notation.

1.5. **The algebra  $\mathcal{A}$  of diagrams.** For some purposes, it is better to consider weight systems as linear functionals over a graded vector space  $\mathcal{A}^c$ .

**Definition 1.7.**  $\mathcal{A}^c$  is the quotient space

$$\mathcal{A}^c = \text{span}(\mathcal{D}^c) / \text{span} \{ \text{all } 4T \text{ relations} \} .$$

Namely,  $\mathcal{A}^c$  is the quotient of the vector space freely generated by all chord diagrams by the subspace spanned by all relations of the form:



As the  $4T$  relation is homogeneous,  $\mathcal{A}^c$  inherits a gradation from  $\mathcal{D}^c$ .

Clearly, a weight system of degree  $m$  is just a linear functional on  $\mathcal{G}_m \mathcal{A}^c$  which vanishes on classes represented by a chord diagram that has an isolated chord.


The space  $\mathcal{A}^c$  has an equivalent description as follows:

**Definition 1.8.** A *Chinese Character Diagram*<sup>6</sup> (CCD) is a connected graph made of a single oriented circle and a certain number of unoriented dashed lines, which are allowed to meet in two types of trivalent vertices:

- (1) *Internal vertices* in which three dashed lines meet. These vertices are *oriented* — one of the two possible cyclic orderings of the arcs emanating from such a vertex is specified.
- (2) *External vertices* in which a dashed line ends on the circle.

The collection of all Chinese character diagrams will be denoted by  $\mathcal{D}^t$  ( $t$  for ‘trivalent’).

By convention, the circle in a CCD is always oriented counterclockwise, and so are the internal vertices. Also, as higher than trivalent vertices are not allowed in a CCD, what appears in a picture to be a vertex of order 4 is not a vertex at all — it is just a pair of arcs passing each other without intersection. An example is in figure 2.



**Figure 2.** A Chinese Character Diagram (CCD) together with the conventional way of drawing it, as outlined in the text.


Clearly, there is an even number of vertices in a CCD. We will use *half* the number of vertices to grade  $\mathcal{D}^t$ . A chord diagram is also a CCD, and it is easy to see that the gradations are compatible:  $\mathcal{G}_m \mathcal{D}^c \subseteq \mathcal{G}_m \mathcal{D}^t$ .

<sup>6</sup>This joke is due to Morton Brown. When the circle is stripped off a Chinese character diagram such as , what remains is a ‘Chinese character’ like .

**Definition 1.9.** Let the vector space  $\mathcal{A}^t$  be the quotient


$$\mathcal{A}^t = \text{span}(\mathcal{D}^t) / \text{span}\{\text{all } STU \text{ relations}\}.$$

An  $STU$  relation is a relation of the form  $S = T - U$ , where the diagrams  $S$ ,  $T$ , and  $U$  are identical outside of a small circle, inside of which they look as in figure 3. As the  $STU$  relation is homogeneous,  $\mathcal{A}^t$  inherits a gradation from  $\mathcal{D}^t$ .



**Figure 3.** The diagrams  $S$ ,  $T$ , and  $U$ , and the  $STU$  relation.


*Remark 1.10.* Notice that the  $4T$  relation holds in  $\mathcal{A}^t$ :



(Both equalities in the above equation hold in  $\mathcal{A}^t$  because of the  $STU$  relation). This implies that the inclusion  $\mathcal{D}^c \hookrightarrow \mathcal{D}^t$  descends to a linear map  $\phi^c : \mathcal{A}^c \rightarrow \mathcal{A}^t$ .

**Theorem 6.** (*Proof on page 15*) The map  $\phi^c : \mathcal{A}^c \rightarrow \mathcal{A}^t$  is an isomorphism. Furthermore, the following two identities hold in  $\mathcal{A}^t$ :

(1) *Antisymmetry of internal vertices:*



(2) *The IHX identity:*

$$\text{IHX diagram} = \text{H diagram} - \text{X diagram}. \tag{4}$$

As  $\mathcal{A}^t$  and  $\mathcal{A}^c$  are anyway isomorphic, we will denote both (either) of them by the same symbol  $\mathcal{A}$ . Various pieces of the following theorem were discovered independently by Bar-Natan [5, 6], Kontsevich [25] and Lin [29]:

**Theorem 7.** (*Proof on page 16*)  $\mathcal{A}$  has a naturally defined product  $\cdot$  and a naturally defined co-product  $\Delta$ , which together make it a commutative and co-commutative Hopf algebra<sup>7</sup>. Furthermore, there is a naturally defined sequence of co-algebra automorphisms  $\{\psi^q\}_{q=-\infty}^{\infty}$  of  $\mathcal{A}$  reminiscent of the Adams operations of  $K$ -theory. These automorphisms satisfy  $\psi^q \circ \psi^p = \psi^{qp}$ .


<sup>7</sup>NOT in the graded (super) sense. Namely, elements in it honestly satisfy  $a \cdot b = b \cdot a$ , with no signs.

1.6. **The primitive elements of  $\mathcal{A}$ .** By the structure theory of Hopf algebras we know that  $\mathcal{A}$  is the symmetric algebra generated by the primitive elements of  $\mathcal{A}$ :

$$\mathcal{A} = \mathcal{S}(\mathcal{P}(\mathcal{A})), \quad \mathcal{P}(\mathcal{A}) = \{a \in \mathcal{A} : \Delta(a) = a \otimes 1 + 1 \otimes a\}. \quad (5)$$

The following description of  $\mathcal{A}$  is better suited for the study of  $\mathcal{P}(\mathcal{A})$ :

**Definition 1.11.** A *Chinese Character* (CC) is a (possibly empty) graph whose vertices are either trivalent and oriented (in the sense of definition 1.8), or are univalent. The trivalent vertices in such a graph are called *internal*, while the univalent vertices are called *external*. The collection of all Chinese characters *that have at least one external vertex in each connected component* will be denoted by  $\mathcal{C}$ . It is graded by half the number of vertices in a character.



**Figure 4.** A Chinese Character (CC) of degree 8. As in the case of CCDs, all trivalent vertices are oriented counterclockwise and all apparent vertices of valency higher than 3 are not vertices at all. This CC has four connected components.

Let  $\mathcal{B}$  be the quotient space

$$\mathcal{B} = \text{span}(\mathcal{C}) / \{\text{anti-symmetric vertices and IHX relations}\}. \quad (6)$$

Namely,  $\mathcal{B}$  is just the vector space generated by Chinese characters with exactly the two relations of theorem 6 imposed.  $\mathcal{B}$  inherits a grading from  $\mathcal{C}$ .

**Theorem 8.** (Kontsevich [25]) (Proof on page 29) *The spaces  $\mathcal{A}$  and  $\mathcal{B}$  are naturally isomorphic via maps  $\bar{\sigma} : \mathcal{A} \rightarrow \mathcal{B}$  and  $\bar{\chi} : \mathcal{B} \rightarrow \mathcal{A}$ . Furthermore, if a space  $\mathcal{P}$  is defined just like  $\mathcal{B}$  only with connected Chinese characters replacing arbitrary Chinese characters, then the above isomorphism identifies  $\mathcal{P}$  and  $\mathcal{P}(\mathcal{A})$ .*

The dual space<sup>8</sup>  $\mathcal{A}^*$  of the Hopf algebra  $\mathcal{A}$  is also a Hopf algebra, and thus one might wish to investigate the primitive elements of  $\mathcal{A}^*$ . Let  $\mathcal{P}'(\mathcal{A}^*)$  be the elements of degree different than 1 in  $\mathcal{P}(\mathcal{A}^*)$ .


**Theorem 9.** (1) (Proof on page 20)  $\mathcal{S}(\mathcal{P}'(\mathcal{A}^*))$  is isomorphic to the space  $\mathcal{W}$  of all weight systems.  
 (2) (Lin [29], Kontsevich [25], Bar-Natan) (Proof on page 34) *Via the correspondence of theorem 1, the weight systems in  $\mathcal{P}'(\mathcal{A}^*)$  correspond to additive Vassiliev invariants — Vassiliev invariants which are additive under the operation of taking the connected sum of two knots.*

<sup>8</sup> $\mathcal{A}^*$  is the graded dual of  $\mathcal{A}$  defined by  $\mathcal{A}^* = \bigoplus (\mathcal{G}_m \mathcal{A})^*$ .

1.7. **How big are  $\mathcal{A}$ ,  $\mathcal{W}$  and  $\mathcal{P}$ ?** One way to answer this question is by explicitly writing all diagrams and all relations, and using a computer to reduce the resulting matrix. The results are summarized in the table in page 34. Computer power is limited, however, and one might hope for better:

**Conjecture 1.** *All weight systems come from Lie algebras as in theorem 4. In other words, the weight systems produced in theorem 4 span the space of all weight systems.*

**Definition 1.12.** A *marked surface* is a compact two dimensional smooth surface with a choice of finitely many tangents (*markings*) to its boundary, regarded up to a diffeomorphism. (See figure 5). Let  $\mathcal{M}$  be the vector space spanned by the set of marked surfaces that have at least one marking on each connected component.



**Figure 5.** A marked surface with two boundary components

**Theorem 10.** (Kontsevich [25], Bar-Natan) (Proof on page 37) *There is a natural linear map  $\Phi : \mathcal{B} \rightarrow \mathcal{M}$ .*

**Theorem 11.** (Proof on page 43) *The pullback  $(\Phi \circ \bar{\sigma})^* \mathcal{M}^*$  of  $\mathcal{M}^*$  via  $\Phi \circ \bar{\sigma}$  is the subring of  $\mathcal{A}^*$  spanned by the linear functionals on  $\mathcal{A}$  generated (as in theorem 4) by Lie groups in the families  $SO$  and  $GL$  and all of their representations.  $(\Phi \circ \bar{\sigma})^* \mathcal{M}^*$  also contains in it the linear functionals on  $\mathcal{A}$  generated by all representations of Abelian, symplectic and Spin groups.*

*Remark 1.13.* The “philosophical father” of theorems 8, 10, and 11 is M. Kontsevich [25]. He suggested a somewhat weaker version of  $\Phi$ , whose image is the space spanned by normalized orientable (in the sense of definition 6.11) marked surfaces, and conjectured theorem 11 for his version of  $\Phi$  and only for the groups in the family  $GL$ . The stronger version of  $\Phi$ , the fact that our  $\Phi$  includes Kontsevich’s (lemma 6.49) and the proofs of theorems 11 and 12 are due to the author.

*Remark 1.14.* There is a natural extension (See section 2.3) of the theory of Vassiliev invariants to framed links. Framed links have various natural *cabling* operations (see definitions 3.13 and 6.21 and exercise 6.43) and, dually, invariants of framed links have various natural cabling operations. These operations take Vassiliev invariants to Vassiliev invariants of the same type (see exercises 3.14, 6.22 and 6.43).


**Theorem 12.** (Proof on page 48) *The ring of Vassiliev framed knot invariants coming from the writhe, the HOMFLY and the Kauffman polynomials and all of their cablings corresponds via theorem 1 to the pullback  $(\Phi \circ \bar{\sigma})^* \mathcal{M}^*$ .*

**Conjecture 2.** (See the discussion in page 48) Linear functionals on  $\mathcal{M}$  separate points in  $\mathcal{A}$ ; that is,  $\mathcal{A}^* = (\Phi \circ \bar{\sigma})^* \mathcal{M}^*$ . Theorem 11 implies that this conjecture is stronger than conjecture 1.


1.8. **Odds and ends.** The last section of the paper contains an incomplete survey of the literature available on Vassiliev invariants, one piece of bad news — that conjecture 1 and the completeness of Vassiliev invariants (problem 1.1) cannot be true at the same time, and some questions.

### 1.9. Summary of spaces and maps.

#### 1.9.1. Spaces.



#### 1.9.2. Dual spaces.



## 2. THE BASIC CONSTRUCTIONS

### 2.1. The classical knot polynomials.

2.1.1. *Proof of theorem 2.* Let  $C(K)(z)$  be the Conway polynomial of a knot  $K$ , and use the same symbol  $C$  to denote the natural extension (2) of the Conway polynomial to knots having self intersections. Then by the definition of the Conway polynomial [13, 21],

$$C \left( \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) = C \left( \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} \right) - C \left( \begin{array}{c} \diagdown \diagup \\ \diagdown \diagup \end{array} \right) = z \cdot C \left( \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} \right).$$

Therefore if  $K$  has more than  $m$  double points,  $C(K)$  is divisible by at least  $z^{m+1}$  and hence the coefficient of  $z^m$  in  $C(K)$  vanishes. This implies that the  $m$ th coefficient of the Conway polynomial is a Vassiliev invariant of type  $m$ .  $\square$

2.1.2. *Proof of theorem 3.* The idea is exactly the same as in the previous proof. In one of its standard parametrizations, the HOMFLY polynomial [19] is a function  $P$  of two parameters  $q$  and  $N$  which satisfies the identity

$$q^{N/2} P \left( \begin{array}{c} \diagup \\ \diagdown \end{array} \right) - q^{-N/2} P \left( \begin{array}{c} \diagdown \\ \diagup \end{array} \right) = (q^{1/2} - q^{-1/2}) P \left( \begin{array}{c} \diagup \\ \diagup \end{array} \right) \left( \begin{array}{c} \diagdown \\ \diagdown \end{array} \right).$$

Clearly if one changes variables to  $q = e^x$  and expands in powers of  $x$ , the above equation can be rewritten in the form

$$P \left( \begin{array}{c} \diagup \\ \diagdown \end{array} \right) - P \left( \begin{array}{c} \diagdown \\ \diagup \end{array} \right) = x \cdot (\text{some mess}).$$

The precise form of the ‘mess’ in the right side of the above equation is immaterial. What ever it is, the same argument as in the previous proof carries through and we see that (in this parametrization) the  $m$ th coefficient of the HOMFLY polynomial is a Vassiliev invariant of type  $m$ . The Jones polynomial is just the  $N \equiv 2$  specialization of the HOMFLY polynomial, and thus the same proof works for the Jones polynomial as well. Similar considerations also work in the case of the Kauffman polynomial.  $\square$

*Remark 2.1.* The Conway polynomial proof and idea to substitute  $q = e^x$  and expand in powers on  $x$  are due to the author. The argument in the HOMFLY case was completed by Birman and Lin in [10].

*Remark 2.2.* Similar arguments can be used to prove theorem 5. A complete proof can be found in Lin, [28] and Piunikhin, [33]. See also remark 4.8.

*Remark 2.3.* In fact, similar arguments can prove a slightly stronger theorem — that every knot invariant coming from a deformation of the identity solution of the Yang-Baxter equation can be re-expressed in terms of Vassiliev invariants.

**2.2. Constructing a weight system from a Vassiliev knot invariant.** Let  $V$  be a knot invariant of type  $m$ , and let  $D \in \mathcal{G}_m \mathcal{D}^c$  be a chord diagram of degree  $m$ . An *embedding* of  $D$  in  $\mathbf{R}^3$  will be an immersion  $K_D : S^1 \rightarrow \mathbf{R}^3$  of the circle into  $\mathbf{R}^3$  whose only singularities are transversal self intersections and which satisfies:

$$K_D(\theta) = K_D(\theta') \iff (\theta = \theta') \text{ or } \left( \begin{array}{l} \theta \text{ and } \theta' \text{ are the two} \\ \text{ends of a chord in } D \end{array} \right).$$

For example,

$$D = \textcircled{\times} \quad \longrightarrow \quad K_D = \text{[diagram of a knot with two crossings]}.$$


If  $K_D$  and  $\bar{K}_D$  are two embeddings of  $D$ , then one can get from one to the other by a sequence of ‘flips’, in which an overcrossing  $\begin{array}{c} \diagup \\ \diagdown \end{array}$  is replaced by an undercrossing  $\begin{array}{c} \diagdown \\ \diagup \end{array}$ . Each such flip *does not* change the value of  $V(K_D)$  — by definition 1.2 the change in the value of  $V(K_D)$  when such a flip is performed is given by  $V$  evaluated on a knot with  $m + 1$  double points, and this is 0 by the Birman-Lin condition of definition 1.3. Thus one can unambiguously set

$$W(D) = W_m(V)(D) = V(K_D).$$

*Proof of part (1) of theorem 1.* We need to show that  $W$  is a weight system. First consider the case where  $D$  has an isolated chord. Then  $K_D$  can be chosen to look like in figure 6. Using equation (2) on the point  $P$  of figure 6, we get


$$W(D) = V(K_D) = V(K_D^o) - V(K_D^u),$$

where  $K_D^o$  ( $K_D^u$ ) is the version of  $K_D$  in which the double point  $P$  was replaced by an overcrossing (undercrossing). But  $K_D^o$  and  $K_D^u$  are ambient isotopic, and therefore  $W(D) = 0$ .



**Figure 6.** The knot corresponding to a diagram having an isolated chord. The ends of the isolated chord are mapped to the point  $P$ .

Next let  $SW$  be the almost saturated (*i.e.* having  $m - 1$  self-intersections) knot shown (partially) in figure 7. Pieces of the  $x$  and  $y$  axes near the origin serve as arcs in that knot, as well as a third line  $z'$  parallel to the  $z$  axis but transversing the  $x - y$  plane South-West of the origin. Let  $NE$  be the same, only with the third line  $z'$  moved to transverse the  $x - y$  plane North-East of the origin. There are two ways to calculate  $V_i(NE)$  in terms of  $V_i(SW)$  and the weights of saturated knots (knots having precisely  $m$  double points) using the flip relation — by moving  $z'$  from  $SW$  to  $NE$  along the two dotted paths in figure 7. The two ways must yield the same answer, and therefore the four saturated knots corresponding to  $z'$  intersecting the  $x$  and  $y$  axes South, East, West and North of the origin have diagrams whose weights are related. With the sign convention of (2), this relation is seen to be (3).  $\square$



**Figure 7.** The graph  $SW$  and the two ways of getting from it to  $NE$ . Notice that  $z'$  is perpendicular to the plane and therefore appears as a single dot.

### 2.3. Framed links.

**Definition 2.4.** A framed knot is a knot  $K$  together with a choice of homotopy class in the space of never vanishing sections of the normal bundle of  $K$ . A framed link is a set of disjoint framed knots.

*Exercise 2.5.* Check that there is no difficulty in extending definitions 1.2 and 1.3 to framed knots. Show that the construction in the previous section carries through with only one change — if  $V$  is a Vassiliev invariant of framed knots and  $D$  has an isolated chord,  $W_m(V)(D)$  does not necessarily vanish.

*Exercise 2.6.* Show that there is a simple extension of definitions 1.2 and 1.3 and of the theory of the previous section to framed links. The main difference is that chord diagrams will now have many circles instead of just one.

**2.4. Constructing invariant tensors from Lie algebraic information.** In this section we will discuss a general method for constructing invariant tensors from a certain type of diagrams and some Lie algebraic information. When restricted to chord diagrams, this construction produces weight systems.

Let  $\mathfrak{g}$  be a finite dimensional Lie algebra over a field  $\mathbf{F}$ ,  $t$  an  $Ad$ -invariant symmetric non-degenerate bilinear form (a *metric*) on  $\mathfrak{g}$  (the Killing form of  $\mathfrak{g}$  is an example if  $\mathfrak{g}$  is semi-simple), and let  $R$  a finite dimensional representation of  $\mathfrak{g}$ . The objects  $(\mathfrak{g}, t, R)$  can all be regarded as tensors:

**The Lie algebra.** A Lie algebra  $\mathfrak{g}$  is a vector space (also denoted by  $\mathfrak{g}$ ) together with a distinguished element (tensor)  $\tilde{f}$  of  $\mathfrak{g}^* \otimes \mathfrak{g}^* \otimes \mathfrak{g}$  called *the Lie bracket*, subject to some well known requirements called *antisymmetry* and the *Jacobi identity*. The bilinear form  $t$  induces an isomorphism of  $\mathfrak{g}$  and  $\mathfrak{g}^*$ , and so  $\tilde{f}$  corresponds to some tensor  $f \in \mathfrak{g}^* \otimes \mathfrak{g}^* \otimes \mathfrak{g}^*$ . The tensor  $f$  is totally antisymmetric. We will represent it by a graph:

$$\begin{array}{c} \mathfrak{g}^* \text{---} \diagup \\ \text{---} \mathfrak{g}^* \text{---} \\ \mathfrak{g}^* \text{---} \diagdown \end{array} \longleftrightarrow f \in \mathfrak{g}^* \otimes \mathfrak{g}^* \otimes \mathfrak{g}^* .$$

(The internal vertex in the above graph should be oriented, otherwise it is not clear which order its external vertices should be taken with so that they correspond to the three  $\mathfrak{g}^*$ 's on the r.h.s. of the correspondence. As usual, if no orientation is specified, we pick the counterclockwise orientation.)

**The bilinear form.** The bilinear form  $t$  is a tensor in  $\mathfrak{g}^* \otimes \mathfrak{g}^*$ . Its inverse is a tensor  $t^{-1} \in \mathfrak{g} \otimes \mathfrak{g}$ . We will represent these two tensors by graphs:

$$\mathfrak{g}^* \text{---} \mathfrak{g}^* \leftrightarrow t \in \mathfrak{g}^* \otimes \mathfrak{g}^* \quad ; \quad \mathfrak{g} \text{---} \mathfrak{g} \leftrightarrow t^{-1} \in \mathfrak{g} \otimes \mathfrak{g} .$$

**The representation.** A representation  $R$  is a vector space (also denoted by  $R$ ) together with a distinguished tensor  $r$  in  $\mathfrak{g}^* \otimes R \otimes R^*$ . We will represent it by a graph:

$$\begin{array}{c} \mathfrak{g}^* \text{---} \uparrow \\ \text{---} R \\ \text{---} R^* \end{array} \longleftrightarrow r \in \mathfrak{g}^* \otimes R \otimes R^* .$$

For completeness, we will also have a graphical representation for the identity  $I \in \text{End}(R)$ :

$$R^* \longrightarrow R \quad \longleftrightarrow \quad I \in R^* \otimes R .$$

Let  $D$  be a diagram made of those components as above — dashed lines, directed full lines, oriented trivalent vertices in which three dashed lines meet, trivalent vertices in which a dashed line ends on a directed full line, ‘beginnings’ of full lines marked by an  $R^*$ , ‘ends’ of full lines marked by an  $R$ , and ‘ends’ of dashed lines marked by either a  $\mathfrak{g}$  or a  $\mathfrak{g}^*$ . To such a diagram we associate a tensor

$$\mathcal{T}(D) = \mathcal{T}_{\mathfrak{g},R}(D) \in \bigotimes_{\substack{\text{univalent vertices} \\ v \text{ of } D}} \left( \begin{array}{c} \text{the vector space} \\ \text{marked near } v \end{array} \right) .$$

The construction of  $\mathcal{T}(D)$  is simple. Simply separate  $D$  to a ‘union’ of its components, consider what you’ve got as a tensor in some higher tensor product of the spaces involved, and contract the obvious pairs of spaces and their duals.

*Example 2.7.* We get a tensor  $\Omega = \sum_{R^*} \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array} \in R \otimes R^* \otimes R \otimes R^*$  via

$$\begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array} \longrightarrow \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} \mathfrak{g}^* \quad \mathfrak{g} \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} \mathfrak{g} \quad \mathfrak{g}^* \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array},$$

surrounding by a box pairs of spaces that are to be contracted, we get

$$\longrightarrow \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} \boxed{\mathfrak{g}^* \mathfrak{g}} \quad \boxed{\mathfrak{g} \mathfrak{g}^*} \\ \uparrow \text{---} \\ R^* \end{array} \begin{array}{c} R \\ \uparrow \text{---} \\ R^* \end{array}.$$

In other words,  $\Omega$  is the image of  $r \otimes t^{-1} \otimes r \in (\mathfrak{g}^* \otimes R \otimes R^*) \otimes (\mathfrak{g} \otimes \mathfrak{g}) \otimes (\mathfrak{g}^* \otimes R \otimes R^*)$  under the map

$$(\mathfrak{g}^* \otimes R \otimes R^*) \otimes (\mathfrak{g} \otimes \mathfrak{g}) \otimes (\mathfrak{g}^* \otimes R \otimes R^*) \xrightarrow[\text{and spaces \#5 and \#6}]{\text{contract spaces \#1 and \#4}} R \otimes R^* \otimes R \otimes R^*.$$

*Example 2.8.* If we start from a *closed* diagram — a diagram with no external vertices, we just get a scalar in the ground field  $\mathbf{F}$ . It is an instructive exercise to verify that if  $\{\mathfrak{g}_a\}_{a=1}^{\dim \mathfrak{g}}$  is an orthonormal basis of  $\mathfrak{g}$  relative to the scalar product  $t$ , then

$$\mathcal{T}(\bigcirc) = \sum_{a,b,c=1}^{\dim \mathfrak{g}} f_{abc} \operatorname{tr}(R(\mathfrak{g}_a)R(\mathfrak{g}_b)R(\mathfrak{g}_c)) \in \mathbf{F},$$

where  $f_{abc}$  are the structure constants of  $\mathfrak{g}$  with respect to the basis  $\{\mathfrak{g}_a\}$ .

*Example 2.9.*  $\mathcal{T}(\bigcirc) = \operatorname{tr} I = \dim R$ .

The Lie algebra  $\mathfrak{g}$  acts on the spaces  $\mathfrak{g}$ ,  $\mathfrak{g}^*$ ,  $R$ ,  $R^*$ , and therefore also on their various tensor products. As  $f$ ,  $t^{-1}$ , and  $r$  are invariant tensors and the contraction operation is invariant, we see that:

**Proposition 2.10.** *The tensors  $\mathcal{T}(D)$  are invariant under the above mentioned  $\mathfrak{g}$ -action.*

The tensors  $\mathcal{T}(D)$  are not necessarily all different. Let  $\mathcal{D}^a$  ( $a$  for ‘all’) denote the collection of all diagrams made of the above mentioned components, and let  $\mathcal{A}^a$  be the vector space

$$\mathcal{A}^a = \operatorname{span}(\mathcal{D}^a) / \{\text{anti-symmetric vertices, } STU \text{ and } IHX \text{ relations}\}.$$

**Proposition 2.11.** *The map  $\mathcal{T} : \mathcal{D}^a \rightarrow \{\text{invariant tensors}\}$  defined by  $D \mapsto \mathcal{T}(D)$  descends to a map  $\mathcal{T} : \mathcal{A}^a \rightarrow \{\text{invariant tensors}\}$ .*

*Proof.* The Jacobi identity is the fact that a certain sum of three quadratic expressions in the structure constants of  $\mathfrak{g}$  vanishes. Each of the diagrams making the *IHX* relation (4) is such a quadratic expression, and it is easy to check that the *IHX* relation (in the context of tensors produced using a Lie algebra) is just a restatement of the Jacobi identity. Similarly, it can easily be seen that the *STU* relation is the fact that ‘representations represent’. Namely, it is just a restatement of the identity  $R([a, b]) = R(a)R(b) - R(b)R(a)$  for  $a, b \in \mathfrak{g}$ . Antisymmetry of internal vertices is the total antisymmetry of the tensor  $f$ .  $\square$

*Proof of part (1) of theorem 4.* Notice that a diagram  $D \in \mathcal{D}^t$  is closed, and so  $\mathcal{T}(D)$  is a scalar. Let  $W : \mathcal{D}^t \rightarrow \mathbf{F}$  denote the map  $D \mapsto \mathcal{T}(D)$ . We have just proven that  $W$  descends to a linear functional  $W \in (\mathcal{A}^t)^*$ . Using remark 1.10 we see that we can restrict this linear functional to  $\mathcal{G}_m \mathcal{A}^c$ . Call the resulting functional  $W_{\mathfrak{g}, R, m}$ .  $\square$


*Remark 2.12.* The discussion in this section has an obvious generalization to the case where each chain or cycle of directed arcs is associated with (is *colored* by) a (possibly) different representation of  $\mathfrak{g}$ . Furthermore, using linear extension cycles of directed arcs can be colored by an arbitrary *virtual* representation (and the resulting extension is consistent in the obvious sense). For what might be an even further generalization, see problem 5.4.

### 3. THE ALGEBRA $\mathcal{A}$ OF DIAGRAMS

In this section we will prove the theorems of section 1.5.

**3.1. Proof of theorem 6.** Let us start by proving that the linear map  $\phi^c : \mathcal{A}^c \rightarrow \mathcal{A}^t$  is an isomorphism. We will do that by constructing an inverse  $\phi^t : \mathcal{A}^t \rightarrow \mathcal{A}^c$  — a map  $\tilde{\phi}^t : \mathcal{D}^t \rightarrow \mathcal{A}^c$  satisfying the *STU* relation and extending the natural projection  $\mathcal{D}^c \rightarrow \mathcal{A}^c$ . To do that, notice that the *STU* relation expresses a diagram with some number  $k$  of internal vertices as a difference of diagrams with just  $k - 1$  internal vertices. Using the *STU* relation repeatedly it is clear how to construct  $\tilde{\phi}^t$  inductively, and the only problem is to show consistency — that if a CCD is reduced via *STU* to a linear combination of chord diagrams in two different ways, the resulting combinations are equivalent mod  $4T$ .

If  $D \in \mathcal{D}^t$  has only one internal vertex, consistency is clear — it is precisely the  $4T$  relation. So let us assume that  $\tilde{\phi}^t$  has been successfully defined on diagrams with less than  $k$  internal vertices,  $k > 1$ , and let  $D$  be a diagram with exactly  $k$  internal vertices. Suppose the *STU* relation is used to express  $D$  as a difference of two diagrams with  $k - 1$  internal vertices in two ways — by applying it to remove the arcs  $i$  or  $j$ , each of which connects the circle to an internal vertex. If  $i$  and  $j$  are not connected to the same internal vertex figure 8 shows why the two ways agree.



**Figure 8.** The consistency proof

If  $i$  and  $j$  are connected to the same internal vertex, pick a third arc  $l$  that connects the circle to a different internal vertex (if possible), and use the transitivity of the equality relation. There is one exceptional case<sup>9</sup> in which such a third arc  $l$  does not exist. It is somewhat more convenient to return to this case later, after the proof of lemma 3.1, on page 17.

<sup>9</sup>Of which I was informed by G. Masbaum after a preliminary version of this paper was circulated.

It remains to show that the  $STU$  relation implies the antisymmetry of internal vertices and the  $IHX$  identity. For the latter, by repeatedly using  $STU$  (if necessary), we can assume that the figures  $I$ ,  $H$ , and  $X$  that we are dealing with touch the circle in one of their corners, say the lower left corner. The proof is now similar to the proof that the commutator in an associative algebra satisfies the Jacobi identity, and is summarized in figure 9.

**Figure 9.**  $IHX$  follows from  $STU$

Similarly, the antisymmetry of internal vertices follows from:

**3.2. Proof of theorem 7.** For the purpose of this proof, it is convenient to ‘linearize’ diagrams:

More precisely, define  $\mathcal{D}^l$  to be the collection of all ‘linear’ diagrams — these are the same as Chinese character diagrams only with the circle replaced by a directed line. Then define  $\mathcal{A}^l$  to be

$$\text{span}(\mathcal{D}^l) / \text{span} \{ \text{all } STU \text{ relations} \} . \quad (8)$$


Clearly there is a map  $\phi^l : \mathcal{A}^l \rightarrow \mathcal{A}$  defined by ‘closing the line into a circle’ as in (7).

**Lemma 3.1.** *The map  $\phi^l$  is an isomorphism.*

*Proof.* The surjectivity of  $\phi^l$  is trivial, and all that is required is to prove its injectivity — that two ‘linear’ diagrams that map to the same ‘circular’ diagram are in fact equal as members of  $\mathcal{A}^l$ :


This is easily accomplished. Disconnect the vertex marked by the letter  $d$  in (9). Add three little ‘right turning hooks’ near each of the remaining vertices. Then put a  $(-)$  sign near

each of the hooks that is connected to a directed arc *leaving* the vertex. What you get looks like this:




Consider the signed sum  $\Sigma$  of all possible ways of connecting the ‘floating end’ (marked by the letter  $f$ ) to one of the 15 (in this case) hooks, taking hooks marked with a  $(-)$  with a negative sign. There are two ways to group the 15 terms in this sum, and comparing these two ways will prove our lemma.

- (1) By arcs: our sign convention and the antisymmetry of internal vertices show that when the terms in  $\Sigma$  are grouped by arcs, all drops out (see figure 10) except for the terms corresponding to the three ‘groupless’ hooks. These are marked by 1, 2, and 3 in (10). The first two of these three terms form exactly the left hand side of (9), while the third vanishes because of the antisymmetry of internal vertices.



**Figure 10.** Grouping by arcs

- (2) By vertices: our sign convention, the antisymmetry of internal vertices, and the  $STU$  and  $IHX$  relations show that when the terms in  $\Sigma$  are grouped by vertices, all drops out (see figure 11).  $\square$



**Figure 11.** Grouping by vertices

*Remark 3.2.* From the knot theory view point, lemma 3.1 is just the fact that the theory of knots with a distinguished point is equivalent to the standard theory of knots. In [10], Birman and Lin have used this view point to prove a special case of lemma 3.1.

**Conclusion of the proof of theorem 6:** The one exceptional case alluded to in the proof of theorem 6 is when the arcs  $i$  and  $j$  are connected to the same internal vertex and

there are no other arcs connecting an internal vertex to the circle. In that case,  $D$  must look as follows:

$$D = \left( \text{diagram of a circle with a blob } B \text{ and vertices } i, j \right) \cdot \left( \begin{array}{l} \text{no trivalent vertices other than inside the} \\ \text{blob } B, \text{ and the one connecting } i \text{ and } j. \end{array} \right).$$

Fortunately, no matter how  $STU$  is used to reduce  $D$  to chord diagrams, the answer is 0 mod  $4T$ . Indeed, applying  $STU$  around (say)  $i$  and then collapsing  $B$  into chord diagrams using  $STU$  relations, we get

$$D = \left( \text{diagram 1} \right) - \left( \text{diagram 2} \right) = \left( \text{diagram 3} \right) - \left( \text{diagram 4} \right) = D_T - D_U.$$

But  $D_T$  and  $D_U$  are equal — to get from  $D_T$  to  $D_U$  one has to pass the end  $f$  of the chord marked  $n$  over the chords in the blob  $B'$ . This is possible (modulo the  $4T$  relation) by an argument parallel to the argument in the proof of lemma 3.1. The only differences are that hooks are put only on the circle, and that instead of summing  $STU$  and  $IHX$  relations over vertices, one sums  $4T$  relations over dashed arcs.  $\square$

**Definition 3.3.** The product  $\cdot : \mathcal{D}^l \otimes \mathcal{D}^l \rightarrow \mathcal{D}^l$  is the operation of *connected sum*,

$$\left( \text{diagram 1} \right) \cdot \left( \text{diagram 2} \right) = \left( \text{diagram 3} \right).$$

**Proposition 3.4.** *The above defined product  $\cdot$  descends to a product  $\cdot : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ . With this product,  $\mathcal{A}$  is an associative and commutative algebra. The empty circle  $\bigcirc$  is a unit in  $\mathcal{A}$ .*

*Proof.* Remembering that  $\mathcal{A} \simeq \mathcal{A}^l$ , the only problem that remains is to show the commutativity of the product as an operation  $\mathcal{A}^l \otimes \mathcal{A}^l \rightarrow \mathcal{A}^l$ . But clearly for  $D_{1,2} \in \mathcal{D}^l$ ,  $D_1 \cdot D_2$  and  $D_2 \cdot D_1$  correspond to the same diagram in  $\mathcal{D}^l$  under the isomorphism  $\phi^l$ .  $\square$

*Remark 3.5.* As an operation on  $\mathcal{D}^t$ , the product is also given as connected sum,

$$\left( \text{diagram 1} \right) \cdot \left( \text{diagram 2} \right) = \left( \text{diagram 3} \right) = \left( \text{diagram 4} \right).$$


It is *not* well defined as an operation  $\mathcal{D}^t \otimes \mathcal{D}^t \rightarrow \mathcal{D}^t$ . As such, it depends on the points on circles used for the surgery involved in the connected sum. However, as an operation  $\mathcal{D}^t \otimes \mathcal{D}^t \rightarrow \mathcal{A}$  it is well defined, as follows from the equivalence of  $\mathcal{A}^l$  and  $\mathcal{A}^t$ .

*Exercise 3.6.* Check that the product of  $\mathcal{A}$  corresponds in a natural way to the *connected sum* operation on knots.

**Definition 3.7.** Let  $\mathcal{D}$  be the vector space generated by the elements of  $\mathcal{D}^t$ . Define a *splitting*  $D_s$  of a diagram  $D \in \mathcal{D}^t$  to be a marking of the dashed arcs in  $D$  by the letters  $l$  and  $r$ , so that if three dashed arcs of  $D$  meet at some vertex, then they are all marked by the same letter. Define  $\Delta : \mathcal{D}^t \rightarrow \mathcal{D} \otimes \mathcal{D}$  by

$$\Delta(D) = \sum_{\substack{\text{all splittings} \\ D_s \text{ of } D}} L(D_s) \otimes R(D_s),$$

where  $L(D_s)$  ( $R(D_s)$ ) is obtained from  $D$  by removing all the arcs marked by  $r$  ( $l$ ) in  $D_s$ . See figure 12.



**Figure 12.** Computing  $\Delta(\text{trivalent vertex})$ .

**Proposition 3.8.**  $\Delta$  descends to a co-commutative co-associative co-product on  $\mathcal{A}$ . The linear functional  $\epsilon \in \mathcal{A}^*$  defined by  $\epsilon(\text{circle}) = 1$ ;  $\epsilon|_{\mathcal{G}_m \mathcal{A}} = 0$ , ( $m > 0$ ) is a co-unit in the resulting co-algebra.

*Proof.* An explicit computation shows that

$$\Delta(\text{trivalent vertex} - \text{trivalent vertex} + \text{trivalent vertex}) = (\text{trivalent vertex} - \text{trivalent vertex} + \text{trivalent vertex}) \otimes (\text{circle}) + (\text{circle}) \otimes (\text{trivalent vertex} - \text{trivalent vertex} + \text{trivalent vertex}).$$

This implies that  $\Delta$  is well defined as a map  $\Delta : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ . The other assertions of the proposition are trivial.  $\square$

**Proposition 3.9.** With the operations defined above,  $\mathcal{A}$  is a Hopf algebra.

*Proof.* Let  $D_{1,2} \in \mathcal{D}^t$ . The compatibility  $\Delta(D_1 D_2) = \Delta(D_1) \Delta(D_2)$  of the product and the co-product is just the statement

$$\sum_{\substack{\text{all splittings} \\ (D_1 D_2)_s \text{ of } D_1 D_2}} L((D_1 D_2)_s) \otimes R((D_1 D_2)_s) = \sum_{\substack{\text{all splittings} \\ D_{1s} \text{ of } D_1 \\ D_{2s} \text{ of } D_2}} L(D_{1s}) L(D_{2s}) \otimes R(D_{1s}) R(D_{2s}),$$

which is rather clear.  $\square$

*Exercise 3.10.* Show that the co-product of  $\mathcal{A}$  corresponds to the multiplication of knot invariants. In other words, let  $V_1$  ( $V_2$ ) be a Vassiliev invariant of type  $m_1$  ( $m_2$ ). Show that  $V_1 \cdot V_2$  is a Vassiliev invariant of type  $m_1 + m_2$  and that

$$W_{m_1+m_2}(V_1 \cdot V_2) = (W_{m_1}(V_1) \otimes W_{m_2}(V_2)) \circ \Delta.$$

**Definition 3.11.** Let  $D \in \mathcal{D}^t$  be a diagram with exactly  $n$  internal trivalent vertices, and let  $q \in \mathbf{Z}$  be a non-zero integer. Define  $\psi^q(D) \in \mathcal{D}$  to be the sum of all possible ways of lifting  $D$  to the  $q$ th cover of the circle. For example,

$$\psi^2(\text{circle with two vertices}) = \left( \left( \text{circle with two vertices} \right) + \left( \text{circle with two vertices} \right) + \dots \right) = \text{circle with two vertices} + \text{circle with two vertices} + \dots = 12 \text{circle with two vertices} + 4 \text{circle with two vertices}.$$

For  $q = 0$ , just define

$$\psi^0(D) = \begin{cases} \text{circle} & \text{if } D = \text{circle}, \\ 0 & \text{otherwise.} \end{cases}$$

*Exercise 3.12.* Show that  $\psi^q$  descends to a co-algebra automorphism of  $\mathcal{A}$  and that for any  $q, p \in \mathbf{Z}$ ,  $\psi^q \circ \psi^p = \psi^{qp}$ .

Notice that propositions 3.4, 3.8, 3.9 and exercise 3.12 prove theorem 7.

**Definition 3.13.** Let  $K : (S^1 = \{z : |z| = 1\}) \rightarrow \mathbf{R}^3$  be a framed knot and let  $n(z)$  ( $z \in S^1$ ) be a section of the normal bundle of  $K$  compatible with its framing. Notice that the normal bundle of  $K$  has a natural complex structure. For a non-zero integer  $q$  define the  $q$ th *connected cabling*  $\psi^q(K)$  of  $K$  to be the knot given by

$$\psi^q(K)(z) = K(z^q) + \epsilon zn(z^q),$$

where  $\epsilon$  is a very small number. Let  $\psi^q(K)$  inherit the framing of  $K$ .

*Exercise 3.14.* Show that the operations  $\psi^q$  on knots and on diagrams correspond. In other words, show that if  $V$  is a Vassiliev invariant of type  $m$  of framed knots then  $V \circ \psi^q$  is also a Vassiliev invariant of type  $m$ , and that  $W_m(V \circ \psi^q) = W_m(V) \circ \psi^q$ .

3.2.1. *Proof of part (1) of theorem 9.*

*Exercise 3.15.* Prove that  $\mathcal{W}$  is a sub-Hopf-algebra of  $\mathcal{A}^*$  and that  $\mathcal{P}(\mathcal{W}) = \mathcal{P}(\mathcal{A}^*) \cap \mathcal{W}$ .

The exercise implies that to prove part (1) of theorem 9 it is enough to show that

$$\mathcal{P}(\mathcal{A}^*) \cap \mathcal{W} = \mathcal{P}'(\mathcal{A}^*).$$

A functional  $W \in \mathcal{A}^*$  is primitive iff  $\Delta(W) = \epsilon \otimes W + W \otimes \epsilon$  iff for all  $D_1$  and  $D_2$ ,  $W(D_1 \cdot D_2) = \epsilon(D_1)W(D_2) + W(D_1)\epsilon(D_2)$ . The last condition means that  $W$  is primitive iff it vanishes on reducible diagrams (those diagrams which are a product in a non-trivial way). Now if  $W \in \mathcal{P}'(\mathcal{A}^*)$  then  $W$  is primitive and  $\deg W > 1$ . This implies that  $W$  vanishes on diagrams having an isolated chord. On the other hand, if  $W \in \mathcal{P}(\mathcal{A}^*) \cap \mathcal{W}$ , then it vanishes on  $\textcircled{\text{---}}$  and therefore it is of degree higher than 1.  $\square$

3.2.2. *Proof of part (2) of theorem 4.* All that that remains to be shown is that there exists a canonical projection

$$(W \rightarrow \hat{W}) : (\mathcal{A}^* = \mathcal{S}(\mathcal{P}(\mathcal{A}^*))) \rightarrow (\mathcal{W} = \mathcal{S}(\mathcal{P}'(\mathcal{A}^*))).$$

Such a projection is induced from the natural projection  $\mathcal{P}(\mathcal{A}^*) \rightarrow \mathcal{P}'(\mathcal{A}^*)$  that maps the primitive of degree 1 to 0 and acts as the identity otherwise.  $\square$

*Exercise 3.16.* Define a map  $\phi : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$  by

$$\phi(D_1 \otimes D_2) = \begin{cases} \left( \textcircled{\text{---}} \right)^{\deg D_1} \cdot D_2 & \text{if } D_1 \text{ is chord diagram,} \\ 0 & \text{if } D_1 \text{ has internal trivalent vertices,} \end{cases}$$

and show that  $\hat{W}(D) = W(\phi(\Delta D))$ .<sup>10</sup>

---

<sup>10</sup>I wish to thank S. Willerton for noticing an inaccuracy in my original formulation of this exercise and for supplying this much cleaner version.

#### 4. KONTSEVICH'S KNIZHNIK-ZAMOLODCHIKOV CONSTRUCTION

In this section we will mostly be concerned with proving the hard part of theorem 1 — with defining the map  $V$ . Recall that the ‘ $m$ th derivatives’ of a Vassiliev invariant of type  $m$  is a system of ‘constants’, and that we called such systems of constants *weight systems*. The aim now is to show that every weight system can be ‘integrated’  $m$  times to give a Vassiliev invariant. An incomplete proof of that fact first appeared in Bar-Natan [6], where it was shown that the combinatorics underlying perturbative Chern-Simons theory is essentially that of weight systems, and that on a naive level, ignoring questions of convergence, perturbative Chern-Simons theory can be used to integrate a weight system to a Vassiliev invariant. The analytical difficulties in [6] were later resolved by Axelrod and Singer [3] and by Kontsevich [26]. Later on, Kontsevich found a second proof based on the Knizhnik-Zamolodchikov equation [23], in which convergence is much easier to show. This is the proof that we will present here.

**4.1. Connections, curvature, and holonomy.** Up to some (important) subtlety, a connection is a 1-form whose values are in the algebra of endomorphisms of the fiber. One would like to know how much of the theory of connections can be generalized to the case of 1-forms with values in an arbitrary associative algebra. As was shown by K-T. Chen [12], much of the theory persists in the more general case. Let us briefly review some aspects of Chen’s theory.

Let  $X$  be a smooth manifold and let  $\mathfrak{A}$  be a topological algebra over the real numbers  $\mathbf{R}$  (or the complex numbers  $\mathbf{C}$ ), with a unit 1. An  $\mathfrak{A}$ -valued connection  $\Omega$  on  $X$  is an  $\mathfrak{A}$ -valued 1-form  $\Omega$  on  $X$ . Its curvature  $F_\Omega$  is the  $\mathfrak{A}$ -valued 2-form  $F_\Omega = d\Omega + \Omega \wedge \Omega$ , where the definitions of the exterior differentiation operator  $d$  and of the wedge product  $\wedge$  are precisely the same as the corresponding definitions in the case of matrix valued forms. The notion of “parallel transport” also has a generalization in the new context: Let  $B : I \rightarrow X$  be a smooth map from some interval  $I = [a, b]$  to  $X$ . Define the *holonomy*  $h_{B,\Omega}$  of  $\Omega$  along  $B$  to be the function  $h_{B,\Omega} : I \rightarrow \mathfrak{A}$  which satisfies

$$h_{B,\Omega}(a) = 1; \quad \frac{\partial}{\partial t} h_{B,\Omega}(t) = \Omega(\dot{B}(t)) h_{B,\Omega}(t), \quad (t \in I)$$

if such a function exists and is unique. In many interesting cases,  $h_{B,\Omega}$  exists and is given (see *e.g.* [12]) by the following “iterated integral” formula:

$$h_{B,\Omega}(t) = 1 + \sum_{m=1}^{\infty} \int_{a \leq t_1 \leq \dots \leq t_m \leq t} (B^*\Omega)(t_m) \cdot \dots \cdot (B^*\Omega)(t_1). \quad (11)$$

(In this formula  $B^*\Omega$  denotes the pullback of  $\Omega$  to  $I$  via  $B$ ). Furthermore, just like in the standard theory of connections, if  $F_\Omega \equiv 0$  (‘ $\Omega$  is flat’), then  $h_{B,\Omega}$  is invariant under homotopies of  $B$  that preserve its endpoints.

In the case of interest for us,  $\mathfrak{A}$  will be the completion of a graded algebra of finite type over the complex numbers — the direct product of the finite dimensional (over  $\mathbf{C}$ ) homogeneous components of a graded algebra. The connection  $\Omega$  will be homogeneous of degree 1. In this case the  $m$ th term  $h_{m,B,\Omega}$  in (11) is homogeneous of degree  $m$ , and there is no problem with the convergence of the sum there. Also, as each term lives in a different degree, Chen’s theory implies that each term is invariant under homotopies of  $B$  that preserve its endpoints.

These assertions are not very hard to verify directly from the definition of  $h_{m,B,\Omega}$  as a multiple integral.

**4.2. The formal Knizhnik-Zamolodchikov connection.** Let  $\mathcal{D}_n^{KZ}$  be the collection of all diagrams made of  $n$  ordered upward pointing arrows, and dashed arcs and oriented vertices as in the definition of  $\mathcal{A}^t$ , with the standard conventions about higher than trivalent vertices and about the orientation of vertices:

$$\mathcal{D}_n^{KZ} = \left\{ \underbrace{\left( \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \cdots \quad \uparrow \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \uparrow \quad \uparrow \quad \uparrow \quad \cdots \quad \uparrow \end{array} \right)}_{n \text{ ordered upward pointing arrows}} \right\}.$$

Let the ground field be  $\mathbf{C}$  and let  $\mathcal{A}_n^{KZ}$  be the quotient

$$\mathcal{A}_n^{KZ} = \text{span}(\mathcal{D}_n^{KZ}) / \{STU \text{ relations}\}.$$

$\mathcal{A}_n^{KZ}$  is an algebra with ‘composition’ as its product:

$$\begin{array}{c} \uparrow \\ \text{---} \\ \uparrow \end{array} \cdot \begin{array}{c} \uparrow \quad \uparrow \\ \text{---} \quad \text{---} \\ \uparrow \quad \uparrow \end{array} = \begin{array}{c} \uparrow \quad \uparrow \\ \text{---} \quad \text{---} \\ \uparrow \quad \uparrow \end{array}, \quad (n=3).$$

$\mathcal{A}_n^{KZ}$  is graded by half the number of vertices in a diagram, excluding the  $2n$  endpoints of the  $n$  arrows; the degree of the above product is 4.

For  $1 \leq i < j \leq n$  define  $\Omega_{ij} \in \mathcal{A}_n^{KZ}$  by

$$\Omega_{ij} = \begin{array}{c} \uparrow \quad \cdots \quad \uparrow^i \quad \text{---} \quad \uparrow^j \quad \cdots \quad \uparrow \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \end{array}.$$

Let  $X_n$  be the configuration space of  $n$  distinct points in  $\mathbf{C}$ ;  $X_n = \{(z_1, \dots, z_n) \in \mathbf{C}^n : z_i = z_j \Rightarrow i = j\}$ , and let  $\omega_{ij}$  be the complex 1-form on  $X_n$  defined by

$$\omega_{ij} = d(\log z_i - z_j) = \frac{dz_i - dz_j}{z_i - z_j}.$$

The formal Knizhnik-Zamolodchikov connection is the  $\mathcal{A}_n^{KZ}$ -valued connection

$$\Omega_n = \sum_{1 \leq i < j \leq n} \Omega_{ij} \omega_{ij}$$

on  $X_n$ .

**Proposition 4.1.** *The formal Knizhnik-Zamolodchikov connection  $\Omega_n$  is flat.*

*Proof.* Clearly  $d\Omega_n = 0$ . Let us check that

$$\Omega_n \wedge \Omega_n = \sum_{i < j; i' < j'} \Omega_{ij} \Omega_{i'j'} \omega_{ij} \wedge \omega_{i'j'} = 0. \quad (12)$$

The above sum can be separated into three parts, according to the cardinality of the set  $\{i, j, i', j'\}$ . If this cardinality is 2 or 4 then  $\Omega_{ij}$  and  $\Omega_{i'j'}$  commute, while  $\omega_{ij}$  and  $\omega_{i'j'}$  anti-commute. It is easy to check that this implies that the corresponding parts of the sum (12)

vanish. The only interesting case is when  $|\{i, j, i', j'\}| = 3$ , say  $\{i, j, i', j'\} = \{1, 2, 3\}$ . In this case,

$$\sum_{\{i, j, i', j'\}=\{1, 2, 3\}} \Omega_{ij} \Omega_{i'j'} \omega_{ij} \wedge \omega_{i'j'} = (\Omega_{12} \Omega_{23} - \Omega_{23} \Omega_{12}) \omega_{12} \wedge \omega_{23} + (\text{cyclic permutations}).$$

By the *STU* relation this is

$$= \Omega_{123} (\omega_{12} \wedge \omega_{23} + (\text{cyclic permutations})) = 0, \quad (13)$$

where  $\Omega_{123}$  is given by

$$\Omega_{123} = \begin{array}{c} \uparrow \\ \uparrow \text{---} \uparrow \\ \uparrow \end{array} \cdot \cdot \cdot \begin{array}{c} \uparrow \\ \uparrow \\ \uparrow \end{array} \in \mathcal{A}_n^{KZ}.$$

The vanishing of  $\omega_{12} \wedge \omega_{23} + (\text{cyclic permutations})$  is called ‘Arnold’s identity’ [1] and can be easily verified by a direct computation.  $\square$

*Remark 4.2.* The connection  $\Omega_n$  has a simple generalization to the case when the underlying algebra is  $\mathcal{A}_{n,n}^{KZ}$ , the algebra generated by diagrams having  $2n$  arrows, whose first  $n$  arrows point upward and whose next  $n$  arrows point downward. The only difference is a sign difference in the application of the *STU* relation in (13). Therefore if one defines

$$\Omega_{n,n} = \sum_{1 \leq i \leq j \leq 2n} s_i s_j \Omega_{ij} \omega_{ij},$$

where  $s_i = \begin{cases} +1 & i \leq n \\ -1 & i > n \end{cases}$ , then the connection  $\Omega_{n,n}$  is flat.


**4.3. Kontsevich’s integral invariants.** Choose a decomposition  $\mathbf{R}^3 = \mathbf{C}_z \times \mathbf{R}_t$  of  $\mathbf{R}^3$  to a product of a complex plane  $\mathbf{C}_z$  parametrized by  $z$  and a real line  $\mathbf{R}_t$  parametrized by  $t$  and let  $K : S^1 \rightarrow \mathbf{R}^3$  be a parametrized knot on which the function  $t$  is a Morse function. Consider the following series, whose precise definition will be discussed below:

$$Z(K) = \sum_{m=0}^{\infty} \frac{1}{(2\pi i)^m} \int_{t_{\min} < t_1 < \dots < t_m < t_{\max}} \sum_{\substack{\text{applicable pairings} \\ P=\{(z_i, z'_i)\}}} (-1)^{\#P_{\downarrow}} D_P \bigwedge_{i=1}^m \frac{dz_i - dz'_i}{z_i - z'_i} \in \mathcal{A}_{\mathbf{C}}^r. \quad (14)$$

In the above equation,



- $t_{\min}$  ( $t_{\max}$ ) is the minimal (maximal) value of  $t$  on  $K$ .
- an ‘applicable pairing’ is a choice of an unordered pair  $(z_i, z'_i)$  for every  $1 \leq i \leq m$ , for which  $(z_i, t_i)$  and  $(z'_i, t_i)$  are *distinct* points on  $K$ .
- $\#P_{\downarrow}$  is the number of points of the form  $(z_i, t_i)$  or  $(z'_i, t_i)$  at which  $K$  is decreasing. Remember that in this article we are only considering *oriented* knots.
- $\mathcal{A}_{\mathbf{C}}^r$  is the quotient of  $\mathcal{A}_{\mathbf{C}}$  by the ideal generated by the diagram  $\bigcirc$ . In other words, diagrams having an isolated chord (in the sense of definition 1.6) are set equal to 0 in  $\mathcal{A}_{\mathbf{C}}^r$ . The subscripts  $\mathbf{C}$  are intended to remind us that the construction is done over the ground field  $\mathbf{C}$ . There is a similar definition for  $\mathcal{A}_{\mathbf{R}}^r$ .
- $D_P$  is the chord diagram naturally associated with  $K$  and  $P$  as in figure 13. It is to be regarded as an element of the quotient  $\mathcal{A}_{\mathbf{C}}^r$ .

- Every pairing defines a map  $\{t_i\} \mapsto \{(z_i, z'_i)\}$  locally around the current values of the  $t_i$ 's. Use this map to pull the  $dz_i$ 's and  $dz'_i$ 's to the  $m$ -simplex  $t_{\min} < t_1 < \dots < t_m < t_{\max}$  and then integrate the indicated wedge product over that simplex.



**Figure 13.**  $m = 4$ : a knot  $K$  with a pairing  $P$  and the corresponding chord diagram  $D_P$ . Notice that  $D_P = 0$  in  $\mathcal{A}_{\mathbb{C}}^r$  due to the isolated chord marked by 1.


4.3.1. *Finiteness.* Properly interpreted, the integrals in (14) are finite. There appears to be a problem in the denominator when  $z_i - z'_i$  is small for some  $i$ . This can happen in either of two ways:

- (1)  in this case the integration domain for  $z_{i+1}$  is as small as  $z_i - z'_i$ , and its 'smallness' cancels the singularity coming from the denominator.
- (2)  in this case the corresponding diagram  $D_P$  has an isolated chord, and so it is 0 in  $\mathcal{A}_{\mathbb{C}}^r$ .

4.3.2. *Invariance under horizontal deformations.* For times  $t_{\min} \leq a < b \leq t_{\max}$  define  $Z(K, [a, b])$  in exactly the same way as (14), only restricting the domain of integration to be  $a < t_1 < \dots < t_m < b$ . Of course,  $Z(K, [a, b])$  will not be in  $\mathcal{A}_{\mathbb{C}}^r$ , but rather in the vector space

$$\mathcal{A}_{\mathbb{C}}^{K, [a, b]} = \text{span} \left\{ \begin{array}{l} \text{diagrams whose solid lines} \\ \text{are as in the part of } K \text{ on} \\ \text{which } a \leq t \leq b \end{array} \right\} / \text{span} \left\{ \begin{array}{l} STU \text{ relations and di-} \\ \text{agrams with subdia-} \\ \text{grams like } \begin{array}{c} \text{---} \circ \text{---} \\ \text{---} \end{array} \end{array} \right\}.$$

For example, if  $t_1, t_4$ , and  $K$  are as in figure 13, then the following is a diagram in  $\mathcal{A}_{\mathbb{C}}^{K, [t_1, t_4]}$ .




The same reasoning as in section 4.3.1 shows that  $Z(K, [a, b])$  is finite. For  $t_{\min} \leq a < b < c \leq t_{\max}$ , there is an obvious product  $\mathcal{A}_{\mathbb{C}}^{K, [a, b]} \otimes \mathcal{A}_{\mathbb{C}}^{K, [b, c]} \rightarrow \mathcal{A}_{\mathbb{C}}^{K, [a, c]}$ , and it is easy to show that with this product  $Z(K, [a, b])Z(K, [b, c]) = Z(K, [a, c])$ .

Let  $t_{\min} < a < b < t_{\max}$  be times for which  $K$  has no critical points in the time slice  $a \leq t \leq b$ , and let  $n$  be the number of upward (or downward) pointing strands of  $K$  in that slice. Then  $\mathcal{A}_{\mathbb{C}}^{K, [a, b]} \equiv \mathcal{A}_{n, n}^{KZ}$ , and comparing with (11) and the definition of  $\Omega_{n, n}$  we see that  $Z(K, [a, b])$  is the holonomy of  $\Omega_{n, n}$  along the braid defined by the intersection of  $K$  with the slice  $a \leq t \leq b$ . The flatness of  $\Omega_{n, n}$  implies that this holonomy is invariant under horizontal deformations of that piece of  $K$ , and together with

$$Z(K) = Z(K, [t_{\min}, t_{\max}]) = Z(K, [t_{\min}, a])Z(K, [a, b])Z(K, [b, t_{\max}])$$

we see that  $Z(K)$  is invariant under horizontal deformations of  $K$  which ‘freeze’ the time slices in which  $K$  has a critical point.

**4.3.3. Moving critical points.** In this section we will show that (subject to some restrictions)  $Z(K)$  is also invariant under deformations of  $K$  that do move critical points. The idea is to narrow critical points to sharp needles using horizontal deformations, and then show that very sharp needles contribute almost nothing to  $Z(K)$  and therefore can be moved around (almost) freely:




**Lemma 4.3.** *If the two knots  $K_{1,2}$  both contain a sharp needle of width  $\epsilon$ , and are the identical except possibly for the length and the directions of their respective needles, then*

$$\|Z_m(K_1) - Z_m(K_2)\| \sim \epsilon$$

where  $Z_m$  is the degree  $m$  piece of  $Z$  and  $\|\cdot\|$  is some fixed norm on  $\mathcal{G}_m \mathcal{A}_{\mathbb{C}}^r$ .

*Proof.* Clearly, the difference between  $Z_m(K_1)$  and  $Z_m(K_2)$  will come only from terms in (14) in which one of the  $z_i$ 's (or  $z'_i$ 's) is on the needle. So let us show that if a knot  $K$  contains a needle  $N$  of width  $\epsilon$ , then such terms in  $Z_m(K)$  are at most proportional to  $\epsilon$ . Without loss of generality we can assume that the needle  $N$  points upward. If the highest pair  $(z_i, z'_i)$  that touches  $N$  connects the two sides of  $N$ , the corresponding diagram is 0 in  $\mathcal{A}_{\mathbb{C}}^r$  and there is nothing to worry about. If there is no pair  $(z_j, z'_j)$  that connects the two sides of  $N$  then again life is simple: in that case there are no singularities in (14) so nothing big prevents



from being small. (Notice that these two terms appear in  $Z(K)$  with opposite signs due to the factor  $(-1)^{\#P_1}$  but otherwise they differ only by something proportional to  $\epsilon$ ). If  $(z_j, z'_j)$  is a pair that does connect the two sides of  $N$ , it has to do so in the top (round) part of  $N$  — otherwise  $dz_j - dz'_j = 0$ .

So the only terms that cause some worry are those that have some  $k > 1$  pairs  $(z_{j_1}, z'_{j_1})$  through  $(z_{j_k}, z'_{j_k})$  on the top part of  $N$ , with  $(z_{j_k}, z'_{j_k})$  being the highest of these pairs and  $(z_{j_1}, z'_{j_1})$  the lowest. We might as well assume that there are no pairs other than  $(z_i, z'_i)$  that touch  $N$  only once — such pairs just shorten the domain of integration in (14) without adding any singularity in the denominator. So what we have looks like:



Writing  $\delta_\alpha = |z_{j_\alpha} - z'_{j_\alpha}|$ , we see that the integral corresponding to (16) is bounded by a constant times


$$\int_0^\epsilon \frac{d\delta_1}{\delta_1} \int_0^{\delta_1} \frac{d\delta_2}{\delta_2} \dots \int_0^{\delta_{k-1}} \frac{d\delta_k}{\delta_k} \int_{z_{j_k}}^{z'_{j_k}} \frac{dz_i - dz'_i}{z_i - z'_i} \sim \epsilon. \quad \square$$

Unfortunately, there is one type of deformation that (15) and lemma 4.3 cannot handle — the total number of critical points in  $K$  cannot be changed:



Even if the hump on the left figure is deformed into a needle and then this needle is removed, a (smaller) hump still remains.

4.3.4. *The correction.* Let the symbol  $\infty$  stand for the embedding



Notice that

$$Z(\infty) = \bigcirc + (\text{higher order terms}) \quad (18)$$

and so using power series  $Z(\infty)$  can be inverted and the following definition makes sense:

**Definition 4.4.** Let  $K$  be a knot embedded in  $\mathbf{C} \times \mathbf{R}$  with  $c$  critical points. Notice that  $c$  is always even and set<sup>11</sup>

$$\tilde{Z}(K) = \frac{Z(K)}{(Z(\infty))^{\frac{c}{2}-1}}.$$


**Theorem 13.**  $\tilde{Z}(K)$  is invariant under arbitrary deformations of the knot  $K$ .

<sup>11</sup>The non-invariance of  $Z(K)$  under the move (17) was first noticed by R. Bott and the author. The correction  $\tilde{Z}(K)$  is due to Kontsevich [25].

*Proof.* Clearly,  $\tilde{Z}(K)$  is invariant under deformations that do not change the number of critical points of  $K$ , and the only thing that remains to be checked is its invariance under the move (17). So let  $K_c$  and  $K_s$  be two knots that are identical other than that in some place  $K_c$  has the figure in the left side of (17) while in the same place  $K_s$  has the figure on the right side of (17). We need to show that in  $\mathcal{A}_{\mathbb{C}}^r$ ,

$$Z(K_c) = Z(\infty)Z(K_s).$$

Using deformations as in section 4.3.3 we can move the ‘humps’ of  $K_c$  to be very far from the rest of the knot, and shrink them to be very small. This done, we can ignore contributions to  $Z(K_c)$  coming from pairings in which any of the pairs connect the humps to the rest of the knot. Hence  $Z(K_c)$  factors to a part which is the same as in  $Z(K_s)$  times contributions that come from pairings that pair the ‘humpy’ part of  $K_c$  to itself. But as the following figure shows, for the same reasons as in section 4.3.3, these contributions are precisely  $Z(\infty)$ :



*Exercise 4.5.* Show that  $\tilde{Z}(K)$  is in fact real:  $\tilde{Z}(K) \in \mathcal{A}_{\mathbb{R}}^r$ .

*Hint 4.6.* Use the fact that the transformation  $t \rightarrow -t$ ,  $z \rightarrow \bar{z}$  maps a knot to an equivalent knot, while mapping  $\Omega_{n,n}$  to minus its conjugate.

*Remark 4.7.* Kontsevich [25], building on some work of Drinfel’d ([16] and [17]), has proven that  $\tilde{Z}(K)$  has rational coefficients.

#### 4.4. Proof of theorem 1.

4.4.1. *Proof of part (2) of theorem 1.* A weight system  $W$  of degree  $m$  is just a linear functional on  $\mathcal{G}_m \mathcal{A}_{\mathbb{R}}^r$ . Extend it by zero to all of  $\mathcal{A}_{\mathbb{R}}^r$ , and define

$$V(W)(K) = W(\tilde{Z}(K)). \quad \square$$

4.4.2. *Proof of part (3) of theorem 1.* Let  $W$  be a degree  $m$  weight system. To show that  $W = W_m(V(W))$  it is enough to show that if  $D \in \mathcal{G}_m \mathcal{D}^c$  is a chord diagram of degree  $m$  and if  $K_D$  is an embedding of  $D$  in the sense of section 2.2, then (for the natural extension of  $\tilde{Z}$  to knots with double points, definition 1.2):

$$\tilde{Z}(K_D) = \bar{D} + (\text{terms of degree } > m),$$

where  $\bar{D}$  is the class of  $D$  in  $\mathcal{A}_{\mathbb{R}}^r$ . In view of (18), it is enough to prove the same for  $Z$  rather than for  $\tilde{Z}$ . If two knots  $K^o$  and  $K^u$  are identical except that two of their strands form an overcrossing in  $K^o$  and an undercrossing in  $K^u$ , it is clear that the only contributions to  $Z(K^o) - Z(K^u)$  come from pairings in which these two strands are paired.  $Z(K_D)$  is a signed sum of  $Z$  evaluated on  $2^m$  knots, and this sum can be partitioned in pairs like the above  $K^{o,u}$  around  $m$  different crossings — and thus contributions to  $Z(K_D)$  come only from pairings that pair the strands near any of the  $m$  double points of  $K_D$ . This implies that the lowest degree contribution to  $Z(K_D)$  is at least of degree  $m$ . In degree  $m$  the pairing  $P$  is determined by the above restriction. It is easy to see that in that case  $D_P = D$ , and therefore the piece of degree precisely  $m$  in  $Z(K_D)$  is proportional to  $\bar{D}$ . It remains to

determine the constant of proportionality. This is a simple computation — in degree 1, the difference between  $Z(K^o)$  and  $Z(K^u)$  comes from the difference between integrating

$$\frac{dz - dz'}{z - z'}$$

along a contour in which  $z$  passes near but above  $z'$  and along a contour in which  $z$  passes near but under  $z'$ . By Cauchy's theorem this is  $2\pi i$ . Repeating this  $m$  times for each of the  $m$  double points of  $K_D$ , we get  $(2\pi i)^m$  and this exactly cancels the  $(2\pi i)^m$  in the denominator of (14). This proves that  $W = W_m(V(W))$ .

Next, let  $V$  be a Vassiliev invariant of type  $m$ . By the above discussion,

$$W_m(V - V(W_m(V))) = W_m(V) - W_m(V(W_m(V))) = W_m(V) - W_m(V) = 0.$$

In other words,  $V$  and  $V(W_m(V))$  differ by an invariant whose  $m$ th derivative vanishes. Namely, by an invariant of type  $m - 1$ .  $\square$

*Remark 4.8.* The argument in section 4.4.2 together with Kohno's theorem [24] immediately imply theorem 5, as was observed by Piunikhin [33].

**Problem 4.9.** Is Kontsevich's construction the same as the Reshetikhin-Turaev construction? In other words, consider the diagram:

$$\begin{array}{ccc} & \{\mathfrak{g}, R, m\} & \\ \text{theorem 4} \swarrow & & \searrow \text{theorem 5} \\ \mathcal{W} & \xleftrightarrow[\text{theorem 1}]{} & \mathcal{V} \end{array} \quad (19)$$

Theorem 5 says that the two ways of getting from a triple  $\{\mathfrak{g}, R, m\}$  to  $\mathcal{W}$  are the same. Is it also true that the two ways of getting from such a triple to  $\mathcal{V}$  are the same? Notice that in the Reshetikhin-Turaev construction  $\mathfrak{g}$ -invariance is broken by a choice of a Cartan sub-algebra (and then the result is shown not to depend on that choice), whereas in Kontsevich's construction (definition 4.4)  $\mathfrak{g}$ -invariance is never broken. This means that if the two constructions do agree, then Kontsevich's is perhaps somewhat "better". *Warning:* Low order computations show that for diagram (19) to commute,  $\tilde{Z}$  should be redefined to be  $Z(K)/Z(\infty)^{c/2}$ . This does not change the conclusions of theorems 13 and 1, but does invalidate theorem 9. It appears that [16, 17] imply a positive answer to this problem, but the details of such an argument are yet to be carefully checked.

**Problem 4.10.** Is Kontsevich's construction the same as the Chern-Simons construction alluded to in the beginning of section 4?

**Problem 4.11.** The statement of theorem 1 is purely combinatorial, and one would expect that the proof would also be combinatorial. I see the analytical proof given here as a temporary argument until a combinatorial proof will be found. See also problem 7.3.

## 5. THE PRIMITIVE ELEMENTS OF $\mathcal{A}$

**5.1. Theorem 8 from the perspective of Lie algebras.** One of the versions of the Poincare-Birkhoff-Witt (PBW) theorem (see *e.g.* [15, sec. 2.4.10]) says that the universal enveloping algebra  $\mathfrak{U}(\mathfrak{g})$  of a Lie algebra  $\mathfrak{g}$  is canonically isomorphic to the symmetric algebra

$\mathfrak{S}(\mathfrak{g})$  on  $\mathfrak{g}$ . The Lie algebra  $\mathfrak{g}$  acts on the two spaces in a compatible way, and therefore the  $\mathfrak{g}$ -invariant pieces of these two spaces are isomorphic:

$$\mathfrak{U}(\mathfrak{g})^{\mathfrak{g}} \simeq \mathfrak{S}(\mathfrak{g})^{\mathfrak{g}}. \quad (20)$$

The next exercise shows that in some sense the space  $\mathcal{A}$  corresponds to  $\mathfrak{U}(\mathfrak{g})^{\mathfrak{g}}$  and the space  $\mathcal{B}$  corresponds to  $\mathfrak{S}(\mathfrak{g})^{\mathfrak{g}}$ . In the light of (20) it is therefore not too surprising to find that  $\mathcal{A} \simeq \mathcal{B}$ .

*Exercise 5.1.* Given a Lie algebra  $\mathfrak{g}$  and a metric  $t$  construct natural maps  $\mathcal{T}_{\mathcal{A}} : \mathcal{A}^l \rightarrow \mathfrak{U}(\mathfrak{g})^{\mathfrak{g}}$  and  $\mathcal{T}_{\mathcal{B}} : \mathcal{B} \rightarrow \mathfrak{S}(\mathfrak{g})^{\mathfrak{g}}$ . Notice that these maps do not preserve the natural gradings of the spaces involved.

*Hint 5.2.* For  $\mathcal{T}_{\mathcal{A}}$  carry out the same procedure as in section 2.2, ignoring those parts of the procedure that mention the representation  $R$ . Given  $D \in \mathcal{D}^l$  get a tensor in some tensor power of  $\mathfrak{g}$ , and use the product of  $\mathfrak{U}(\mathfrak{g})$  to land in  $\mathfrak{U}(\mathfrak{g})^{\mathfrak{g}}$ . The map  $\mathcal{T}_{\mathcal{B}}$  is even easier, only that there is no natural ordering for the external vertices of a diagram  $C \in \mathcal{C}$ .

**Problem 5.3.** The maps  $\mathcal{T}_{\mathcal{A}}$  and  $\mathcal{T}_{\mathcal{B}}$  are not isomorphisms (see *e.g.* (30)), but I do not know how far they are from being isomorphisms.

**Problem 5.4.** Exercise 5.1 shows that to get a linear functional on  $\mathcal{A}^l$  it is sufficient to choose a linear functional on the center of  $\mathfrak{U}(\mathfrak{g})$  for some Lie algebra  $\mathfrak{g}$  that has an invariant metric. If  $\mathfrak{g}$  is semi-simple, it is well known that the space of such linear functionals is spanned by traces in finite dimensional representations and thus linear functionals on  $\mathfrak{U}(\mathfrak{g})^{\mathfrak{g}}$  give the same  $\mathcal{T}$ 's as in section 2.4. I do not know if this is still true if  $\mathfrak{g}$  is not semi-simple.

The proof of theorem 8 is more or less a direct translation of the proof of the above version of the PBW theorem to our language. It is instructive to read the following proof keeping Lie algebras in mind and noticing that  $\mathcal{T}_{\mathcal{A}}$  and  $\mathcal{T}_{\mathcal{B}}$  carry the isomorphism that we construct to the standard isomorphism (20).

**5.2. Proof of theorem 8.** We will construct a map  $\bar{\chi} : \mathcal{B} \rightarrow \mathcal{A}$  and a map  $\bar{\sigma} : \mathcal{A} \rightarrow \mathcal{B}$ , and then show that they are each other's inverse. For a Chinese character  $C \in \mathcal{C}$  that has  $k$  external vertices, define  $\chi(C)$  to be the sum of the  $k!$  ways of enclosing  $C$  in a circle. Each of these  $k!$  ways is obtained by choosing a bijection between the external vertices of  $C$  and the collection of  $k$ th roots of unity  $\{z : z^k = 1\}$ , and then gluing each external vertex of  $C$  to the corresponding point on the unit circle. For example,

$$\chi(\text{---}) = 2\text{---}, \quad \chi(\text{---}) = 16\text{---} + 8\text{---}, \quad \chi(\text{---}) = 3\text{---} + 3\text{---} = 0.$$

Theorem 6 implies that  $\chi$  descends to a map  $\bar{\chi} : \mathcal{B} \rightarrow \mathcal{A}$ .

The idea of the proof is to try to invert  $\bar{\chi}$ . The image of  $\bar{\chi}$  are the *symmetric* linear combinations of diagrams — those combinations that are stable under permuting external vertices. So given any diagram we will try to show that it is equivalent (modulo *STU* relations) to a symmetric combination. The obvious way to do that is to *symmetrize* — to compare each diagram with the sum of all of its ‘permuted versions’. We will show that the difference between a diagram and a permuted version of it is equivalent (mod *STU*) to a sum of diagrams with a lower number of external vertices (but this sum is *not* uniquely determined), and so we can ‘push the problems down’ and prove by induction. The most central of the technical details that have to be checked is the uniqueness of the procedure







Hence  $\Sigma 2$  follows from the anti-symmetry of internal vertices in  $\mathcal{B}$ , from the fact (23) that  $\Gamma_{D_{\pm}}$  does not touch the internal vertices of  $D_{\pm}$ , and from the induction hypothesis. The proof of  $\Sigma 3$  is similar.

**Proof of  $\Sigma 4$ :** Using the notation of (22), we need to show that  $\sigma_k(S_i D) = \sigma_k(TD - U_i D)$  for a diagram  $D \in \mathcal{D}_k$ :

$$\sigma_k(TD) - \sigma_k(U_i D) = \frac{1}{k!} \left( D^{CC} - (U_i D)^{CC} + \sum_{\pi \in \mathcal{S}_k} (\Lambda_D(\pi) - \Lambda_{U_i D}(\pi)) \right)$$

using the cocycle property (24) this is

$$= \frac{1}{k!} \sum_{\pi \in \mathcal{S}_k} (\Lambda_D(\pi) - \Lambda_D(U_i \pi) + \Lambda_D(U_i)) = \Lambda_D(U_i) = \sigma_{k-1}(\Gamma_D(U_i)) = \sigma_k(S_i D).$$

**Proof of  $\Sigma 5$ :** Let  $D \in \mathcal{D}_k$  be regarded as an element of  $\mathcal{D}_k/\mathcal{I}_k$ . Then

$$\bar{\chi}(\bar{\sigma}_k(D)) = \frac{1}{k!} \bar{\chi} \left( D^{CC} + \sum_{\pi \in \mathcal{S}_k} \Lambda_D(\pi) \right) = \frac{1}{k!} \sum_{\pi \in \mathcal{S}_k} (\pi D + (\bar{\chi} \circ \bar{\sigma}_{k-1})(\Gamma_D(U_{\pi})))$$

using the induction hypothesis ( $\Sigma 5$ ) and then lemma 5.5 this is

$$= \frac{1}{k!} \sum_{\pi \in \mathcal{S}_k} (\pi D + \Gamma_D(U_{\pi})) = \frac{1}{k!} \sum_{\pi \in \mathcal{S}_k} (\pi D + D - \pi D) = D.$$

This concludes the proof of the equivalence of  $\mathcal{A}$  and  $\mathcal{B}$ .

Let  $C = \bigcup_{i \in I} C_i$  be a presentation of a diagram  $C \in \mathcal{C}$  as the union of its connected components. Define a co-product  $\Delta_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C} \otimes \mathcal{C}$  by

$$\Delta_{\mathcal{C}}(C) = \sum_{J \subseteq I} \left( \bigcup_{i \in J} C_i \right) \otimes \left( \bigcup_{i \in I \setminus J} C_i \right).$$

It is easy to verify that

$$(\chi \otimes \chi) \circ \Delta_{\mathcal{C}} = \Delta \circ \chi. \quad (26)$$

This implies that under the equivalence of  $\mathcal{A}$  and  $\mathcal{B}$ ,  $\Delta_{\mathcal{C}}$  corresponds to the co-product of  $\mathcal{A}$ . From the definition of  $\Delta_{\mathcal{C}}$  it is easy to check that the primitive elements of the co-algebra  $\mathcal{B}$  are the equivalence classes of connected diagrams in  $\mathcal{C}$ . This concludes the proof of theorem 8.  $\square$

*Remark 5.6.* The statement of theorem 8 is due to M. Kontsevich [25], as well as the definition of the map  $\chi$ . The above proof is due to the author.

*Remark 5.7.* The product  $\cdot$  on  $\mathcal{A}$  does not (!) correspond to taking the union of diagrams in  $\mathcal{C}$ . Using disjoint union to make  $\mathcal{B}$  an algebra, this means that the maps  $\mathcal{A} \leftrightarrow \mathcal{B}$  are vector space isomorphisms but not algebra isomorphisms. Pulling back to  $\mathcal{A}$  the product of  $\mathcal{B}$ , we see that  $\mathcal{A}$  is an algebra in two different ways, both compatible with its co-product.


*Exercise 5.8.* Show that in the case of Lie algebras, the proof of theorem 8 can be changed in a minor way to yield a proof of the full PBW theorem

$$\mathfrak{U}(\mathfrak{g}) \simeq \mathfrak{S}(\mathfrak{g})$$

The idea is to allow an additional type of univalent vertices in  $\mathcal{A}^l$  and  $\mathcal{B}$ , marked by elements of  $\mathfrak{g}$ . Then mod out by a few more relations, and repeat essentially the same argument as above. Notice that, by a small further modification, this can be done even if the Lie algebra  $\mathfrak{g}$  does not carry a metric.

### 5.3. Proof of part (2) of theorem 9.

Let  $K_1$  and  $K_2$  be knots, and let  $K_1\#K_2$  denote their connected sum. Draw  $K_1\#K_2$  as on the right, and use the same argument as in the proof of theorem 13 to show that  $Z(K_1\#K_2) = Z(K_1)Z(K_2)$ , and therefore, after carefully counting critical points in  $K_1\#K_2$ ,  $\tilde{Z}(K_1\#K_2) = \tilde{Z}(K_1)\tilde{Z}(K_2)$ . If  $W$  is a primitive weight system, it satisfies  $\Delta(W) = W \otimes \epsilon + \epsilon \otimes W$ , and then by the definition of  $\Delta$  on  $\mathcal{A}^*$  we get:



$$\begin{aligned} V(W)(K_1\#K_2) &= W(\tilde{Z}(K_1\#K_2)) = W(\tilde{Z}(K_1)\tilde{Z}(K_2)) = \Delta(W)(\tilde{Z}(K_1) \otimes \tilde{Z}(K_2)) \\ &= W(\tilde{Z}(K_1))\epsilon(\tilde{Z}(K_2)) + \epsilon(\tilde{Z}(K_1))W(\tilde{Z}(K_2)) \\ &= V(W)(K_1) + V(W)(K_2). \quad \square \end{aligned}$$

## 6. HOW BIG ARE $\mathcal{A}$ , $\mathcal{W}$ AND $\mathcal{P}$ ?

6.1. **Numerical results.** 2,106 lines of C++ code and about 10 days of CPU time yield the following results for the dimensions of the various spaces involved [7]:

$m$	0	1	2	3	4	5	6	7	8	9
$\dim \mathcal{G}_m \mathcal{A}$	1	1	2	3	6	10	19	33	60	104
$\dim \mathcal{G}_m \mathcal{W}$	1	0	1	1	3	4	9	14	27	44
$\dim \mathcal{G}_m \mathcal{P}$	1	1	1	1	2	3	5	8	12	18
$\dim \mathcal{G}_m Lie$	1	1	2	3	6	10	19	33	60	104
diagrams	1	0	1	2	7	36	300	3,218	42,335	644,808
relations	0	0	0	2	15	144	1,645	21,930	334,908	5,056,798


*Remark 6.1.* • The space  $\mathcal{G}_m Lie$  is the subspace of  $(\mathcal{G}_m \mathcal{A})^*$  spanned by the Lie algebraic weight systems of theorem 4. In fact, I have only computed the weight systems corresponding to the classical groups and their representations, and so the above numbers also prove conjecture 2 up to degree 9.

- The numbers in the last two rows indicate the size of the matrices that had to be row reduced in order to compute  $\dim \mathcal{G}_m \mathcal{W}$ . These numbers have no real significance — they somewhat depend on the details of the algorithm chosen — and are displayed only so as to give an impression of the complexity of the problem.
- The dimensions of  $\mathcal{G}_m \mathcal{A}$  and  $\mathcal{G}_m \mathcal{P}$  were deduced from  $\dim \mathcal{G}_m \mathcal{W}$  using equation (5) and theorem 9.
- I wish to thank V. I. Arnold for correcting an earlier mistake I made in computing  $\dim \mathcal{G}_8 \mathcal{P}$ .
- The problem is highly exponential and it is unlikely that it will be possible to use the same techniques to compute  $\dim \mathcal{G}_{10} \mathcal{A}$ .

The last point makes it evident that a computer search is not the best way to generate weight systems. In the next section, I will present a very general construction of weight systems, which conceivably generates all of them.


6.2. **Marked surfaces.** In this section, we will construct the map  $\Phi : \mathcal{B} \rightarrow \mathcal{M}$ , promised in theorem 10. The easiest way to do so, is to factor it through the space  $\mathcal{B}^m$  of equivalence classes of *marked diagrams*.

**Definition 6.2.** A *marked diagram* is a Chinese character (see definition 1.11) whose arcs are marked either by the symbol '=' or by the symbol '×'. Sometimes we will mark an arc with more than one symbol. In this case, it is understood that an even number of ×'s (and an arbitrary number of ='s) on a single arc is equivalent to a single '=' on that arc, while an odd number of ×'s (and an arbitrary number of ='s) is equivalent to a single '×'. (See figure 14). The collection of all marked diagrams is denoted by  $\mathcal{C}^m$  and is graded in the same way as  $\mathcal{C}$ .




**Figure 14.** Two versions of the same degree 2 marked diagram.

**Definition 6.3.**  $\mathcal{B}^m$  is the quotient of  $\text{span } \mathcal{C}^m$  by the subspace spanned by the relations:



In these marked diagrams the symbol 'o' stands for an arbitrary additional marking, which is the same on corresponding arcs on both sides of the same equation.

Notice that the following two additional relations are satisfied in  $\mathcal{B}^m$ :



**Definition 6.4.** Let  $\mu : \mathcal{C} \rightarrow \mathcal{B}^m$  be the map defined by

$$(\text{-----}) \rightarrow (\text{-----}) - (\text{-----}).$$

In other words, a diagram  $C$  in  $\mathcal{C}$  having  $e$  edges is mapped to the (signed) sum of the  $2^e$  ways of marking  $C$ .

**Proposition 6.5.** The map  $\mu$  descends to a map  $\mu : \mathcal{B} \rightarrow \mathcal{B}^m$ .



**Definition 6.9.** Define the map  $\Phi : \mathcal{B} \rightarrow \mathcal{M}$  to be the composition  $\Phi = \tau \circ \mu$ .

For the purpose of conjecture 2, much of the information in  $\mathcal{M}$  is superfluous:

*Exercise 6.10.* For a marked surface  $M \in \mathcal{M}$  define a linear functional  $\check{M} \in \mathcal{M}^*$  by

$$\check{M}(M') = \begin{cases} 1 & M' \simeq M, \\ 0 & \text{otherwise.} \end{cases}$$

Let  $M_{\uparrow}$  and  $M_{\downarrow}$  be two marked surfaces that differ only by the orientation of one of their marks. Then the corresponding linear functionals on  $\mathcal{B}$  differ only by a sign:

$$\Phi^* \check{M}_{\uparrow} + \Phi^* \check{M}_{\downarrow} = 0.$$

**Definition 6.11.** Each tangent to the boundary of a connected surface  $M$  defines an orientation on the boundary component on which it lives, and if  $M$  is orientable, such a tangent taken together with the outward pointing normal defines an orientation on  $M$ . Call a connected marked surface  $M$  *normalized* if either of the following holds:

- $M$  is orientable and all its markings generate the same orientation on  $M$ .
- $M$  is non-orientable and its marking generate *consistent* orientations on each boundary component separately.

A general marked surface is called *normalized* if all of its connected components are normalized. It is called *normalized orientable* if all of its connected components are normalized and orientable.

*Remark 6.12.* Exercise 6.10 shows that in order to study  $\Phi^* \mathcal{M}^*$  it is sufficient to consider only the linear functionals corresponding to normalized marked surfaces.

*Exercise 6.13.* Let  $M$  be a connected marked surface of Euler characteristic  $\chi$  and genus  $g$ . If  $M$  has  $b$  boundary components and  $n$  tangents marked on its boundary define the degree  $m$  of  $M$  by


$$m = n - \chi = \begin{cases} 2g + b + n - 2 & \text{if } M \text{ is orientable,} \\ g + b + n - 2 & \text{otherwise.} \end{cases}$$

Define the degree of a general marked surface to be the sum of the degrees of its connected components, and check that with this grading on  $\mathcal{M}$ , the map  $\Phi$  preserves degrees.

*Remark 6.14.* The definition of the thickening map has a natural extension (also denoted by  $\tau$ ) to marked diagrams which are allowed to have some  $p$  cycles of directed full lines. Simply thicken dashed lines as before, and thicken directed lines to semi-open strips, so that every full line  $l$  is replaced by  $l \times [0, 1)$ . The result is a *punctured* marked surface — a marked surface with  $p$  points removed. See figure 16.

*Exercise 6.15.* Check that the extended map  $\tau$  satisfies the *STU* relation and thus descends to a map defined on the relevant quotient space.

**6.3. The classical Lie algebras.** The purpose of this section is to show how to compute  $\mathcal{T}(D)$  for an arbitrary diagram  $D \in \mathcal{D}^t$  and a large number of representations of the algebras in the families *gl*, *su*, *sp*, and *so*. I will show in detail the computations for *so*, and just state the results for *gl*, *su*, and *sp*.



**Figure 16.** Thickening full lines. *A* shows how to thicken dashed-full vertices, in *B* the resulting surface is a punctured sphere with three holes while in *C* it is a punctured torus. In all cases a dotted line represents an ‘open end’ of the surface.

6.3.1. *Chord diagrams in the  $N$  dimensional representation of  $so(N)$ .* For a start, let us consider only chord diagrams. Let  $\Lambda^1$  be the defining representation of  $so(N)$  for some  $N$ , and let  $t$  be given by matrix trace in that representation<sup>12</sup>,

$$t(a, b) = \text{tr} \Lambda^1(a) \Lambda^1(b).$$

The basic building block of chord diagrams is the tensor

$$\mathcal{T}_{\beta\delta}^{\alpha\gamma} = r_{b\beta}^{\alpha} (t^{-1})^{ab} r_{a\delta}^{\gamma} = \begin{array}{c} \alpha \\ \downarrow \text{---} \uparrow \\ \beta \end{array} \begin{array}{c} \delta \\ \downarrow \text{---} \uparrow \\ \gamma \end{array} \in ((\Lambda^1)^*)^{\otimes 2} \otimes (\Lambda^1)^{\otimes 2},$$

where  $\alpha$  and  $\gamma$  run over a basis of  $(\Lambda^1)^*$ ,  $\beta$  and  $\delta$  run over a basis of  $\Lambda^1$ ,  $a$  and  $b$  run over a basis of  $so(N)$ , and  $r_{b\beta}^{\alpha}$  and  $(t^{-1})^{ab}$  are the tensors considered in section 2.4, expressed in coordinates.

A convenient choice of generators for  $so(N)$  are the  $N \times N$  matrices  $M_{ij}$  ( $i < j$ ), given by

$$(M_{ij})_{\alpha\beta} = \delta_{i\alpha} \delta_{j\beta} - \delta_{i\beta} \delta_{j\alpha}.$$

That is, the  $ij$  entry of  $M_{ij}$  is  $+1$ , the  $ji$  entry of  $M_{ij}$  is  $-1$ , and all other entries of  $M_{ij}$  are zero. The invariant bilinear form that we pick on  $so(N)$  is the matrix trace in the defining representation, and so

$$t_{(ij)(kl)} \stackrel{\text{def}}{=} \text{tr} (M_{ij} M_{kl}) = -2\delta_{ik} \delta_{jl}.$$

Inverting the  $\frac{N(N-1)}{2} \times \frac{N(N-1)}{2}$  matrix  $t_{(ij)(kl)}$  we get

$$(t^{-1})^{(ij)(kl)} = -\frac{1}{2} \delta^{ik} \delta^{jl}, \quad (27)$$

and so

$$\mathcal{T}_{\beta\delta}^{\alpha\gamma} = \sum_{i < j; k < l} (t^{-1})^{(ij)(kl)} (M_{ij})_{\alpha\beta} (M_{ij})_{\gamma\delta}. \quad (28)$$

Using (27) and some algebraic manipulations we can simplify (28), and then represent it by a diagram:

$$(28) = \frac{1}{2} (\delta_{\alpha\delta} \delta_{\beta\gamma} - \delta_{\alpha\gamma} \delta_{\beta\delta}) = \frac{1}{2} \left( \begin{array}{c} \alpha \quad \delta \\ \frown \quad \smile \\ \beta \quad \gamma \end{array} - \begin{array}{c} \alpha \quad \delta \\ \times \\ \beta \quad \gamma \end{array} \right). \quad (29)$$

The last thing to note is that  $\mathcal{T}(k \text{ disjoint circles}) = N^k$ . These rules are sufficient to compute  $\mathcal{T}(D)$  for any chord diagram  $D$ .

*Example 6.16.*

$$\begin{array}{c} \circ \times \end{array} \rightarrow \frac{1}{4} \begin{array}{c} \circ \otimes \end{array} - \frac{1}{2} \begin{array}{c} \circ \otimes \end{array} + \frac{1}{4} \begin{array}{c} \circ \otimes \end{array} \rightarrow \frac{N}{4} - \frac{N}{2} + \frac{N^2}{4} = \frac{N(N-1)}{4}.$$

<sup>12</sup>In a simple Lie algebra, an invariant metric is always a multiple of the Killing form. It is easy to check that if  $t \rightarrow \kappa t$ , then  $\mathcal{T}(D) \rightarrow \kappa^{-\text{deg } D} \mathcal{T}(D)$ , and so our choice of  $t$  is as good as any other choice.

6.3.2. *Diagrams with trivalent vertices.* By proposition 2.10 the tensor  $\text{---}\bigcirc\text{---}$  is invariant, and therefore if the Lie algebra  $\mathfrak{g}$  is simple, it must be a multiple of  $t = \text{---}$ . In that case, let  $\kappa$  be the constant for which

$$\text{---}\bigcirc\text{---} = \kappa \text{---}. \quad (30)$$

Using this definition, diagrams that have trivalent vertices can be reduced to diagrams without trivalent vertices:

$$\text{---}\bigcirc\text{---} = \frac{1}{\kappa^3} \text{---}\bigcirc\text{---} \stackrel{STU}{=} \frac{1}{\kappa^3} \left( \text{---}\bigcirc\text{---} - \text{---}\bigcirc\text{---} \right). \quad (31)$$

(The above equality should be read as an equality between the tensors represented by these diagrams.)

In the case of the  $N$  dimensional representation of  $so(N)$ , a comparison of  $\text{---}\bigcirc\text{---}$  and  $\text{---}\bigcirc\text{---}$  (using (29)) shows that  $\kappa = 1$ . Furthermore, in this case (31) can be further reduced using (29). The result is:

$$\text{---}\bigcirc\text{---} = \frac{1}{4} \left( \text{---}\bigcirc\text{---} - \text{---}\bigcirc\text{---} - \text{---}\bigcirc\text{---} + \text{---}\bigcirc\text{---} - \text{---}\bigcirc\text{---} + \text{---}\bigcirc\text{---} + \text{---}\bigcirc\text{---} - \text{---}\bigcirc\text{---} \right). \quad (32)$$

In other words for  $D \in \mathcal{D}^t$ ,

$$\mathcal{T}(D) = 2^{v-e} \sum_M s_M N^{b(\tau D_M)}, \quad (33)$$

where

- $v$  ( $e$ ) is the number of internal vertices (edges) in  $D$ .
- The sum is over all possible markings  $M$  of the dashed lines of  $D$ , as in definition 6.4.
- $s_M$  is the sign corresponding to  $M$  as in definition 6.4.
- $b(\tau D_M)$  is the number of boundary components of the thickening  $\tau(D_M)$  of  $\tau D_M$ , defined in remark 6.14.

*Example 6.17.*  $\mathcal{T}(\text{---}\bigcirc\text{---})$  can be computed in the following manner:

$$2^{1-3} \left( \text{---}\bigcirc\text{---} - 3 \text{---}\bigcirc\text{---} + 3 \text{---}\bigcirc\text{---} - \text{---}\bigcirc\text{---} \right) \rightarrow 2^{-2} (N^3 - 3N^2 + 3N - N) = \frac{N(N-1)(N-2)}{4}.$$

6.3.3. *Tensor products of representations.* Recall that as in remark 2.12 we can consider diagrams in which each directed arc is colored by a (possibly) different representation. The following proposition can be used to reduce computations with arcs colored by a tensor product  $R_1 \otimes R_2$  to computations with arcs colored by  $R_1$  or  $R_2$ :

**Proposition 6.18.** *Let  $R_1$  and  $R_2$  be two representations of some Lie algebra  $\mathfrak{g}$ . The following is an equality between tensors in  $\mathfrak{g}^* \otimes R_1 \otimes R_2 \otimes R_1^* \otimes R_2^*$ :*

$$\mathfrak{g}^* \text{---} \left( \begin{array}{c} R_1 \otimes R_2 \\ \text{---} \\ (R_1 \otimes R_2)^* \end{array} \right) = \mathfrak{g}^* \left( \begin{array}{c} R_1 \\ \text{---} \\ R_1^* \end{array} \right) \left( \begin{array}{c} R_2 \\ \text{---} \\ R_2^* \end{array} \right) + \mathfrak{g}^* \left( \begin{array}{c} R_1 \\ \text{---} \\ R_1^* \end{array} \right) \left( \begin{array}{c} R_2 \\ \text{---} \\ R_2^* \end{array} \right).$$

*Proof.* Follows from the definition of the action of  $\mathfrak{g}$  on a tensor product,  $(R_1 \otimes R_2)(\mathfrak{g}_a) = R_1(\mathfrak{g}_a) \otimes 1 + 1 \otimes R_2(\mathfrak{g}_a)$ .  $\square$

*Example 6.19.* Let us compute  $\mathcal{T}(\text{---}\bigcirc\text{---})$  in the second tensor power of the defining representation of  $so(N)$ . In the formula below, full lines are colored by the *defining* representation.

$$\mathcal{T}(\text{---}\bigcirc\text{---}) = \mathcal{T}(\text{---}\bigoplus\text{---} + 2\text{---}\bigoplus\text{---} + \text{---}\bigoplus\text{---}) = \frac{1}{2}N^2(N-1) + 0 + \frac{1}{2}N^2(N-1) = N^2(N-1).$$

*Exercise 6.20.* Verify that  $\mathcal{T}_{(\Lambda^1)^{\otimes 2}}(\text{---}\bigcirc\text{---}) = N(N-1)(N^2 - N + 2)$ .

**Definition 6.21.** Let  $K : (S^1 = \{z : |z| = 1\}) \rightarrow \mathbf{R}^3$  be a framed knot and let  $n(z)$  ( $z \in S^1$ ) be a section of the normal bundle of  $K$  compatible with its framing. For a non-zero integer  $q$  define the  $q$ th *disconnected cabling*  $K^{\otimes q}$  of  $K$  to be the  $q$ -component link whose  $j$ th component  $K_j^{\otimes q}$  is given by

$$K_j^{\otimes q}(z) = K(z + j\epsilon n(z)),$$

where  $\epsilon$  is a very small number. Let  $K^{\otimes q}$  inherit the framing of  $K$ .

*Exercise 6.22.* Show that if  $V$  is a Vassiliev invariant of type  $m$  of framed knots then  $V \circ (K \rightarrow K^{\otimes q})$  is also a Vassiliev invariant of type  $m$ . Show that the operations  $K \rightarrow K^{\otimes q}$  on knots and  $R \rightarrow R^{\otimes q}$  on representations are adjoints of each other — show that if  $V$  is a Vassiliev invariant of type  $m$  of framed knots for which  $W_m(V) = W_{\mathfrak{g}, R, m}$ , then  $W_m(V \circ (K \rightarrow K^{\otimes q})) = W_{\mathfrak{g}, R^{\otimes q}, m}$ .

6.3.4. *The Adams operations.* Let  $q$  be a non-negative integer. If  $\chi$  is a virtual character on a compact Lie group  $G$  (*i.e.*, a conjugation invariant  $L^2$  function on  $G$ ), then so is  $\bar{\psi}^q \chi$ , which is defined by  $(\bar{\psi}^q \chi)(g) = \chi(g^q)$ . This defines an operation  $\bar{\psi}^q$  called *the  $q$ 'th Adams operation* on the representation ring  $R(G)$  of  $G$  (see *e.g.* [11, pp. 104]). We are interested in this operation on the Lie algebra level. Recall that every character (and hence every virtual character  $\chi$ ) induces a linear functional  $\phi(\chi)$  on the universal enveloping algebra  $\mathfrak{U}(\mathfrak{g})$  of the Lie algebra  $\mathfrak{g}$  of  $G$ .

*Exercise 6.23.* Let  $m$  denote the product of and let  $\Delta_{\mathfrak{u}}$  denote the co-product (see *e.g.* [31]) of  $\mathfrak{U}(\mathfrak{g})$ , and let  $\hat{\psi}^q : \mathfrak{U}(\mathfrak{g}) \rightarrow \mathfrak{U}(\mathfrak{g})$  be the composition

$$\mathfrak{U}(\mathfrak{g}) \xrightarrow{\Delta_{\mathfrak{u}}} \mathfrak{U}(\mathfrak{g}) \otimes \mathfrak{U}(\mathfrak{g}) \xrightarrow{1 \otimes \Delta_{\mathfrak{u}}} \dots \xrightarrow{1^{\otimes (q-2)} \otimes \Delta_{\mathfrak{u}}} \mathfrak{U}(\mathfrak{g})^{\otimes q} \xrightarrow{m} \mathfrak{U}(\mathfrak{g}).$$

Then the operations  $\bar{\psi}^q$  and  $\hat{\psi}^q$  correspond under  $\phi$ . In other words,

$$\phi(\bar{\psi}^q \chi) = (\hat{\psi}^q)^* \phi(\chi).$$

*Exercise 6.24.* Fix a Lie algebra  $\mathfrak{g}$  and a metric  $t$ , let  $D$  be a diagram with a distinguished cycle of directed lines, let  $\mathcal{T}_{\chi}(D)$  denote the tensor associated with the diagram  $D$  when its distinguished cycle is colored with the virtual representation  $\chi$ , and recall that  $\psi^k D$  was defined in definition 3.11. Show that

$$\mathcal{T}_{(\hat{\psi}^q)^* \chi}(D) = \mathcal{T}_{\chi}(\psi^q D).$$

*Example 6.25.* For  $\mathfrak{g} = so(N)$  and  $\Lambda^1$  the defining representation,

$$\mathcal{T}_{(\hat{\psi}^2)^* \Lambda^1}(\text{---}\bigcirc\text{---}) = \mathcal{T}_{\Lambda^1}(\psi^2 \text{---}\bigcirc\text{---}) = 12\mathcal{T}_{\Lambda^1}(\text{---}\bigcirc\text{---}) + 4\mathcal{T}_{\Lambda^1}(\text{---}\bigotimes\text{---}) = N(N-1)(3N-2).$$

In [2] it is shown that in an arbitrary *special*  $\lambda$ -ring the operations  $\psi^q$  are given by universal polynomials in the ‘exterior power’ operations  $\Lambda^q$ . In fact, the opposite is also true and the  $\Lambda^q$ 's can also be expressed in terms of the  $\psi^q$ 's. We have simple combinatorial algorithms



6.3.6. *Other Lie algebras.* So far we have dealt only with a single Lie algebra, and varied its representation. Let us now try to vary the algebra:

*Exercise 6.32.* Let  $(\mathfrak{g}_{\mathbf{C}}, R_{\mathbf{C}})$  be the complexification of a pair  $(\mathfrak{g}, R)$  consisting of a real Lie algebra  $\mathfrak{g}$  and a representation  $R$  of  $\mathfrak{g}$  in a real vector space. Show that  $W_{\mathfrak{g}_{\mathbf{C}}, R_{\mathbf{C}}, m} = W_{\mathfrak{g}, R, m}$  for every  $m$  (and therefore no new information can be gained by studying, say, the various real forms of  $so(N, \mathbf{C})$ ).

*Exercise 6.33.* External tensor products correspond to the co-product on  $\mathcal{A}$  (or equivalently, to the product in  $\mathcal{A}^*$ ). In other words, let  $\mathfrak{g}_1$  and  $\mathfrak{g}_2$  be Lie algebras, let  $t_1$  and  $t_2$  be metrics and let  $R_1$  and  $R_2$  be representations of  $\mathfrak{g}_1$  and  $\mathfrak{g}_2$  respectively. Consider the Lie algebra  $\mathfrak{g}_1 \oplus \mathfrak{g}_2$  with the metric  $t_1 \oplus t_2$  and the external tensor product representation  $R_1 \otimes R_2$ . Then the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\Delta} & \mathcal{A} \otimes \mathcal{A} \\ \mathcal{T}_{R_1 \otimes R_2} \downarrow & & \downarrow \mathcal{T}_{R_1} \otimes \mathcal{T}_{R_2} \\ \mathbf{C} & \xleftarrow{m} & \mathbf{C} \otimes \mathbf{C} \end{array}$$

*Exercise 6.34.* Show that a linear functional  $\omega$  on  $\mathcal{A}^*$  associated to some one dimensional representation of a one dimensional Abelian Lie algebra is an (easily describable) unit in the algebra  $\mathcal{A}^*$ . Show that the same linear functionals on  $\mathcal{G}_m \mathcal{A}^*$  appear as  $W_m(V)$  where  $V$  is the coefficient of  $x^m$  in a framed-knot invariant of the form  $K \rightarrow \exp(rxw(K))$ ,  $r$  is an arbitrary constant, and  $w(K)$  is the *writhe* (see e.g. [22]) of  $K$ . Show that  $\omega \in (\Phi \circ \bar{\sigma})^* \mathcal{M}^*$ .

*Exercise 6.35.* Check that higher dimensional Abelian Lie algebras give the same linear functionals on  $\mathcal{A}$  as one dimensional Lie algebras.

Exercises 6.33 and 6.34 taken together show that (in our context) studying the *sl* family is essentially equivalent to studying the *gl* family. So let us pick one of them:

*Exercise 6.36.* The  $gl(N)$  analogs of equations (29), (32) and (33) are

$$\mathcal{T}_{\beta\delta}^{\alpha\gamma} = \begin{array}{c} \alpha \\ | \\ \beta \end{array} \begin{array}{c} \delta \\ \cdots \\ \gamma \end{array} = \begin{array}{c} \alpha \quad \delta \\ \frown \quad \smile \\ \beta \quad \gamma \end{array}, \quad (34)$$

$$\begin{array}{c} \curvearrowright \\ \vdots \\ \curvearrowleft \end{array} = \begin{array}{c} 0 \\ \text{---} \\ \text{---} \\ \text{---} \end{array} - \begin{array}{c} 1 \\ \text{---} \\ \text{---} \\ \text{---} \end{array}, \quad (35)$$

and

$$\mathcal{T}(D) = \sum_M (-1)^{s_M} N^{b(\tau D_M)}, \quad (36)$$

where here

- The sum is over all possible markings  $M$  of the internal vertices of  $D$  by the digits 0 and 1.
- $s_M$  is the sum of all the digits in  $M$ .
- $b(\tau D_M)$  is the number of boundary components of the thickening  $\tau(D_M)$  of  $\tau D_M$ . Internal vertices are thickened depending on the digit marked on them using one of the two possibilities in (35) and chords are always thickened as if they were marked by an '=' symbol.

Notice that the exterior powers of the defining representation of  $gl(N)$  generate its representation ring, and thus we have effective computation techniques for  $gl(N)$  in all of its representations.

*Exercise 6.37.* Find the analogs of equations (29), (32) and (33) for the Lie algebra  $sp(N)$  or read them in Bar-Natan, [5, 6]. Use straightforward (but rather tricky) combinatorics to show that if  $D$  is a diagram with no univalent vertices but with an arbitrary number of cycles of directed lines marked by the defining representation, then

$$\mathcal{T}_{sp(N)}(D) = \mathcal{T}_{so(-2N)}(D) \quad (37)$$

where  $\mathcal{T}_{so(-2N)}$  is defined as in (33), only with  $(-2N)$  replacing  $N$ .

**Problem 6.38.** I learned about equation (37) from M. Kontsevich [25]. He claims that the above result follows from studying the super Lie algebra  $Osp(m, n)$ . I believe, but as of now I can not reproduce his result.

Notice that the exterior powers of the defining representation of  $sp(N)$  generate its representation ring, and thus we have effective computation techniques for  $sp(N)$  in all of its representations.

**Problem 6.39.** The results of Kuperberg [27] can be used to derive an explicit algorithm for computations using the rank 2 exceptional Lie algebra  $G_2$  (see also [35]). I do not know whether the results are inside  $(\Phi \circ \bar{\sigma})^* \mathcal{M}^*$ .

**6.4. Marked surfaces and the classical Lie algebras.** In this section we will prove theorem 11 and a few related results. We will do so by constructing a vector space  $\mathcal{L}$  and a pairing  $\langle \cdot, \cdot \rangle : \mathcal{M} \otimes \mathcal{L} \rightarrow \mathbf{F}$  that has the following properties:

**Proposition 6.40.** (Proof on page 45) For an element  $L \in \mathcal{L}$  denote by  $\Psi(L)$  the pullback to  $\mathcal{B}^*$  via the map  $\Phi$  of the linear functional  $(M \mapsto \langle M, L \rangle) \in \mathcal{M}^*$ . Then for every list of triples  $\{(\mathfrak{g}_i, R_i, m_i)\}$  where each  $\mathfrak{g}_i$  is either an  $so$  or a  $gl$  algebra with  $R$  a single-valued representation of the corresponding group there is a (non-unique) generator  $L \in \mathcal{L}$  for which

$$\Psi(L) = \bar{\chi}^* \prod_i W_{\mathfrak{g}_i, R_i, m_i}. \quad (38)$$

Conversely, for every generator  $L \in \mathcal{L}$  there is a (canonically determined) list of triples  $\{(\mathfrak{g}_i, R_i, m_i)\}$  for which (38) holds. (For the definition of the isomorphism  $\bar{\chi}$  see page 29).

**Proposition 6.41.** (Follows from (37) and exercises 6.34 and 6.51) Let  $m$  be a non-negative integer. If  $\mathfrak{g}$  is a symplectic or an Abelian algebra and  $R$  is a representation of  $\mathfrak{g}$  or if  $\mathfrak{g}$  is in the family  $so$  and  $R$  is a spin representation, then  $W_{\mathfrak{g}, R, m} \in (\Phi \circ \bar{\sigma})^* \mathcal{M}^*$ .


**Proposition 6.42.** (Proof on page 46) The pairing  $\langle \cdot, \cdot \rangle$  is non-degenerate in the sense that if  $\bar{M}_1$  and  $\bar{M}_2$  are linear combinations of normalized marked surfaces and  $\langle \bar{M}_1, L \rangle = \langle \bar{M}_2, L \rangle$  for every  $L \in \mathcal{L}$ , then  $\bar{M}_1 = \bar{M}_2$ .

Clearly, these three propositions together with remark 6.12 imply theorem 11.

**6.4.1. The pairing  $\langle \cdot, \cdot \rangle$ .** By a *covering* of the circle we will mean a not necessarily connected oriented covering of the oriented circle, regarded up to an orientation preserving homeomorphism. Such coverings can be identified with non-empty finite sets<sup>14</sup> of integers measuring

<sup>14</sup>In this section the word ‘set’ means ‘an unordered collection with multiplicities’. I.e.,  $\{2, 2, 1\} = \{2, 1, 2\} = \{1, 2^2\} \neq \{1, 2\}$ . Notice that for a set  $\{q, \dots, q\}$  of size  $\beta$  we sometimes use the notation  $\{q^\beta\}$ .

the *multi-degree* of a cover:



*Exercise 6.43.* Explain how coverings of the circle correspond to general cablings of framed knots, generalizing definitions 3.13 and 6.21. Generalize exercises 3.14 and 6.22 to general cablings.

A *labeled covering* will be a quadruple  $L = (s, N, L_0, m)$  where  $L_0$  is a covering of the circle,  $m$  is a non-negative integer called ‘the degree of  $L$ ’,  $s$  is called ‘the symbol of  $L$ ’ and is either the symbol ‘ $gl$ ’ or the symbol ‘ $so$ ’, and  $N$  is an integer. A *covering bouquet* is a finite non-empty set of labeled coverings of the circle — something that looks like

$$\{(gl, 54, \{2, 2, 1\}, 6), (so, 9, \{7, 3\}, 14), \dots\}.$$

**Definition 6.44.** Let  $\mathcal{L}$  be the vector space spanned by the collection of all covering bouquets.

**Definition 6.45.** Let  $L = (s, N, L_0 = \{q_1, \dots, q_k\}, m)$  be a labeled covering of the circle and let  $M$  be a marked surface with  $n$  marked tangents. Define

$$\langle M, L \rangle = \begin{cases} 0 & \text{if } \deg M \neq m, \\ 0 & \text{if } s = 'gl' \text{ and } M \text{ is not} \\ & \text{normalized orientable,} \\ \sum_{\pi \in \mathcal{S}_n} \sum_{l \text{ of } \pi} N^{b(\tau L_0 \cup_l M)} & \text{otherwise,} \end{cases} \quad (39)$$

where:

- The outer sum is over the  $n!$  ways (each indexed by a permutation  $\pi \in \mathcal{S}_n$ ) of arranging the  $n$  markings of  $M$  around the base  $S^1$  of the covering  $L_0$ .
- The inner sum is over the  $(\sum |q_i|)^n$  ways of lifting  $\pi$  from the base  $S^1$  to the covering itself.
- $\tau L_0$  is the thickening of  $L_0$  into a disjoint union of  $k$  punctured disks as described in exercise 6.14.
- $\tau L_0 \cup_l M$  is the punctured surface (with boundary) obtained by gluing  $M$  to  $\tau L_0$  in the  $n$  sites specified by  $l$ , in such a way that in each such site the orientation given on  $L_0$  matches with the corresponding marking of  $M$ .
- $b(\tau L_0 \cup_l M)$  is the number of boundary components of  $\tau L_0 \cup_l M$ .

*Example 6.46.* Let us compute the pairing of a disk marked by three aligned tangents with the labeled covering  $(gl, N, \{1, 1\}, 2)$ . The sum is over 48 terms, but after dividing by the 3-fold cyclic symmetry of the circle and by the possibility to exchange the two leaves of the

covering, we get only 4 types of contributions. Observed from above, this is how it looks:

$$\langle \text{Y-junction with circle}, \text{circle with dot} \rangle \rightarrow 6 \text{ (Term 1)} + 18 \text{ (Term 2)} + 6 \text{ (Term 3)} + 18 \text{ (Term 4)}$$

$$\rightarrow 6N^4 + 18N^2 + 6N^2 + 18N^2 = 6N^2(N^2 + 7).$$

**Definition 6.47.** Let  $\{M_i\}_{i=1}^c$  be the connected components of a general marked surface  $M \in \mathcal{M}$  and  $L = \{L_k\}_{k=1}^\gamma$  be a presentation of a covering bouquet as a set of labeled coverings. Define

$$\langle M, L \rangle = \sum_{k_1, \dots, k_c=1}^\gamma \prod_{k=1}^\gamma \left\langle \bigcup_{\{i:k_i=k\}} M_i, L_k \right\rangle,$$

and extend  $\langle \cdot, \cdot \rangle$  linearly to all of  $\mathcal{M} \otimes \mathcal{L}$ .

6.4.2. *Proof of proposition 6.40.*

**Lemma 6.48.** For any  $m$  and  $N$ ,  $\Psi(\{(so, N, \{1\}, m)\}) = W_{so(N), \Lambda^1, m}$ .

*Proof.* Follows immediately from the definition of the pairing  $\langle \cdot, \cdot \rangle$ , from the definition (6.9) of the map  $\Phi$ , from the definition of  $\bar{\chi}$ , and from (33).  $\square$

**Lemma 6.49.** For any  $m$  and  $N$ ,  $\Psi(\{(gl, N, \{1\}, m)\}) = W_{gl(N), \Lambda^1, m}$ .

*Proof.* Consider a Chinese character  $C$  as a one dimensional CW-complex. A marking  $\tilde{C}$  of  $C$  defines a co-chain  $\beta(\tilde{C}) \in C^1(C, \mathbf{Z}/2\mathbf{Z})$  — for an edge  $l$  in  $C$  define

$$\beta(\tilde{C})(l) = \begin{cases} 0 & l \text{ is marked by an '=' symbol,} \\ 1 & l \text{ is marked by a 'x' symbol.} \end{cases}$$

It is easy to check that the thickening  $\tau\tilde{C}$  of  $\tilde{C}$  is orientable iff  $\beta(\tilde{C})$  is a co-boundary, *i.e.* iff there exists an  $\alpha \in C^0(C, \mathbf{Z}/2\mathbf{Z})$  for which  $\beta(\tilde{C}) = d\alpha$ . Also,  $\tau\tilde{C}$  is normalized orientable iff such an  $\alpha$  can be found which vanishes on the external (univalent) vertices of  $C$ . In such a case,  $\alpha$  is uniquely determined by  $\beta(\tilde{C})$ . In computing  $\Psi(\{(gl, N, \{1\}, m)\})$  a signed summation over normalized orientable thickenings of  $C$  is performed (see definition 6.4 and 6.45), and by the above discussion it can be replaced by a signed summation over  $\alpha \in C^0(\text{int } C, \mathbf{Z}/2\mathbf{Z})$ . Comparing with (36), we see that we are done.  $\square$

*Exercise 6.50.* Let  $\Lambda^1$  be the defining representation of  $so(N)$  or  $gl(N)$ , and let  $P^q$  denote the virtual representation  $(\hat{\psi}^q)^* \Lambda^1$ . Use proposition 6.18, exercises 6.23 and 6.33 and equation (26) to prove the obvious generalization of the statement:

$$\Psi(\{(gl, 54, \{2, 2, 1\}, 6), (so, 9, \{7, 3\}, 14)\}) = \bar{\chi}^*(W_{gl(54), P^2 \otimes P^2 \otimes P^1, 6} \cdot W_{so(9), P^7 \otimes P^3, 14}).$$

Clearly, lemmas 6.48 and 6.49 and exercise 6.50 prove proposition 6.40. The following exercise is the remaining ingredient for the proof of proposition 6.41:

*Exercise 6.51.* The crux of the argument leading to computations for the Lie algebra  $so$  is equation (32), used to simplify internal vertices of a diagram. This equation is valid no matter what representation is used on the external circle surrounding a diagram in  $\mathcal{D}^t$ . Use this fact to show that if  $R$  is a spin representation of  $so(N)$  (or, in fact, any representation of  $so(N)$ ), then  $W_{so(N), R, m} \in (\Phi \circ \bar{\sigma})^* \mathcal{M}^*$ .

6.4.3. *Proof of proposition 6.42.* The idea is to show that all relevant information about a normalized marked surface  $M$  can be read from the numbers  $\langle M, L \rangle$  for various  $L$ 's. Keeping exercise 6.13 and the classification of 2D surfaces (see *e.g.* [30]) in mind we see that (for a connected normalized surface) it is enough to read its degree, whether it is orientable, and the number of markings on each of its boundary components.

*Exercise 6.52.* Let  $M$  be a disk with  $n$  aligned tangents marked on its boundary (so that  $\deg M = n - 1$ ). Prove that the highest power of  $N$  in  $\langle M, (so \text{ or } gl, N, \{1\}, n - 1) \rangle$  is  $N^n$  and that in the sum (39) this power is achieved only when the lifting  $l$  arranges the markings of  $M$  along the leaf  $S^1$  of the covering  $\{1\}$  in order. Conclude that

$$\langle M, (so \text{ or } gl, N, \{1\}, n - 1) \rangle = nN^n + (\text{lower order terms}).$$

*Exercise 6.53.* The same is still true for the cover  $\{q\}$ . Only that in this case there are more liftings of the markings of  $M$  to the leaf of the cover that keep them in order. More precisely, let  $p$  be the number of non-decreasing sequences of length  $n - 1$  of non-negative integers smaller than  $q$ . Prove that

$$\begin{aligned} \langle M, (so \text{ or } gl, N, \{q\}, n - 1) \rangle &= qpN^n + \dots = q \binom{q + n - 2}{n - 1} N^n + \dots \\ &= \left( \frac{q^n}{(n - 1)!} + (\text{lower powers of } q) \right) N^n + \dots \end{aligned}$$

*Exercise 6.54.* Let  $M = \bigcup_1^b M_i$  be a disjoint union of  $b$  disks with  $n_i$  aligned tangents marked on the boundary of the  $i$ th disk,  $1 \leq i \leq b$ . Let  $m$  be the degree of  $M$ , let  $n = \sum n_i$  be the total number of markings on  $M$  and let  $L^\beta(N)$  be the labeled covering  $(so \text{ or } gl, N, \{1^\beta\}, m)$ . Prove that the highest power of  $N$  in  $\langle M, L^\beta(N) \rangle$  is  $N^{\beta - b + n}$ , and it is attained in the sum (39) only in the cases were the lifting  $l$  satisfies:

- For each  $i$ , all the markings on  $M_i$  lift to the same component of  $L^\beta$ , and the cyclic ordering induced on them from the cyclic ordering the points of that component agrees with their cyclic order on the boundary of  $M_i$ .
- If, for some  $i$  and  $j$ , the disks  $M_i$  and  $M_j$  lift to the same component  $L_0$  of  $L^\beta$ , then they are *non-interlaced* —  $L_0$  can be cut into two disjoint arcs  $A_i$  and  $A_j$ , so that  $A_i$  ( $A_j$ ) is connected only to  $M_i$  ( $M_j$ ).

**Definition 6.55.** The *width* of a monomial  $Q$  in the variables  $q_1, \dots, q_\beta$  is the number of the  $q_i$ 's that appear in  $Q$  in a positive power. For example, the width of the monomial  $x^2y^3z$  is 3. The *maximal width* of a polynomial  $P$  is the maximal width of a monomial in  $P$ .

*Exercise 6.56.* Prove that if  $M$  is as in the previous exercise and  $L^\beta(N)$  is the labeled covering  $(so \text{ or } gl, N, \{q_1, \dots, q_\beta\}, m)$ , then the same conclusion as in the previous exercise holds. Let  $Q$  be the coefficient of  $N^{\beta - b + n}$  in  $\langle M, L^\beta(N) \rangle$ . Prove that  $Q$  is a polynomial of maximal width  $\min(b, \beta)$  in  $q_1, \dots, q_\beta$ , and that if in addition it is given that  $\beta \geq b$  then

$$Q = \left( b! \sum_{1 \leq j_1 < \dots < j_b \leq \beta} \prod_{i=1}^b \frac{q_{j_i}^{n_i}}{(n_i - 1)!} \right) + \left( \begin{array}{c} \text{terms of lower width or} \\ \text{order in } q_1, \dots, q_\beta \end{array} \right). \quad (40)$$

Notice that the above exercise shows that if  $M$  is a connected normalized surface than its isomorphism class can be read from the numbers  $\langle M, L \rangle$  for various  $L$ . The degree of  $M$  can be read from the degree of any  $L$  for which  $\langle M, L \rangle \neq 0$ .  $M$  is *not* orientable iff  $\langle M, L \rangle = 0$  for all  $L$  of symbol  $gl$ ,  $b$  can be read by increasing  $\beta$  until the width of  $Q$  stops growing, and then the number of markings on each component of the boundary of  $M$  can be read from the powers of the  $q_j$ 's appearing in (40).

If  $M$  is a general normalized surface, we will show that its isomorphism class can be read from the numbers  $\langle M, L \rangle$  where  $L$  is a bouquet  $L = \cup L_k^{\beta_k}(N_k)$  where each  $L_k^{\beta_k}(N_k)$  is a labeled covering of the type considered in exercise 6.56:

$$L = \{(s_1, N_1, \{q_{11}, \dots, q_{1\beta_1}\}, m_1), \dots, (s_\gamma, N_\gamma, \{q_{\gamma 1}, \dots, q_{\gamma\beta_\gamma}\}, m_\gamma)\},$$

here each  $s_k$  is either  $so$  or  $gl$ .

*Exercise 6.57.* For fixed  $s_k$ 's and fixed  $m_k$ 's, show that  $P = \langle M, L \rangle$  is a polynomial in the variables  $N_1, \dots, N_\gamma, q_{11}, \dots, q_{\gamma\beta_\gamma}$  and that this polynomial is divisible by  $\prod_{k=1}^\gamma N_k$ .

We will now define an ordering  $\preceq$  among the monomials that appear in  $\langle M, L \rangle$  and show that from the  $\preceq$ -maximal monomial in  $\langle M, L \rangle$  the isomorphism class of  $M$  can be read.

**Definition 6.58.** Order monomials in the variables  $N_1, \dots, N_\gamma, q_{11}, \dots, q_{\gamma\beta_\gamma}$  which are divisible by  $\prod N_k$  lexicographically according to the following parameters of such a monomial  $Q$ :

- The total degree of  $Q$  in the variables  $N_1, \dots, N_\gamma$ .
- The width of  $Q / (\prod N_k)$  in the variables  $N_1, \dots, N_\gamma$ .
- The width of  $Q$  in the variables  $q_{11}, \dots, q_{\gamma\beta_\gamma}$ .
- The power of  $q_{11}$  in  $Q$ .
- The power of  $q_{12}$  in  $Q$ .
- ⋮
- The power of  $q_{\gamma\beta_\gamma}$  in  $Q$ .

*Exercise 6.59.* Let  $M$  be a normalized marked surface. For a fixed  $\gamma$ , set  $s_k = so$ , ( $1 \leq k \leq \gamma$ ). Find some values for  $m_1, \dots, m_\gamma$  for which  $\langle M, L \rangle$  is non-zero (show that this is possible whenever  $\gamma$  is sufficiently large), and call the resulting polynomial  $P_\gamma$ . Let  $c$  be the maximal value of  $\gamma$  for which the maximal width  $w_\gamma$  of  $P_\gamma / (\prod N_k)$  is still equal to  $\gamma$  (show that such a value of  $\gamma$  exists and that  $w_\gamma$  is constant for  $\gamma \geq c$ ). For that value of  $\gamma$ , fix  $\beta_1, \dots, \beta_\gamma$  for which the maximal width of  $P_\gamma$  in the variables  $q_{11}, \dots, q_{\gamma\beta_\gamma}$  has reached its maximal possible value (show that this 'settling of width' indeed happens). Let  $Q$  be a  $\preceq$ -maximal monomial in  $P_\gamma$ . Show that  $M$  is the union of  $c$  connected components (orientable or not)  $M_1, \dots, M_c$  satisfying:

- $\deg M_k = m_k$ .
- The width  $b_k$  of  $Q$  in the variables  $q_{k1}, \dots, q_{k\beta_k}$  is equal to the number of boundary components of  $M_k$ .
- Order the boundary components of  $M_k$  in some way  $B_{k1}, \dots, B_{kb_k}$  so that the number  $n_{kj}$  of markings on  $B_{kj}$  is a non-increasing function of  $j$ . Then  $n_{kj}$  is equal to the degree of  $q_{kj}$  in  $Q$ .

*Exercise 6.60.* For a fixed  $\gamma$  and  $m_1, \dots, m_\gamma$  as before, find a maximal set  $S$  of  $s_k$ 's that can be changed from *so* to *gl* so that the  $\preceq$ -order of the resulting  $Q$  remains the same as in the previous exercise (such a set might be empty). Show that the components  $M_k$  corresponding to the  $s_k$ 's in  $S$  are orientable.

The last two exercises show that the isomorphism class of  $M$  can be read from the numbers  $\langle M, L \rangle$ . We now just need to show that if  $\bar{M}$  is a linear combination of normalized marked surfaces and  $\langle \bar{M}, L \rangle = 0$  for all  $L$ 's, then  $\bar{M} = 0$ .

*Exercise 6.61.* Show that this is indeed the case.

*Hint 6.62.* Define an ordering relation  $\ll$  on normalized marked surfaces in a way similar to the definition of  $\preceq$ . Show that the coefficient of the  $\ll$ -maximal surface appearing in  $\bar{M}$  with a non-zero coefficient can be read from  $\langle \bar{M}, L \rangle$  using the previous two exercises.  $\square$

**6.5. Marked surfaces and the classical knot polynomials.** Using everything that we already know, the simplest way to prove theorem 12 is to recall (see Turaev [35, 37, 41]) that the HOMFLY polynomial is built from the Lie algebra  $sl(N)$  (or from the essentially equivalent (see exercise 6.34) Lie algebra  $gl(N)$ ), that the Kauffman polynomial is built from the Lie algebra  $so(N)$ , and that cabling these polynomials corresponds to taking higher representations of those groups. Remembering all that, theorem 12 follows immediately from theorem 11. Alternatively, if one does not want to use theorem 5, it is easy to get equations (34) and (36) from the power series expansion in the variable  $x$  of a framing dependent version of the HOMFLY polynomial,

$$P^f \left( \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) - P^f \left( \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \right) = (e^{x/2} - e^{-x/2}) P^f \left( \begin{array}{c} \diagup \\ \diagdown \end{array} \right) \hat{\circ} \left( \begin{array}{c} \diagdown \\ \diagup \end{array} \right);$$

$$P^f \left( \begin{array}{c} \uparrow \\ \circ \end{array} \right) = e^{Nx/2} P^f \left( \begin{array}{c} \uparrow \\ \uparrow \end{array} \right),$$

and to get equations (29) and (33) from the power series expansion in the variable  $x$  of a framing dependent version of the Kauffman polynomial,

$$F^f \left( \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) - F^f \left( \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \right) = (e^{x/2} - e^{-x/2}) (F^f \left( \begin{array}{c} \diagup \\ \diagdown \end{array} \right) \hat{\circ} \left( \begin{array}{c} \diagdown \\ \diagup \end{array} \right) - F^f \left( \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \right));$$

$$F^f \left( \begin{array}{c} \uparrow \\ \circ \end{array} \right) = e^{\frac{N-1}{2}x} F^f \left( \begin{array}{c} \uparrow \\ \uparrow \end{array} \right). \quad \square$$

**6.6. More on conjecture 2.** Recalling that  $\Phi \circ \bar{\sigma} = \tau \circ \mu \circ \bar{\sigma}$  and that  $\bar{\sigma}$  and  $\tau$  are isomorphisms (theorems 8 and 14), we see that conjecture 2 is true iff the map  $\mu$  is one-to-one iff its adjoint  $\mu^*$  is onto. Keeping in mind theorem 12, we see that conjecture 2 is essentially the assertion:

- Vassiliev invariants are precisely as powerful as the HOMFLY and Kauffman polynomials and all of their cablings.

Whether or not this assertion is true, asking whether  $\mu$  is 1-1 is clearly a simpler way of stating it, perhaps making it easier to resolve.

I have the following evidence to support conjecture 2:

- It had been verified up to degree 9.

- Its natural generalization had been verified for *homotopy link invariants*<sup>15</sup> (see [8]). A homotopy link invariant is a link invariant that does not change when an overcrossing in which only one component of the link is involved is replaced by an undercrossing.
- Conjecture 2 is similar in form to the question of whether any weight system can be integrated to a knot invariant, which, after some hard work in section 4, was answered affirmatively. In both cases the question is whether a certain map is onto; in the case of conjecture 2 the map is the adjoint of

$$(\text{-----}) \rightarrow (\text{-----}) - (\text{-----} \times \text{-----}), \quad (41)$$

while in the case of section 4 the map is the adjoint of

$$\times \rightarrow \times - \times.$$

- To prove conjecture 2 one needs to show that it is possible to extend any functional  $\phi$  defined on  $\mathcal{B}$  to a functional (also called  $\phi$ ) defined on the space  $\mathcal{B}^{pm}$  of partially marked diagrams — diagrams in which only a part of the arcs are marked — so that the extended  $\phi$  maps (41) to an equality. It is not hard to show that over  $\mathbf{Z}/2\mathbf{Z}$  it is possible to consistently extend every functional defined on unmarked diagrams to diagrams marked exactly once.

## 7. ODDS AND ENDS

### 7.1. Some questions.

7.1.1. *Why Lie algebras?* In some sense (theorem 1) studying the algebra  $\mathcal{A}$  is *exactly* the same as studying Vassiliev invariants, and indeed, some of the structure of  $\mathcal{A}$  can be understood in terms of knot theory (remark 3.2, exercises 3.6, 3.10, 3.14). The algebra  $\mathcal{A}$  (or, in fact,  $\mathcal{A}^*$ ) also has a (*weaker*) relation with Lie algebras (theorem 4 and conjecture 1), a relation which is not 1-1 (exercises 6.28 and 6.35) and not known to be onto. However, many of the results about  $\mathcal{A}$  have a Lie theoretic interpretation but seem to have no knot theoretic interpretation.

**Problem 7.1.** Interpret theorems 6 and 8 and remark 5.7 in terms of knot theory.

7.1.2. *Why surfaces.* A Short glance at the diagram in section 1.9 shows that to every marked surface naturally corresponds a knot invariant, and that the knot invariants thus obtained are rather strong.

**Problem 7.2.** Understand why do marked surfaces appear in knot theory. Find a direct topological construction for the invariant corresponding to a marked surface.

7.1.3. *The ground ring.*

**Problem 7.3.** How much of the theory of sections 4, 5 and 6 carries over to an arbitrary ground ring? A computer search (*e.g.*  $\dim \mathcal{G}_m \mathcal{A}$  was re-computed over fields of small prime characteristics) suggests that perhaps everything carries over.

---

<sup>15</sup>I wish to thank C. Day for teaching me about this class of invariants.

7.1.4. *Higher products.* On  $\mathcal{B}$  (and therefore also on  $\mathcal{A}$ ) there is a second grading: by the number of external vertices in a Chinese character. Furthermore,  $\mathcal{B}$  has various *higher products*; for a positive integer  $j$  and a pair of Chinese characters  $C_1, C_2$  define  $m_j(C_1, C_2)$  to be the sum over all possible ways of sewing  $C_1$  and  $C_2$  along  $j$  external vertices:

$$m_1 \left( \begin{array}{c} \circ \\ \circ \end{array}, \begin{array}{c} \circ \\ \circ \end{array} \right) = 4 \begin{array}{c} \circ \\ \circ \end{array}; \quad m_2 \left( \begin{array}{c} \circ \\ \circ \end{array}, \begin{array}{c} \square \\ \square \end{array} \right) = 8 \begin{array}{c} \circ \\ \square \\ \circ \end{array} + 4 \begin{array}{c} \square \\ \square \end{array}.$$

Superficially,  $m_1$  and the product of  $\mathcal{A}$  are defined in similar ways. However, there appears to be no obvious relation between the two — their images are not even in the same degree!

**Problem 7.4.** Understand the second grading of  $\mathcal{A}$  and the higher products  $m_j$  in terms of knot theory. Investigate the relations between the  $m_j$  and the other structures introduced in this paper.

7.2. **Some bad news.** The algebra  $\mathcal{A}$  has a natural involution — simply map every chord diagram  $D \in \mathcal{D}^c$  into the same diagram, only with the orientation of its circle reversed. In the language definition 3.11, this is just the operation  $\psi^{-1}$ , and therefore by exercise 3.14 it corresponds to the operation of orientation reversal for knots (reversal of the orientation of the knot itself, not of the ambient space!). Being an involution,  $\psi^{-1}$  ( $(\psi^{-1})^*$ ) decomposes  $\mathcal{A}$  ( $\mathcal{A}^*$ ) into a sum of a  $+1$  eigenspace and a  $-1$  eigenspace. Call a diagram  $D \in \mathcal{A}$  (a functional  $W \in \mathcal{A}^*$ ) *even* if it satisfies  $\psi^{-1}D = D$  ( $(\psi^{-1})^*W = W$ ) in  $\mathcal{A}$  ( $\mathcal{A}^*$ ), and call it *odd* if it satisfies  $\psi^{-1}D = -D$  ( $(\psi^{-1})^*W = -W$ ).

**Problem 7.5.** Are there any odd diagrams? (This question was first raised by J. Birman)

A computer search has shown that all diagrams are even up to degree 9. This is rather disturbing because it means that all Vassiliev invariants up to degree 9 are even, *i.e.*, cannot tell a knot from its *inverse* (its orientation reversed version). Even worse than that is true — if conjecture 2 or even conjecture 1 is true, then *all* Vassiliev invariants are even, contradicting the hope implicitly expressed in problem 1.1:

*Exercise 7.6.* Show that via the isomorphism  $\mathcal{A} \leftrightarrow \mathcal{B}$ , an even (odd) diagram is carried to a Chinese character with an even (odd) number of external vertices.

*Exercise 7.7.* Without using theorems 11 or 12, show that if  $C \in \mathcal{B}$  is a Chinese character with an odd number of external vertices, then  $\Phi(C) = 0$ .

*Exercise 7.8.* Show that if  $\mathfrak{g}$  is a semi-simple Lie algebra and  $R$  is an arbitrary representation of  $\mathfrak{g}$ , then  $W_{\mathfrak{g}, R, m}$  is even for every  $m$ .

*Hint 7.9.* Use exercise 6.32 and the fact that every Lie algebra over the complex numbers has a conjugate-linear involution (*The Cartan involution*) that carries every representation to its complex conjugate.

7.3. **Bibliographical remarks.** In addition to the works mentioned in the body of this paper, the following papers also discuss Vassiliev invariants: J. C. Baez [4] (Vassiliev invariants for braids, relations with Chern-Simons theory and with quantum gravity) C. Day [14] (Vassiliev invariants of links), M. Gusarov [18], A. B. Sossinsky [38], T. Stanford [39] (Vassiliev invariants of links and graphs), and T. Stanford [40] (Vassiliev invariants for braids, examples for knots that cannot be separated apart by Vassiliev invariants of a fixed type).

## REFERENCES

- [1] V. I. Arnold, *The cohomology ring of the group of dyed braids* (Russian), *Mat. Zametki* **5** (1969) 227–231.
- [2] M. F. Atiyah and D. O. Tall, *Group representations,  $\lambda$ -rings and the  $J$ -homomorphism*, *Topology* **8** (1969), 253–297.
- [3] S. Axelrod and I. M. Singer, *Chern-Simons perturbation theory*, M.I.T. preprint, October 1991.
- [4] J. C. Baez, *Link invariants of finite type and perturbation theory*, Wellesley college preprint, July 1992.
- [5] D. Bar-Natan, *Weights of Feynman diagrams and the Vassiliev knot invariants*, February 1991, Preprint.
- [6] ———, *Perturbative aspects of the Chern-Simons topological quantum field theory*, Ph.D. thesis, Princeton Univ., June 1991, Dep. of Mathematics.
- [7] ———, *Computer data files*, <http://www.ma.huji.ac.il/~drorbn/papers/OnVassiliev/>.
- [8] ———, *Vassiliev Homotopy String Link Invariants*, *Jour. of Knot Theory and its Ramifications*, to appear.
- [9] ———, *To Appear*.
- [10] J. S. Birman and X-S. Lin, *Knot polynomials and Vassiliev’s invariants*, Columbia Univ. preprint, 1991.
- [11] T. Bröcker and T. tom Dieck, *Representations of compact Lie groups*, Springer-Verlag GTM **98**, New-York 1985.
- [12] K-T. Chen, *Iterated path integrals*, *Bull. Amer. Math. Soc.* **83** (1977) 831–879.
- [13] J. H. Conway, *An enumeration of knots and links and some of their algebraic properties*, in *Computational Problems in Abstract Algebra*, 329–358, Pergamon, New-York 1970.
- [14] C. Day, *Vassiliev invariants for links*, Univ. of North Carolina at Chapel Hill preprint, 1992.
- [15] J. Dixmier, *Enveloping algebras*, North-Holland, Amsterdam 1977.
- [16] V. G. Drinfel’d, *Quasi-Hopf algebras*, *Leningrad Math. J.* **1** (1990) 1419–1457.
- [17] ———, *On quasitriangular Quasi-Hopf algebras and a group closely connected with  $Gal(\bar{\mathbb{Q}}/\mathbb{Q})$* , *Leningrad Math. J.* **2** (1991) 829–860.
- [18] M. Gusarov, *A new form of the Conway-Jones polynomial of oriented links*, *Nauchn. Sem. Len. Otdel. Mat. Inst. Steklov* **193** (1991), 4–9.
- [19] J. Hoste, A. Ocneanu, K. Millett, P. Freyd, W. B. R. Lickorish and D. Yetter, *A new polynomial invariant of knots and links*, *Bull. Amer. Math. Soc.* **12** (1985) 239-246.
- [20] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, *Bull. Amer. Math. Soc.* **12** (1985) 103–111.
- [21] L. H. Kauffman, *On knots*, Princeton Univ. Press, Princeton, 1987.
- [22] ———, *An invariant of regular isotopy*, *Trans. Amer. Math. Soc.* **312** (1990) 417–471.
- [23] V. G. Knizhnik and A. B. Zamolodchikov, *Current algebra and Wess-Zumino model in two dimensions*, *Nucl. Phys.* **B247** (1984) 83–103.
- [24] T. Kohno, *Monodromy representations of braid groups and Yang-Baxter equations*, *Ann. Inst. Fourier* **37** (1987) 139–160.
- [25] M. Kontsevich, *Private communication*.
- [26] ———, *Graphs, homotopical algebra and low-dimensional topology*, preprint, 1992.
- [27] G. Kuperberg, *The quantum  $G_2$  link invariant*, Univ. of California at Berkeley preprint, August 1990 (to appear in *Int. Jour. of Math.*).
- [28] X-S. Lin, *Vertex models, quantum groups and Vassiliev’s knot invariants*, Columbia Univ. preprint, 1991.
- [29] ———, *Private communication*.
- [30] W. S. Massey, *A basic course in algebraic topology*, Springer-Verlag GTM **127**, New-York 1991.
- [31] J. Milnor and J. Moore, *On the structure of Hopf algebras*, *Annals of Math.* **81** (1965) 211–264.
- [32] R. C. Penner, *The decorated Teichmüller space of punctured surfaces*, *Commun. Math. Phys.* **113** (1987) 299–339.
- [33] S. Piunikhin, *Weights of Feynman diagrams, link polynomials and Vassiliev knot invariants*, Moscow State Univ. preprint, 1992.
- [34] J. H. Przytycki, *Talk given at the Georgia topology conference*, August 1992.
- [35] N. Yu. Reshetikhin, *Quantized universal enveloping algebras, the Yang-Baxter equation and invariants of links (I & II)*, LOMI preprints **E-4-87** & **E-17-87**, Leningrad 1988.

- [36] ———, *Quasitriangle Hopf algebras and invariants of tangles*, Leningrad Math. J. **1** (1990) 491–513.
- [37] ——— and V. G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, Commun. Math. Phys. **127** (1990) 1–26.
- [38] A. B. Sossinsky, *Feynman diagrams and Vassiliev invariants*, IHES preprint **IHES/M/92/13**, February 1992.
- [39] T. Stanford, *Finite type invariants of knots, links, and graphs*, Columbia Univ. preprint, July 1992.
- [40] ———, *Braid commutators and Vassiliev invariants*, Columbia Univ. preprint, August 1992.
- [41] V. G. Turaev, *The Yang-Baxter equation and invariants of links*, Invent. Math. **92** (1988) 527–553.
- [42] V. A. Vassiliev, *Cohomology of knot spaces*, Theory of Singularities and its Applications (Providence) (V. I. Arnold, ed.), Amer. Math. Soc., Providence, 1990.
- [43] V. A. Vassiliev, *Complements of discriminants of smooth maps: topology and applications*, Trans. of Math. Mono. **98**, Amer. Math. Soc., Providence, 1992.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138  
*E-mail address:* drorbn@math.toronto.edu