

Bihamiltonian Structures and Quadratic Algebras in Hydrodynamics and on Non-Commutative Torus

B. Khesin^{1,2}, A. Levin^{1,3}, M. Olshanetsky^{1,4}

¹ Max Planck Institute of Mathematics, Bonn, Germany

² University of Toronto, Toronto, Canada. E-mail: khesin@math.toronto.edu

³ Institute of Oceanology, Moscow, Russia. E-mail: alevin@wave.sio.rssi.ru

⁴ Institute of Theoretical and Experimental Physics, Moscow, Russia. E-mail: olshanet@gate.itep.ru

Received: 4 September 2003 / Accepted: 12 February 2004

Published online: 12 August 2004 – © Springer-Verlag 2004

Abstract: We demonstrate the common bihamiltonian nature of several integrable systems. The first one is an elliptic rotator that is an integrable Euler-Arnold top on the complex group $GL(N, \mathbb{C})$ for any N , whose inertia ellipsoid is related to a choice of an elliptic curve. Its bihamiltonian structure is provided by the compatible linear and quadratic Poisson brackets, both of which are governed by the Belavin-Drinfeld classical elliptic r -matrix. We also generalize this bihamiltonian construction of integrable Euler-Arnold tops to several infinite-dimensional groups, appearing as certain large N limits of $GL(N, \mathbb{C})$. These are the group of a non-commutative torus (NCT) and the group of symplectomorphisms $SDiff(T^2)$ of the two-dimensional torus. The elliptic rotator on symplectomorphisms gives an elliptic version of an ideal 2D hydrodynamics, which turns out to be an integrable system. In particular, we define the quadratic Poisson algebra on the space of Hamiltonians on T^2 depending on two irrational numbers. In conclusion, we quantize the infinite-dimensional quadratic Poisson algebra in a fashion similar to the corresponding finite-dimensional case.

Contents

1. Introduction	582
2. Main Results	583
3. $GL(N, \mathbb{C})$ -Elliptic Rotators	586
3.1. Lax representation and integrals of motion.	586
3.2. The classical $GL(N, \mathbb{C})$ Feigin-Odesski algebras.	590
3.3. The bihamiltonian structure.	591
4. Elliptic Rotators on Non-Commutative Torus	593
4.1. The Lax equation and integrals of motion.	593
4.2. Quadratic Poisson algebras and the bihamiltonian structure on NCT.	596
5. Elliptic Rotators on $SDiff(T^2)$	597
5.1. Description of the hierarchy.	597

5.2. Quadratic Poisson algebras, and the bihamiltonian structure on $SDiff(T^2)$	600
5.3. Rational limit of the elliptic hydrodynamics.	600
6. Quantum Counterparts	602
Appendices	605
Appendix A. Elliptic functions	605
Appendix B. Elliptic constants related to $gl(N, \mathbb{C})$	608
Appendix C. Non-commutative torus	609

1. Introduction

One of the most interesting problems in the theory of integrable systems is a description of infinite-dimensional integrable systems with two or more space variables. In this paper we consider integrable hierarchies in the $(2 + 1)$ dimension, which we call *elliptic* and *modified hydrodynamics*. These are close relatives of the ideal hydrodynamics on a two-dimensional torus, which is known to be a highly non-integrable system. For instance, even the 4-vortex approximation of the ideal fluid dynamics is non-integrable [1]. We obtain these integrable versions of hydrodynamics starting from finite-dimensional integrable systems whose dimension of the phase space goes to infinity. Our approach is somewhat similar to the derivation of Toda field theory from the open Toda chain. In contrast with the Toda theory, however, we obtain non-local systems.

The classical Euler (or, rather, Helmholtz) equation for the motion of an ideal fluid on the standard two-torus is

$$\partial_t \mathbf{S} = \{\mathbf{S}, \Delta^{-1} \mathbf{S}\},$$

where \mathbf{S} is the vorticity function of the fluid flow, $\{ \}$ is the Poisson bracket, and Δ is the Laplace operator on T^2 . The modified and elliptic hydrodynamics are defined on an elliptic curve, i.e., on a torus with a complex structure fixed. The corresponding equations are, respectively,

$$\partial_t \mathbf{S} = \{\mathbf{S}, \bar{\partial}^{-2} \mathbf{S}\}$$

and

$$\partial_t \mathbf{S} = \{\mathbf{S}, \wp(\bar{\partial}) \mathbf{S}\},$$

where $\bar{\partial}$ is the corresponding operator of the complex structure, and \wp is the Weierstrass \wp -function. (We postpone the precise description of $\wp(\bar{\partial})$ till Sect. 5.) Note that to define the Laplace operator one needs to choose a metric, while for the operator $\bar{\partial}$ in the modified hydrodynamics is defined by a complex structure on T^2 .

Such a modification of the fluid inertia operator from the Laplace operator $\Delta = \partial \bar{\partial}$, which depends on a metric on T^2 , to $\bar{\partial}^2$ or $\wp(\bar{\partial})^{-1}$, both of which depend on a complex structure on T^2 , brings in the integrability and even the bihamiltonian structure for the systems. We construct an infinite set of involutive integrals of motion with respect to two Poisson brackets. One of the brackets is the standard linear Lie-Poisson brackets on the dual space to the algebra of the divergence-free vector fields on T^2 . The other Poisson structure is a quadratic Poisson algebra on Hamiltonians of vector fields. These two brackets are compatible and governed by the same classical r -matrix. Furthermore, we describe a recursion procedure for constructing the sequence of Hamiltonians for this linear-quadratic bihamiltonian structure of the hierarchy of the elliptic hydrodynamics, which thereby exhibits “the strongest form of integrability.”

We come to this construction through the non-commutative deformation of T^2 to the non-commutative torus (NCT). The non-commutative deformation of the Lie algebra of

vector fields is isomorphic to the Lie algebra of NCT. The latter is a special large N limit of $\mathfrak{gl}(N, \mathbb{C})$. We start with the elliptic rotators on $\mathrm{GL}(N, \mathbb{C})$ [2, 3] and develop their bihamiltonian structure based on the Belavin-Drinfeld classical r -matrix [4]. The corresponding quadratic Poisson algebra is the classical limit of the Feigin-Odesski algebra [5]. We describe natural extensions of the elliptic rotators to the infinite-dimensional groups, which preserve their main properties. This brings us to bihamiltonian systems and integrable hierarchies on the NCT, which are of interest by themselves.

As a byproduct, we also describe a quantization of the quadratic Poisson algebra on NCT. In this way we obtain an infinite-dimensional associative algebra with quadratic relations depending on the complex structure on T^2 and the Planck constant $\hbar \in T^2$.

Finally, the commutative (“dispersionless”) limit of the elliptic rotators on NCT leads to the desirable bihamiltonian hierarchies of the elliptic and modified hydrodynamics.

2. Main Results

1. The general setup [6, 7]. Let G be a Lie group and \mathfrak{g} its Lie algebra. Consider an invertible linear operator \mathbf{J} that maps the coalgebra \mathfrak{g}^* to \mathfrak{g} . Its inverse operator \mathbf{J}^{-1} is called *the inertia tensor*. The Euler-Arnold top corresponding to the group G is the Hamiltonian system on \mathfrak{g}^* with respect to the linear Lie-Poisson brackets on \mathfrak{g}^* and the Hamiltonian function given by the quadratic form

$$H = -\frac{1}{2} \langle \mathbf{S}, \mathbf{J}(\mathbf{S}) \rangle, \quad \mathbf{S} \in \mathfrak{g}^*,$$

where $\langle \cdot, \cdot \rangle$ stands for the pairing between \mathfrak{g} and \mathfrak{g}^* . Namely, the variation ∇H can be regarded as an element of \mathfrak{g} and the corresponding Hamiltonian equation of motion is as follows:

$$\partial_t \mathbf{S} = \{H, \mathbf{S}\} := \mathrm{ad}_{\nabla H}^* \mathbf{S}.$$

Recall that the Lie-Poisson brackets are degenerate on \mathfrak{g}^* and their symplectic leaves are coadjoint orbits of G . To descend to a particular coadjoint orbit \mathcal{O} one should fix the values of Casimirs for the linear bracket.

For some special choices of \mathbf{J} the system becomes completely integrable. Some of the examples are Manakov’s tops on $SO(N)$ [8], their limit $N \rightarrow \infty$ found by Ward [9], the Korteweg–de Vries equation [10, 11], as well as the Camassa–Holm and Hunter–Saxton equations on the Virasoro group [12]. (In the infinite-dimensional case, the invertibility of \mathbf{J} is understood as $\mathrm{Ker} \mathbf{J} = 0$.)

2. Elliptic Rotators on $\mathrm{GL}(N, \mathbb{C})$. The elliptic rotator (ER) on $\mathrm{GL}(N, \mathbb{C})$ is an integrable Euler-Arnold top on this group, whose inertia operator is constructed with the help of the Weierstrass \wp -function of an auxiliary elliptic curve, see [2, 3, 13]. Namely, let T_α , $\alpha = (\alpha_1, \alpha_2)$ be the basis of $\mathfrak{sl}(N, \mathbb{C})$ (B.4.) $\alpha_j = (0, 1, \dots, N-1)$, $\alpha \neq (0, 0)$. The structure constants in this basis are

$$\mathbf{C}_\theta(\alpha, \beta) = \frac{1}{\pi\theta} \sin \pi\theta(\alpha \times \beta), \quad \theta = k/N,$$

where $1 \leq k < N$ and k, N are coprime, see (B.8). For $\mathbf{S} = \sum_\alpha S_{-\alpha} T_\alpha \in \mathfrak{sl}(N, \mathbb{C})^*$ the linear Poisson brackets assume the form

$$\{S_\alpha, S_\beta\} = \mathbf{C}_\theta(\alpha, \beta) S_{\alpha+\beta}. \quad (2.1)$$

Let $\wp(x; \tau)$ be the Weierstrass function on the elliptic curve $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$. Consider its values $\wp_\theta(\alpha) = \wp((\alpha_1 + \alpha_2\tau)\theta; \tau)$ on the lattice parametrized by α , the labels of the $\mathfrak{sl}(N, \mathbb{C})$ -basis. The inverse inertia operator is defined as

$$\mathbf{J} : S_\alpha \rightarrow J_\alpha S_\alpha, \quad J_\alpha = \wp_\theta(\alpha). \quad (2.2)$$

It was proved in Ref.[2] that the equations of motion

$$\partial_t S_\alpha = \sum_{\gamma} S_{\alpha-\gamma} \wp_\theta(\gamma) S_\gamma, \quad (2.3)$$

defined by the Hamiltonian $H = -\frac{1}{2}\langle \mathbf{S}, \mathbf{J}(\mathbf{S}) \rangle = 2\pi^2\theta^2 \text{tr}(\mathbf{S} \cdot \mathbf{J}(\mathbf{S}))$ and the brackets (2.1), have the Lax representation with the Lax operator $L^{rot}(\mathbf{S}, z) \in \mathfrak{sl}(N, \mathbb{C})$ depending on the spectral parameter $z \in E_\tau$. The traces $\text{tr}(L^{rot}(\mathbf{S}, z))^k$, $k = 2, \dots, N$ being expanded in the basis of elliptic functions on E_τ produce the involutive integrals of motion and the Hamiltonian H is among them.

We call the commuting flows defined by these integrals *the ER hierarchy*.

Let $r(z)$, $z \in E_\tau$ be the Belavin-Drinfeld classical elliptic r -matrix, see [13, 4] and Sect. 3 below. The main result, which we will also generalize to the infinite-dimensional situation, can be formulated as follows.

Theorem 2.1. *i) In terms of the r -matrix, the linear brackets (2.1) can be written in the form*

$$\{L_1^{rot}(z), L_2^{rot}(w)\}_1 = [r(z-w), L_1^{rot}(z) + L_2^{rot}(w)]. \quad (2.4)$$

ii) Consider the phase space extended by a new variable S_0 , which has zero bracket with all the rest. The same r -matrix defines the quadratic Poisson algebra $P_{N,\theta,\tau}$ related to $\text{GL}(N, \mathbb{C})$ (3.19), (3.20):

$$\begin{aligned} \{L_1(z), L_2(w)\}_2 &= [r(z-w), L_1(z) \otimes L_2(w)], \quad \text{where } L(z) \\ &= S_0 Id + L^{rot}(z). \end{aligned} \quad (2.5)$$

iii) The above two brackets are compatible, i.e. any linear combination of them is a Poisson bracket.

iv) There exists a sequence of integrals of motion in involution with respect to each of the two brackets $\{h_j, h_k\}_{1,2} = 0$ (here the lower indices refer to the linear and the quadratic brackets respectively). They provide the bihamiltonian structure of the elliptic rotator (ER) hierarchy

$$\{h_{j+1}, \mathbf{S}\}_2 = -\{h_j, \mathbf{S}\}_1.$$

The first two statements of the theorem for $\text{GL}(N, \mathbb{C})$ are well known [5, 14]. Apparently, the bihamiltonian structure of the GL_N elliptic rotators is new, and it gives the following

Corollary 2.1. *The Casimirs with respect to one of the brackets generate an integrable dynamics with respect to the other.*

In particular, the functional S_0 , being the Casimir element of the linear brackets (2.1), leads to the equations

$$\partial_t \mathbf{S} = \{S_0, \mathbf{S}\}_2, \quad (2.6)$$

that coincide with (2.3).

It turns out that Theorem 2.1 also holds in the infinite-dimensional situation presented below.

3. *Elliptic Rotators on the Non-commutative Torus.* Consider the \sin -algebra \sin_θ , called also the Lie algebra of the non-commutative torus (NCT). The Poisson brackets on the Lie coalgebra \sin_θ^* of the NCT has the form (2.1) with irrational number $0 \leq \theta < 1$ and the basis $T_\alpha = \text{const} \cdot \exp(2\pi i \alpha_1 x_1) * \exp(2\pi i \alpha_2 x_2)$ is parameterized by the infinite lattice $\alpha \in \mathbb{Z} \oplus \mathbb{Z}$. Here $\exp(2\pi i x_1)$, $\exp(2\pi i x_2)$ are the generators of the NCT and $*$ is the Moyal multiplication (see Appendix C). On the NCT we introduce the complex structure $\bar{\partial}_{\epsilon, \tau}$ depending on τ ($\text{Im } \tau > 0$), and two real numbers $\epsilon = (\epsilon_1, \epsilon_2)$, such that $\theta \epsilon_a$ are irrational and $0 < \theta \epsilon_a \leq 1$ (see (4.3)). The inverse inertia operator \mathbf{J} is the pseudo-differential operator

$$\mathbf{J}(\mathbf{S})(x) = \wp(\theta \bar{\partial}_{\epsilon, \tau}) \mathbf{S}(x),$$

where now $\mathbf{S}(x) = \sum_\alpha S_{-\alpha} T_\alpha(x)$. The equation of motion in the form of the Moyal brackets has the form

$$\partial_t \mathbf{S} = \{\mathbf{S}, \wp(\theta \bar{\partial}_{\epsilon, \tau}) \mathbf{S}\}^\theta, \quad (2.7)$$

where $\{\cdot\}^\theta$ is the Lie–Poisson bracket on \sin_θ^* . In Proposition 3.1 we find the Lax form of (2.7). We construct the classical r -matrix (4.11) and prove the counterpart of the theorem. The quadratic Poisson algebra $P_{\theta, \epsilon, \tau}$ (4.19), (4.20) on the NCT gives rise to the infinite bihamiltonian hierarchy, where Eqs. (2.7) correspond to the quadratic Hamiltonian functional with respect to the linear brackets. At the same time Eqs. (2.7) can be interpreted as the Hamiltonian equations (2.6) in the quadratic Poisson algebra $P_{\theta, \epsilon, \tau}$ with a linear Hamiltonian. For $\text{SL}(2)$ this representation was found in [15] and for the general case in [18].

4. *Elliptic Rotators on $S\text{Diff}(T^2)$.* In the limit $\theta \rightarrow 0$ the Lie algebra of the NCT becomes isomorphic to the Poisson algebra \mathcal{A} of smooth functions on T^2 modulo constants $\text{Ham}(T^2) \sim C^\infty(T^2)/\mathbb{C}$. The algebra of Hamiltonians can also be described by the corresponding Hamiltonian (or, divergence-free) vector fields on the torus. More precisely, the Lie algebra $S\text{Vect}(T^2)$ of divergence-free vector fields on T^2 is the (universal) cocentral extension of $\text{Ham}(T^2)$, defined by the exact sequence

$$0 \rightarrow \text{Ham}(T^2) \rightarrow S\text{Vect}(T^2) \rightarrow \mathbb{C}^2 \rightarrow 0,$$

where the image of $S\text{Vect}(T^2)$ in \mathbb{C}^2 is generated by the two fluxes $\epsilon_1 \partial_1$, $\epsilon_2 \partial_2$. For $\psi \in \text{Ham}(T^2)$ we have

$$V_1(\psi) = -\frac{1}{4\pi^2} \partial_2 \psi, \quad V_2(\psi) = \frac{1}{4\pi^2} \partial_1 \psi,$$

and $V_{\{\psi, \psi'\}} = [V(\psi), V(\psi)']$. We construct the elliptic rotator on the Lie group $S\text{Diff}(T^2)$ of area-preserving diffeomorphisms, corresponding to the Lie algebra $S\text{Vect}(T^2)$.

Consider the dual space of linear functionals $\text{Ham}(T^2)^*$ in the Fourier basis

$$\{\mathbf{e}(\alpha \cdot x) := \exp 2\pi i (\alpha \cdot x) \mid (x = (x_1, x_2), (\alpha \cdot x) = \alpha_1 x_1 + \alpha_2 x_2, \alpha_j \in \mathbb{Z}), \\ \mathbf{S} = \sum_\alpha S_\alpha \mathbf{e}(-(\alpha \cdot x)) \in \text{Ham}(T^2)^*.$$

The Poisson structure on \mathcal{A}^* assumes the form (cf. (2.1))

$$\{S_\alpha, S_\beta\}_1 = (\alpha \times \beta) S_{\alpha+\beta}.$$

This Poisson structure is degenerate and has an infinite set of Casimirs

$$C_k = \int_{T^2} \mathbf{S}^k, \text{ where } k = 2, 3, \dots$$

On a coadjoint orbit $\mathcal{O} \subset Ham(T^2)^*$ of $SDiff(T^2)$ the brackets are non-degenerate and values of the Casimirs are fixed.

Define the operator \mathbf{J} (after the rescaling) as the pseudo-differential operator $\wp(\bar{\partial}_{\epsilon, \tau})$, such that the Hamiltonian has the form

$$H = -\frac{1}{2} \int_{T^2} \mathbf{S} \cdot \wp(\bar{\partial}_{\epsilon, \tau}) \mathbf{S}. \quad (2.8)$$

We prove that the corresponding Hamiltonian system admits an infinite set of commuting integrals, see Sect. 5.

Recall that these elliptic rotators are parameterized, in particular, by the auxiliary elliptic curve. Consider the simplest version of the ER, corresponding to the rational degeneration of the elliptic curve E_τ . In the limit the (inverse) inertia operator becomes $\mathbf{J} = \bar{\partial}_\tau^{-2}$ and the limiting Hamiltonian (2.8) takes the form

$$H = -\frac{1}{2} \int_{T^2} \mathbf{S} \cdot \bar{\partial}_\tau^{-2} \mathbf{S},$$

see (5.23) and [15].

It is interesting to compare this with the Euler equation for an ideal fluid on a torus, where the inertia operator is the Laplacian (i.e., $\mathbf{J} = \Delta^{-1} = (\bar{\partial}\partial)^{-1}$), and the Hamiltonian is

$$H = -\frac{1}{2} \int_{T^2} \mathbf{S} \cdot \Delta^{-1} \mathbf{S},$$

where \mathbf{S} plays the role of the vorticity [7]. While the Euler hydrodynamics equation is highly non-integrable, the modification of the (inverse) inertia operator from $\bar{\partial}\partial$ to $\bar{\partial}_\tau^2$ leads to an integrable hierarchy. We call the rational limit *the modified hydrodynamics* on T^2 . (In a sense, these systems are similar in spirit to the Etingof–Frenkel current algebras [16] on T^2 , which make use of the complex structure on the elliptic curve to construct a central extension of smooth currents on the torus.)

Finally, we prove Theorem 2.1 for the generic systems of *the elliptic hydrodynamics* (2.8). In particular, the quadratic Poisson algebra $\mathcal{P}_{0, \tau}$ (5.27), (5.28) on T^2 , along with the linear Poisson bracket, provide the bihamiltonian structure of the hierarchies of the elliptic and modified hydrodynamics.

3. $GL(N, \mathbb{C})$ -Elliptic Rotators

3.1. Lax representation and integrals of motion.

1. The system description. The elliptic $GL(N, \mathbb{C})$ -rotator is defined on the dual spaces of $\mathfrak{gl}(N, \mathbb{C})$ and $\mathfrak{sl}(N, \mathbb{C})$. Let

$$\mathbf{S} = \sum_{\alpha \in \mathbb{Z}_N^{(2)}} S_{-\alpha} T_\alpha,$$

where T_α is the basis of $\mathfrak{sl}(N, \mathbb{C})$, see (B.4) and (B.5). In what follows we assume that the Greek indices belong to the lattice $\tilde{\mathbb{Z}}_N^{(2)}$ (B.5). Then the Poisson structure on the dual space $\mathfrak{sl}(N, \mathbb{C})^*$ is given by the linear Lie-Poisson brackets (B.8)

$$\{S_\alpha, S_\beta\}_1 = \mathbf{C}_\theta(\alpha, \beta) S_{\alpha+\beta}, \quad (3.1)$$

where $\mathbf{C}_\theta(\alpha, \beta)$ are the structure constants of $\mathfrak{sl}(N, \mathbb{C})$ (B.8). The Hamiltonian has the form

$$H^{rot} = 2\pi^2 \theta^2 \text{tr}(\mathbf{S} \cdot \mathbf{J}(\mathbf{S})) \equiv -\frac{1}{2} \sum_{\gamma \in \tilde{\mathbb{Z}}_N^{(2)}} S_\gamma J_\gamma S_{-\gamma}. \quad (3.2)$$

It defines the equations of motion

$$\partial_t \mathbf{S} = [\mathbf{J}(\mathbf{S}), \mathbf{S}]. \quad (3.3)$$

Let $\mathbf{J} : S_\alpha \rightarrow J_\alpha S_\alpha$, be the (inverse) inertia operator with $J_\alpha = \wp_\theta(\alpha)$, where $\wp_\theta(\alpha)$ are the values of the Weierstrass \wp -function and defined by (B.11). Then (3.3) assumes the form

$$\partial_t S_\alpha = \sum_{\gamma \in \tilde{\mathbb{Z}}_N^{(2)}} S_{\alpha-\gamma} S_\gamma \wp_\theta(\gamma) \mathbf{C}_\theta(\gamma, \alpha). \quad (3.4)$$

2. The linear brackets and r -matrix.

Proposition 3.1. (cf. [2]). *The equations of motion (3.2) have the Lax form*

$$\partial_t L^{rot}(z) = [L^{rot}(z), M(z)], \quad (3.5)$$

where

$$L^{rot} = - \sum_{\alpha} S_\alpha \varphi_\theta(\alpha, z) T_\alpha, \quad (3.6)$$

$$M = - \sum_{\alpha} S_\alpha f_\theta(\alpha, z) T_\alpha, \quad (3.7)$$

and the functions φ_θ, f_θ are given by (B.13), (B.14).

Proof. Substituting (3.6) and (3.7) in the Lax equation we obtain

$$\partial_t S_\alpha \varphi_\theta(\alpha, z) = \sum_{\gamma} S_{\alpha-\gamma} S_\gamma (\varphi_\theta(\alpha - \gamma, z) f_\theta(\gamma, z) - \varphi_\theta(\gamma, z) f_\theta(\alpha - \gamma, z)).$$

Now, using the explicit expressions (B.13) and (B.14) for φ_θ and f_θ respectively, as well as the Calogero functional equation (A.22), we come to

$$\partial_t S_\alpha = \sum_{\gamma} S_{\alpha-\gamma} S_\gamma \varphi_\theta(\alpha, z) (\wp_\theta(\alpha - \gamma) - \wp_\theta(\gamma)),$$

which coincides with (3.4). \square

The Lie-Poisson brackets (3.1) admit the following r -matrix description [4, 13, 14, 17]. Define the classical r -matrix by

$$r_{N,\theta,\tau}(z-w) = \sum_{\gamma} \varphi_\theta(\gamma, z-w) T_\gamma \otimes T_{-\gamma}, \quad (3.8)$$

where φ_θ is defined by (B.13).

Lemma 3.1. [14]. *The r -matrix (3.8) satisfies the classical Yang-Baxter equation*

$$\begin{aligned} [r_{N,\theta,\tau}^{(12)}(z-w), r_{N,\theta,\tau}^{(13)}(z)] + [r_{N,\theta,\tau}^{(12)}(z-w), r_{N,\theta,\tau}^{(23)}(w)] \\ + [r_{N,\theta,\tau}^{(13)}(z), r_{N,\theta,\tau}^{(23)}(w)] = 0. \end{aligned} \quad (3.9)$$

Proof. We reduce the YB equation (3.9) to the following functional equation

$$\varphi_\theta(\gamma, z-w)\varphi_\theta(\alpha, z) - \varphi_\theta(\alpha + \gamma, z-w)\varphi_\theta(\alpha, w) + \varphi_\theta(\alpha + \gamma, z)\varphi_\theta(-\gamma, w) = 0,$$

by using the commutation relations in $\mathfrak{sl}(N, \mathbb{C})$ (B.7). It is easy to see that it can be rewritten for the functions ϕ_θ (B.12), since all three terms have the common exponent $\mathbf{e}_\theta((\gamma_2 + \alpha_2)z - \gamma_2 w)$. Now, it coincides with the Fay identity (A.21), where we put

$$u_1 = (\gamma + \alpha)\theta, \quad u_2 = -\gamma\theta, \quad z_1 = z, \quad z_2 = w. \quad \square$$

Proposition 3.2. *In terms of the Lax operator (3.7) the brackets (3.1) are equivalent to the following relation for the Lax operator*

$$\{L_1^{rot}(z), L_2^{rot}(w)\}_1 = [r_{N,\theta,\tau}(z-w), L^{rot}(z) \otimes Id + Id \otimes L^{rot}(w)], \quad (3.10)$$

$$L_1 = L \otimes Id, \quad L_2 = Id \otimes L.$$

Proof. To prove (3.10) we rewrite it in the basis $T_\alpha \otimes T_\beta$:

$$\begin{aligned} \{S_\alpha, S_\beta\}_1 \varphi_\theta(\alpha, z)\varphi_\theta(\beta, w) \\ = S_{\alpha+\beta} C_\theta(\alpha, \beta) (\varphi_\theta(-\beta, z-w)\varphi_\theta(\alpha + \beta, z) - \varphi_\theta(\alpha, z-w)\varphi_\theta(\alpha + \beta, w)). \end{aligned}$$

The same Fay identity (A.21) implies that

$$\varphi_\theta(\alpha, z)\varphi_\theta(\beta, w) = \varphi_\theta(-\beta, z-w)\varphi_\theta(\alpha + \beta, z) - \varphi_\theta(\alpha, z-w)\varphi_\theta(\alpha + \beta, w).$$

Thus we come to (3.1). \square

3. The hierarchy of the Lax equations. The Lax operator L^{rot} (3.6) has the following properties:

i) L^{rot} is an $\mathfrak{sl}(N, \mathbb{C})$ -valued meromorphic function on E_τ with a simple pole at the origin satisfying

$$Res L^{rot}(z)|_{z=0} = \sum_{\alpha} S_{\alpha} T_{\alpha}.$$

ii) L^{rot} satisfies the quasi-periodicity conditions

$$L^{rot}(z+1) = QL^{rot}(z)Q^{-1}, \quad L^{rot}(z+\tau) = \Lambda L^{rot}(z)\Lambda^{-1}.$$

These properties imply that $Tr(L(z))^k$ are doubly periodic functions with the poles up to the order k . Thereby, they can be expanded in the basis of the Weierstrass function and its derivatives

$$tr(L^{rot}(z))^k = I_{0,k} + I_{2,k}\wp(z) + \dots + I_{k,k}\wp^{(k-2)}(z). \quad (3.11)$$

In particular, in this way we obtain the Hamiltonian

$$tr(L^{rot}(z))^2 = I_{0,2} + I_{2,2}\wp(z), \quad I_{0,2} = 2 \left(\frac{i}{2\pi\theta} \right)^2 H, \quad I_{2,2} = tr S^2.$$

Proposition 3.3. *The coefficients $I_{s,k}$ are in involution with respect to the linear bracket (3.1)*

$$\{I_{s,k}, I_{m,j}\}_1 = 0. \quad (3.12)$$

Proof. We have

$$\{\mathrm{tr}(L_1^{\mathrm{rot}}(z))^k, \mathrm{tr}(L_2^{\mathrm{rot}}(w))^j\}_1 = \mathrm{tr}\{(L_1^{\mathrm{rot}}(z))^k, (L_2^{\mathrm{rot}}(w))^j\}_1.$$

Then, it follows from (3.10) that these functionals Poisson commute. Using the expansion (3.11) we arrive at the involutivity of the coefficients (3.12). \square

In particular, all functions $I_{s,k}$ Poisson commute with the Hamiltonian H (3.2). Therefore, they play the role of conservation laws of the elliptic rotator hierarchy on $\mathrm{GL}(N, \mathbb{C})$. We have a tower of $\frac{N(N+1)}{2}$ independent integrals of motion

$$\begin{array}{ccccccc} I_{0,2} & I_{2,2} & & & & & \\ I_{0,3} & I_{2,3} & I_{3,3} & & & & \\ \dots & \dots & \dots & \dots & & & \\ I_{0,n} & I_{2,n} & \dots & \dots & I_{N,N} & & \end{array}$$

Note that $I_{k,k}$, $k = 0, 2, 3, \dots, N$ are the Casimirs corresponding to the coadjoint orbit

$$\mathcal{R}^{\mathrm{rot}} = \{\mathbf{S} \in \mathfrak{gl}(N, \mathbb{C}), \mathbf{S} = g^{-1} \mathbf{S}^{(0)} g\}.$$

The conservation laws $I_{s,k}$ generate commuting flows on $\mathcal{R}^{\mathrm{rot}}$,

$$\partial_{s,k} \mathbf{S} = \{I_{s,k}, \mathbf{S}\}_1, \quad (\partial_{s,k} := \partial_{t_{s,k}}),$$

that we call ER_N hierarchy.

In what follows we will need another set of the conservation laws coming from the coefficients of the spectral curve

$$\mathcal{C} : F(\lambda, z) \equiv \det(\lambda \cdot Id + L^{\mathrm{rot}}(z)) = 0. \quad (3.13)$$

Here

$$\begin{aligned} F(\lambda, z) &= \lambda^N + \lambda^{N-2} a_2(z) + \dots + a_N(z), \\ a_k(z) &= J_{0k} + J_{2k} \wp(z) + \dots + J_{kk} \wp^{(k-2)}(z). \end{aligned} \quad (3.14)$$

Since $a_k(z)$ and $\mathrm{Tr}(L(z))^k$ are related by the Newton formula, one can recursively represent $J_{s,k}$ as polynomials of $I_{m,j}$, $j \leq k$, $m \leq j$,

$$J_{s,k} = \sum a_{l,m,j}^{s,k} \prod_{\sum j_i m_i = k, l_i \leq j_i} (I_{l_i, j_i})^{m_i}, \quad (\mathbf{l} = (l_1, l_2, \dots)). \quad (3.15)$$

For example,

$$\begin{aligned} J_{s,2} &= -I_{s,2}, \quad (s = 0, 2); \quad J_{s,3} = -I_{s,3}, \quad (s = 0, 2, 3); \\ J_{0,4} &= \frac{3}{2} I_{0,4} - I_{0,2}^2 - \frac{g_2}{12} I_{2,2}^2, \quad (g_2 = 60 \sum_{m,n}^l (m + n\tau)^{-4}); \\ J_{2,4} &= \frac{3}{2} I_{2,4} - 2I_{0,2} I_{2,2}; \quad J_{3,4} = \frac{3}{2} I_{3,4} \quad J_{4,4} = \frac{3}{2} I_{4,4} - I_{2,2}^2, \end{aligned} \quad (3.16)$$

where in the second line we have exploited the relation $12\wp^2 = 2\wp'' + g_2$.

3.2. The classical $GL(N, \mathbb{C})$ Feigin-Odesski algebras. Here we consider the Feigin-Odesski Poisson brackets related to $GL(N, \mathbb{C})$. In notations of Ref. [5] it is the classical limit of the quadratic associative algebra $\mathcal{Q}_{N^2, k}$.

It turns out that the same r -matrix (3.8) defines a quadratic Poisson algebra. We start with the Lax operator L^{rot} (3.6). Modify it in the following way:

$$L(z) = -S_0 Id + L^{rot}(z), \quad (3.17)$$

where S_0 commute with S_α with respect to the linear brackets (3.1). Define the brackets between the entries of L as follows

$$\{L_1(z), L_2(w)\}_2 = [r_{N, \theta, \tau}(z - w), L_1(z) \otimes L_2(w)], \quad (3.18)$$

(see Ref. [13, 14]). These brackets are Poisson, since the Jacobi identity is provided by the classical Yang-Baxter equation (3.9). A finite-dimensional quadratic Poisson algebra is extracted from (3.18) by eliminating the dependence on the spectral parameters z, w .

Proposition 3.4. *The quadratic Poisson algebra on $\mathfrak{gl}(N, \mathbb{C})^*$ has the form*

$$\{S_\alpha, S_0\}_2 = \sum_{\gamma \neq \alpha} S_{\alpha-\gamma} S_\gamma (\wp_\theta(\gamma) - \wp_\theta(\alpha - \gamma)) \mathbf{C}_\theta(\alpha, \gamma), \quad (3.19)$$

$$\{S_\alpha, S_\beta\}_2 = S_0 S_{\alpha+\beta} \mathbf{C}_\theta(\alpha, \beta) + \sum_{\gamma \neq \alpha, -\beta} S_{\alpha-\gamma} S_{\beta+\gamma} \mathbf{f}_\theta(\alpha, \beta, \gamma) \mathbf{C}_\theta(\gamma, \alpha - \beta), \quad (3.20)$$

where $\mathbf{C}_\theta(\alpha, \beta)$ are the $\mathfrak{sl}(N, \mathbb{C})$ structure constants (B.8),

$$\begin{aligned} \mathbf{f}_\theta(\alpha, \beta, \gamma) &= \zeta_\theta(\gamma) + \zeta_\theta(\beta - \alpha + \gamma) - \zeta_\theta(\beta + \gamma) + \zeta_\theta(\alpha - \gamma) \\ &= -\frac{\vartheta'(0) \vartheta_\theta(\alpha) \vartheta_\theta(\beta) \vartheta_\theta(\beta - \alpha + 2\gamma)}{\vartheta_\theta(\alpha - \gamma) \vartheta_\theta(\beta + \gamma) \vartheta_\theta(\beta - \alpha + \gamma) \vartheta_\theta(\gamma)}, \end{aligned} \quad (3.21)$$

and $\vartheta_\theta, \zeta_\theta$ are the zeta constants (B.10), (B.11).

Proof. Equation (3.18) yields the following form of the brackets

$$\begin{aligned} \{S_a, S_b\}_2 \varphi_\theta(a, z) \varphi_\theta(b, w) &= \sum_{\gamma} S_{a-\gamma} S_{b+\gamma} \varphi_\theta(\gamma, z - w) \varphi_\theta \\ &\quad \times (a - \gamma, z) \varphi_\theta(b + \gamma, w) \mathbf{C}_\theta(\gamma, a - b). \end{aligned} \quad (3.22)$$

Here we assume that the Latin indices (a, b, \dots) belong to the lattice $\mathbb{Z}_N^{(2)} = \tilde{\mathbb{Z}}_N^{(2)} \cup (0, 0)$, while the Greek indices belong to $\tilde{\mathbb{Z}}_N^{(2)}$ as above. Note first that the common exponents in the expressions of the functions φ_θ (B.13) coincide in the left and the right sides. This allows one to pass to the functions ϕ_θ and in this way to use the Fay-type identities. We rewrite the last expression in the form

$$\begin{aligned} \{S_a, S_b\}_2 \phi_\theta(a, z) \phi_\theta(b, w) &= \sum_{\gamma} S_{a-\gamma} S_{b+\gamma} \mathbf{C}_\theta(\gamma, a - b) \\ &\quad \times (\phi_\theta(\gamma, z - w) \phi_\theta(a - \gamma, z) \phi_\theta(b + \gamma, w) \\ &\quad - \phi_\theta(a - b - \gamma, z - w) \phi_\theta(b + \gamma, z) \phi_\theta(a - \gamma, w)). \end{aligned}$$

Consider first the case $b = 0$. Then we can apply (A.30) for $u = (a_1 + a_2\tau)\theta$, $v = (\gamma_1 + \gamma_2)\theta$. It immediately leads us to (3.19). If $a, b \neq 0$ we have the terms of two types. For $\gamma \neq a, -b$ one uses (A.27), (A.28) and finds the second term in the right side of (3.20). If $\gamma = a$ or $\gamma = -b$ one should take into account that $\phi_\theta(0, z) = 1$ and then use the Fay identity (A.21). This gives us the first term. The identity (3.21) follows from (A.29). \square

We denote by $P_{N,\theta,\tau}$ the quadratic Poisson algebra (3.19), (3.20). Recall that τ is the modular parameter of the auxiliary curve E_τ and $\theta = k/N$. The algebra $P_{2,\frac{1}{2},\tau}$ is the classical Sklyanin algebra [14]. In this case the r.h.s. of (3.20) contains only the first term.

As for the linear brackets, there exist N Casimirs $C_0^{(2)}, C_2^{(2)}, \dots, C_N^{(2)}$ of the algebra $P_{N,\theta,\tau}$ [18]. They can be read off from the expansion of the spectral curve (3.13) in the basis of elliptic functions

$$\det(S_0 \cdot Id + L^{rot}(z)) = C_0^{(2)} + \sum_{k=2}^N C_k^{(2)} \wp^{(k-2)}(z). \quad (3.23)$$

It follows from (3.13), (3.14) and (3.17) that they are linear combinations of the integrals $J_{s,k}$,

$$C_0^{(2)} = S_0^N + \sum_{m=2}^N S_0^{N-m} J_{0,m}, \quad (3.24)$$

$$C_k^{(2)} = \sum_{m=0}^{N-k} S_0^{N-k-m} J_{k,k+m}, \quad (k = 2, \dots, N). \quad (3.25)$$

3.3. The bihamiltonian structure. Two Poisson structures are called *compatible* (or, form a *Poisson pair*) if their linear combinations are Poisson structures as well.

Proposition 3.5. *The linear and quadratic Poisson brackets on $\mathfrak{gl}(N, \mathbb{C})$ are compatible.*

Proof. Let us replace $S_0 \rightarrow S_0 + \lambda$ in the quadratic brackets (3.19), (3.20). Then (3.19) does not change, while (3.20) acquires an additional linear term. In this way we define a one-parameter family of the brackets

$$\{\mathbf{S}, \mathbf{S}\}_\lambda := \{\mathbf{S}, \mathbf{S}\}_2 + \lambda \{\mathbf{S}, \mathbf{S}\}_1, \quad (3.26)$$

where $\{\mathbf{S}, \mathbf{S}\}_1$ denote the linear brackets (3.1). Therefore, the linear combination $\{\mathbf{S}, \mathbf{S}\}_\lambda$ of two brackets can be obtained from the quadratic bracket by a simple shift of S_0 . The result of the shift, of course, still satisfies the Jacobi identity, and hence it is a Poisson bracket for any λ . \square

We denote this family of the quadratic Poisson algebras by $P_{N,\theta,\tau,\lambda}$. The algebras are isomorphic for different λ and degenerate to the linear Poisson algebra $\mathfrak{gl}(N, \mathbb{C})^*$ as $\lambda \rightarrow \infty$.

Consider the Casimir functions of the Poisson algebra $P_{N,\theta,\tau,\lambda}$,

$$h_k(\lambda) = C_k^{(2)}(S_0 + \lambda) \quad (k = 0, 2, \dots, N).$$

It follows from (3.24) and (3.25) that $h_k(\lambda)$ are polynomials in λ ,

$$h_k(\lambda) = \sum_{m=0}^{N-k} h_{m,k} \lambda^m. \quad (3.27)$$

The coefficients are defined as

$$h_{a,k} = \frac{1}{a!} \partial_\lambda^{(a)} C_k^{(2)}(S_0 + \lambda)_{\lambda=0}.$$

It implies that

$$h_{a,0} = \frac{N!}{(N-a)!a!} S_0^{N-a} + \sum_{m=2}^{N-a} \frac{(N-m)!}{(N-m-a)!a!} S_0^{N-m-a} J_{0,m}, \quad (3.28)$$

$$h_{a,k} = \sum_{m=0}^{N-k-a} \frac{(N-k-m)!}{(N-k-m-a)!a!} S_0^{N-k-m-a} J_{k,k+m}, \quad (k = 2, \dots, N). \quad (3.29)$$

In particular,

$$h_{N-1,0} = N S_0, \quad h_{N-2,0} = \frac{N(N-1)}{2} S_0^2 + J_{0,2}. \quad (3.30)$$

Conversely, one can express $J_{m,k+m}$ as a linear combination of $h_{N-m-s,m}$, $s = 0, \dots, k$ from the relations (3.28) and (3.29). For example,

$$\begin{aligned} J_{j,j} &= h_{N-j,j}, \quad J_{j,j+1} = h_{N-j-1,j} - (N-j) S_0 h_{N-j,j}, \\ J_{j,j+2} &= h_{N-j-2,j} - (N-j) S_0 h_{N-j-1,j} + \frac{(N-j)(N-j-1)}{2} S_0^2 h_{N-j,j}. \end{aligned}$$

We arrange the quantities $h_{m,k}$ in the triangular tableau

$$\begin{array}{ccccccccccccccc} h_{0,0} & h_{1,0} & \dots & \dots & \dots & \dots & \dots & \dots & \dots & & h_{N-1,0} & h_{N,0} \\ & h_{0,2} & h_{1,2} & & \dots & \dots & \dots & & h_{N-3,2} & & h_{N-2,2} & \\ & & h_{0,3} & h_{1,3} & \dots & \dots & \dots & h_{N-4,4} & & h_{N-3,3} & & \\ & & & \dots & \dots & \dots & \dots & & \dots & & & \\ & & & & h_{0,N-1} & & & h_{1,N-1} & & & & \\ & & & & & h_{0,N} & & & & & & \end{array} \quad (3.31)$$

Note that the second line corresponding to $h_{s,1}$ is absent since $h_1(\lambda) = 0$. The left side of the triangle contains the Casimirs $h_{0,j} = C_j^{(2)}$ (3.24), (3.25) of $P_{N,\theta,\tau}$, while the right side represents the Casimirs $h_{N-j,j} = J_{j,j}$ of the linear brackets on $\mathfrak{gl}(N, \mathbb{C})^*$.

The remarkable property of the quantities $h_{a,k}$ is that they are in involution with respect to both the linear and quadratic brackets:

$$\{h_{a,k}, h_{b,j}\}_{1,2} = 0,$$

see, e.g., [19]. This fact follows from the identities

$$\{h_k(\lambda), h_j(\lambda)\}_\lambda = 0, \quad (3.32)$$

held for each λ .

Proposition 3.6. *The integrals $h_{a,k}$ provide the recurrence relation for the elliptic rotator hierarchy ER_N ,*

$$\{h_{a,k}, \mathbf{S}\}_1 = -\{h_{a+1,k}, \mathbf{S}\}_2. \quad (3.33)$$

Proof. Since $h_k(\lambda)$ are the Casimirs, we have $\{h_k(\lambda), \mathbf{S}\}_\lambda = 0$. Substitute the representation (3.27) into this equation. Then the recurrence relation (3.33) comes from the coefficient in front of λ^a . \square

In this way one can start with the Casimirs of the linear brackets and produce a non-trivial dynamical system using the quadratic brackets

$$\{J_{k,k}, \mathbf{S}\}_2 = -\{J_{k,k+1}, \mathbf{S}\}_1, \quad (3.34)$$

(see (3.30)). In particular, the flow (3.4) corresponding to $H = I_{0,2}$ can be represented by the quadratic brackets with S_0 :

$$\partial_t \mathbf{S} = \{S_0, \mathbf{S}\}_2.$$

Alternatively, one can start with the Hamiltonians in the left side of the table (3.31) and the linear brackets.

This allows us to conclude the proof of Theorem 2.1 for the hierarchy ER_N . Namely, Propositions 3.2, 3.4, 3.5, and 3.6 are equivalent, respectively, to i), ii), iii), and iv).

4. Elliptic Rotators on Non-Commutative Torus

4.1. The Lax equation and integrals of motion.

1. The system description. The elliptic rotator on the non-commutative torus \mathcal{A}_θ is a generalization of the elliptic rotator on $GL(N, \mathbb{C})$. We consider the Euler-Arnold top on the group SIN_θ of the NCT algebra \mathcal{A}_θ (Appendix C). The Lie coalgebra \sin_θ^* is equipped with the linear Poisson brackets

$$\{S_\alpha, S_\beta\}_1 = C_\theta(\alpha, \beta) S_{\alpha+\beta}, \quad (\alpha \in \tilde{\mathbb{Z}}^{(2)}), \quad (4.1)$$

where $\mathbf{S} = \sum S_{-\alpha} T_\alpha$ and

$$\tilde{\mathbb{Z}}^{(2)} = \{\alpha = (\alpha_1, \alpha_2), \alpha_j \in \mathbb{Z}, \alpha \neq (0, 0)\}.$$

The inverse inertia tensor \mathbf{J} maps $\sin_\theta^* \rightarrow \sin_\theta$, and depends on four parameters: $\theta, \tau \in \mathbb{C}$, ($\Im m \tau > 0$) and $\epsilon = (\epsilon_1, \epsilon_2)$, where $0 < \theta_{\epsilon_a} \leq 1$ and $\epsilon_a \theta$ are irrational numbers. The components of \mathbf{J} are given by the elliptic constants $J_\alpha = \wp_{\theta, \epsilon}(\alpha)$ (C.18),

$$\mathbf{J}(\mathbf{S}) = \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} \wp_{\theta, \epsilon}(\alpha) S_{-\alpha} T_\alpha.$$

The Hamiltonian has the form

$$H = 2\pi^2 \theta^2 \int_{\mathcal{A}_\theta} \mathbf{S} \cdot \mathbf{J}(\mathbf{S}) = -\frac{1}{2} \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} \wp_{\theta, \epsilon}(\alpha) S_\alpha S_{-\alpha}. \quad (4.2)$$

Introduce a complex structure on the NCT \mathcal{A}_θ depending on $\epsilon = (\epsilon_1, \epsilon_2)$ and τ such that for $X = \sum_a c_a T_a$ one has:

$$\bar{\partial}_{\epsilon, \tau} X = \sum_a (\epsilon_1 a_1 + \epsilon_2 a_2 \tau) c_a T_a. \quad (4.3)$$

Then, in the Moyal representation, the operator \mathbf{J} can be identified with the pseudo-differential operator

$$\mathbf{J}(\mathbf{S})(x) = \wp(\theta \bar{\partial}_{\epsilon, \tau}) \mathbf{S}(x), \quad (4.4)$$

and

$$H = 2\pi^2 \theta^2 \int_{\mathcal{A}_\theta} \mathbf{S} \cdot \wp(\theta \bar{\partial}_{\epsilon, \tau}) \mathbf{S}. \quad (4.5)$$

The equation of motion in the form of the Moyal brackets has the standard form

$$\partial_t \mathbf{S} = \{\mathbf{S}, (\wp(\theta \bar{\partial}_{\epsilon, \tau}) \mathbf{S})\}^\theta := \text{ad}_{\mathbf{J}(\mathbf{S})}^* \mathbf{S}, \quad (4.6)$$

or, in the components,

$$\partial_t S_\alpha = \sum_{\gamma \in \tilde{\mathbb{Z}}^{(2)}} S_\gamma \wp_{\theta, \epsilon}(\gamma) S_{\alpha - \gamma}. \quad (4.7)$$

2. The r -matrix and the Lax equation. Define formally the Lax operator

$$L^{rot}(z) = - \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} S_\alpha \varphi_{\theta, \epsilon}(\alpha, z) T_\alpha, \quad (4.8)$$

where $\varphi_{\theta, \epsilon}$ is given by (C.19). Below we formulate the conditions on the phase space, which show that $L^{rot}(z)$ is well defined for $z \in E_\tau$, $z \neq 0$. Note that

$$\int_{\mathcal{A}_\theta} L^{rot}(z) = 0.$$

The following proposition is an infinite-dimensional analog of Proposition 3.1.

Proposition 4.1. *The equations of motion (4.7) have the Lax form*

$$\partial_t L^{rot} = [L^{rot}, M^{rot}] := \{L^{rot}, M^{rot}\}^\theta \quad (4.9)$$

with

$$M^{rot}(z) = - \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} S_\alpha f_{\theta, \epsilon}(\alpha, z) T_\alpha, \quad (4.10)$$

and $f_{\theta, \epsilon}$ (C.21).

The proof is analogous to the finite-dimensional case. It is based on the Calogero functional equation (A.22).

Introduce the classical r -matrix on $\sin_\theta \otimes \sin_\theta$ as the following sum:

$$r_{\theta, \epsilon, \tau}(z - w) = \sum_{\gamma \in \tilde{\mathbb{Z}}^{(2)}} \varphi_{\theta, \epsilon}(\gamma, z - w) T_\gamma \otimes T_{-\gamma}, \quad (4.11)$$

Lemma 3.1 and Proposition 3.2 have the following analogs for the NCT:

Proposition 4.2. *a) The r -matrix (4.11) satisfies the classical Yang-Baxter equation (3.9).*

b) The r -matrix defines the Poisson brackets for the entries of the Lax operator (4.8),

$$\{L_1^{rot}(z), L_2^{rot}(w)\}_1 = [r_{\theta, \epsilon, \tau}(z - w), L^{rot}(z) \otimes Id + Id \otimes L^{rot}(w)]. \quad (4.12)$$

They are equivalent to the linear brackets (4.1) on \sin_θ^ .*

The proof of Proposition 4.2(a, b) is similar to those of Lemma 3.1 and Proposition 3.2 and is based on the Fay identity (A.21).

3. The hierarchy of the Lax equations. In order to construct integrals of motion we first discuss the properties of the Lax operator. This operator is the meromorphic quasi-periodic function on E_τ with a simple pole

$$\begin{aligned} \text{Res } L^{rot}(z)|_{z=0} &= \mathbf{S}, \\ L^{rot}(z+1) &= U_1^{\epsilon_1} L^{rot}(z) U_1^{-\epsilon_1}, \quad L^{rot}(z+\tau) = U_2^{\epsilon_2} L^{rot}(z) U_2^{-\epsilon_2}, \end{aligned}$$

where U_1, U_2 are the generators of NCT \mathcal{A}_θ (C.1). (This follows from (4.8), (A.9) and (B.15).) It means that the integrals

$$\int_{\mathcal{A}_\theta} (L^{rot}(z))^j$$

are doubly periodic functions on E_τ with poles up to order j . Thereby, we have the expansion

$$\int_{\mathcal{A}_\theta} (-L^{rot}(z))^j = I_{0,j} + \sum_{r=2}^j I_{r,j} \wp^{(r-2)}(z). \quad (4.13)$$

In particular,

$$\int_{\mathcal{A}_\theta} (L_\theta)^2(z) = I_{0,2} + \wp(z) \int_{\mathcal{A}_\theta} \mathbf{S}^2, \quad \text{where } I_{0,2} = 2 \left(\frac{i}{2\pi\theta} \right)^2 H.$$

Note that

$$I_{j,j} = \int_{\mathcal{A}_\theta} \mathbf{S}^j$$

are the Casimirs of the linear brackets.

The phase space \mathcal{R} of the elliptic rotator is determined by the following properties of the integrals $I_{s,j}$:

$$\mathcal{R} = \{ \mathbf{S} \in \mathcal{A}_\theta^* \mid i) I_{s,j} < \infty, \quad ii) \lim_{j \rightarrow \infty} \frac{I_{s,j+1}(\mathbf{S})}{I_{s,j}(\mathbf{S})} < 1 \text{ for all } s \leq j \}. \quad (4.14)$$

The first condition means that the traces (4.13) of the Lax operator are well defined on E_τ for $z \neq 0$. In particular, we have

$$I_{0,2} < \infty, \quad \text{i.e.} \quad \sum \wp_\theta(\alpha) S_\alpha S_{-\alpha} < \infty. \quad (4.15)$$

We will use the second condition below to define the Casimir functions for the family of quadratic Poisson brackets.

Further, due to Lemma 3.1, an analog of Proposition 3.2 is the following.

Proposition 4.3. *The quantities $I_{s,k}$ pairwise Poisson commute.*

The conservation laws $I_{s,k}$ generate commuting flows on the phase space \mathcal{R} with respect to the linear brackets

$$\partial_{s,k} \mathbf{S} = \{I_{s,k}, \mathbf{S}\}_1, \quad (4.16)$$

where $\partial_{s,k}$ stands for the corresponding time derivative $\partial_{t_{s,k}}$. We call these equations *ER_θ hierarchy on the NCT*.

One can show that all flows can be represented in the Lax form with L^{rot} (4.8) and the corresponding $M_{s,k}$. Furthermore, there exists a set of the integrals of $J_{s,k}$ -type. They are related to the integrals $I_{m,j}$ by the same formulae (3.15), (3.16).

4.2. Quadratic Poisson algebras and the bihamiltonian structure on NCT. To define the quadratic Poisson algebra on the phase space \mathcal{R} (4.14) we modify the Lax operator (4.8):

$$L = -S_0 \cdot T_0 + L^{rot} = -S_0 \cdot T_0 - \sum_{\alpha} S_{\alpha} \varphi_{\theta, \epsilon}(\alpha, z) T_{\alpha}, \quad (4.17)$$

where S_0 commutes with S_{α} with respect to the linear brackets. Due to the YB equation, the r -matrix (4.11) defines the quadratic Poisson brackets

$$\{L_1(z), L_2(w)\}_2 = [r_{\theta, \epsilon, \tau}(z - w), L_1(z) \otimes L_2(w)]. \quad (4.18)$$

Again, we can get rid of the dependence on the spectral parameter.

Proposition 4.4. *The quadratic Poisson brackets on the phase space \mathcal{R} are defined as follows*

$$\{S_{\alpha}, S_0\}_2 = \sum_{\gamma \neq \alpha} S_{\alpha - \gamma} S_{\gamma} (\wp_{\theta, \epsilon}(\gamma) - \wp_{\theta, \epsilon}(\alpha - \gamma)) \mathbf{C}_{\theta}(\alpha, \gamma), \quad (4.19)$$

$$\begin{aligned} \{S_{\alpha}, S_{\beta}\}_2 &= S_0 S_{\alpha + \beta} \mathbf{C}_{\theta}(\alpha, \beta) + \sum_{\gamma \neq \alpha, -\beta} S_{\alpha - \gamma} S_{\beta + \gamma} \mathbf{f}_{\theta, \epsilon}(\alpha, \beta, \gamma) \\ &\quad \times \mathbf{C}_{\theta}(\gamma, \alpha - \beta), \end{aligned} \quad (4.20)$$

where $\mathbf{C}_{\theta}(\alpha, \gamma)$ is a structure constant of the \sin_{θ} -algebra,

$$\begin{aligned} \mathbf{f}_{\theta, \epsilon}(\alpha, \beta, \gamma) &= \zeta_{\theta, \epsilon}(\gamma) + \zeta_{\theta, \epsilon}(\beta - \alpha + \gamma) - \zeta_{\theta, \epsilon}(\beta + \gamma) + \zeta_{\theta, \epsilon}(\alpha - \gamma) \\ &= -\frac{\vartheta'(0) \vartheta_{\theta, \epsilon}(\alpha) \vartheta_{\theta, \epsilon}(\beta) \vartheta_{\theta, \epsilon}(\beta - \alpha + 2\gamma)}{\vartheta_{\theta, \epsilon}(\alpha - \gamma) \vartheta_{\theta, \epsilon}(\beta + \gamma) \vartheta_{\theta, \epsilon}(\beta - \alpha + \gamma) \vartheta_{\theta, \epsilon}(\gamma)}, \end{aligned} \quad (4.21)$$

and $\vartheta_{\theta, \epsilon}, \zeta_{\theta, \epsilon}$ are the constants (C.17), (C.18).

Proof. The equality (4.18), being reduced to the coefficients in the front of $T_a \otimes T_b$, can be rewritten in the form (3.22) where the functions $\varphi_{\theta}(a, z)$ are replaced on $\varphi_{\theta, \epsilon}(a, z)$. Then using the cubic functional equations (A.27) and (A.28) we obtain to the algebra (4.19), (4.20). Due to the definition of the phase space (4.15) the series in the right-hand sides of (4.19), (4.20) converge. \square

We denote this Poisson algebra by $P_{\theta, \epsilon, \tau}$. Set now $\epsilon_1 = \epsilon_2 = 1$. The corresponding algebra $P_{\theta, 1, \tau}$ can be considered as a special large N limit of the finite-dimensional algebras $P_{N, \theta, \tau}$ (3.19), (3.20). For this we just replace the rational number $\theta = k/N$ by an arbitrary irrational number $0 \leq \theta < 1$ in the algebra $P_{N, \theta, \tau}$.

Proposition 4.5. *a) The linear (4.1) and quadratic (4.19), (4.20) Poisson brackets defined on \mathcal{R} are compatible.*

b) The ER_θ hierarchy admits the bihamiltonian structure.

Proof. One can shift $S_0 + \lambda$ and define the family of Poisson algebras $P_{\theta, \epsilon, \tau, \lambda}$. To define the Casimir elements on $P_{\theta, \epsilon, \tau, \lambda}$ we consider

$$\log \det(S_0 + \lambda + L^{rot}(z)) = \log(S_0 + \lambda) + \sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{k(S_0 + \lambda)^k} \text{tr} L(z)^k.$$

Then we come to the infinite set of Casimirs ¹

$$\begin{aligned} \tilde{C}_0^{(2)}(S_0 + \lambda) &= \log(S_0 + \lambda) + \sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{k(S_0 + \lambda)^k} I_{0,k}, \\ \tilde{C}_j^{(2)}(S_0 + \lambda) &= \sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{k(S_0 + \lambda)^k} I_{j,k}, \quad (j = 2, 3, \dots). \end{aligned}$$

These functionals are well-defined on the phase space (4.14). This allows us to introduce the new set of conserved quantities

$$\begin{aligned} \tilde{C}_0^{(2)}(S_0 + \lambda) &= \sum_s \tilde{h}_{s,0} \lambda^s, \\ \tilde{C}_j^{(2)}(S_0 + \lambda) &= \sum_s \tilde{h}_{s,j} \lambda^s. \end{aligned}$$

The latter Poisson commute with respect to both types of brackets and give rise to the bihamiltonian structure of the hierarchy

$$\{\tilde{h}_{s,j}, \mathbf{S}\}_1 = -\{\tilde{h}_{s+1,j}, \mathbf{S}\}_2. \quad (4.22)$$

In particular, we can represent the flow (4.6) in the form

$$\partial_t \mathbf{S} = \{\mathbf{S}_0, \mathbf{S}\}_2.$$

(see (3.30)). \square

This proposition concludes the proof of Theorem 2.1 for the hierarchy ER_θ .

5. Elliptic Rotators on $SDiff(T^2)$

5.1. Description of the hierarchy. In the "dispersionless" limit $\theta \rightarrow 0$ the Lie algebra \sin_θ turns to the Lie algebra $Ham(T^2)$ of Hamiltonians on a two-dimensional torus, see (C.22). This algebra of Hamiltonian functions can be represented by the Lie algebra of the corresponding divergence-free vector fields $SVect(T^2)$. More precisely, to pass from $Ham(T^2)$ to $SVect(T^2)$ one has to discard the constant Hamiltonians, but add the "flux" vector fields $\partial/\partial x_1$ and $\partial/\partial x_2$ corresponding to multivalued Hamiltonian functions x_1 and x_2 on the torus.

We define the elliptic rotator system ER on the Lie group $SDiff(T^2)$ by considering the limit $\theta \rightarrow 0$ of the ER_θ -system described above. Let $\theta \rightarrow 0$, $\epsilon_{1,2} \rightarrow \infty$, such that

$$\lim_{\theta \rightarrow 0} (\theta \epsilon_{1,2}) = \epsilon'_{1,2} < 1, \quad \epsilon'_{1,2} \text{ are irrational}. \quad (5.1)$$

In what follows we drop the superscript $'$.

¹ Since we pass from \det to $\log \det$ the basis of the Casimir functions differs from (3.24).

Let $\mathbf{S} = \sum_{\alpha} S_{-\alpha} \mathbf{e}(\alpha \cdot x) \in \text{Ham}^*(T^2)$, where $\mathbf{e}(\alpha \cdot x)$ is the Fourier basis (C.23) of $\text{Ham}^*(T^2)$. In the Fourier basis, the linear Poisson bracket assumes the form

$$\{S_{\alpha}, S_{\beta}\}_1 = (\alpha \times \beta) S_{\alpha+\beta}, \quad \alpha \times \beta = \alpha_1 \beta_2 - \alpha_2 \beta_1. \quad (5.2)$$

The inverse inertia operator $\mathbf{J} : \text{Ham}^*(T^2) \rightarrow \text{Ham}(T^2)$ becomes

$$\mathbf{J} : S_{\alpha} \rightarrow \wp_{\epsilon}(\alpha) S_{\alpha}, \quad \wp_{\epsilon}(\alpha) = \wp(\epsilon_1 \alpha_1 + \epsilon_2 \alpha_2 \tau; \tau),$$

where $\alpha \in \tilde{\mathbb{Z}}^{(2)}$ (B.5). The operator is well defined since ϵ_j are irrational. In other words, the operator \mathbf{J} is the pseudo-differential operator

$$\mathbf{J} : \mathbf{S}(x) \rightarrow \wp(\bar{\partial}_{\epsilon, \tau}) \mathbf{S}(x),$$

where $\bar{\partial}_{\epsilon, \tau}$ is the operator of the complex structure on the elliptic curve, commutative torus T^2 . In fact, the complex structure depends on the ratio $\tau \epsilon_2 / \epsilon_1$.

The quadratic Hamiltonian of the system is

$$H = -\frac{1}{2} \sum_{\gamma} S_{\gamma} \wp_{\epsilon}(\gamma) S_{-\gamma} = -\frac{1}{2} \int_{T^2} \mathbf{S}(\wp(\bar{\partial}_{\epsilon, \tau}) \mathbf{S}), \quad (5.3)$$

and the corresponding equations of motion are

$$\partial_t \mathbf{S} = \{\mathbf{S}, \wp(\bar{\partial}_{\epsilon, \tau}) \mathbf{S}\}_1. \quad (5.4)$$

This equation is an elliptic analog of the hydrodynamics equation on the torus T^2 , regarded as an elliptic curve, see the Remark below. We call it *the elliptic hydrodynamics*.

Define the Lax operator

$$L^{rot}(x; z) = - \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} S_{\alpha} \varphi_{\epsilon}(\alpha, z) \mathbf{e}(\alpha \cdot x), \quad \varphi_{\epsilon}(\alpha, z) = \varphi(\epsilon_1 \alpha_1 + \tau \epsilon_2 \alpha_2, z). \quad (5.5)$$

The conditions on the phase space formulated below, see (5.12), ensure that the operator $L^{rot}(x, z)$ is well defined for $z \in E_{\tau}$, $z \neq 0$. Note that

$$\int_{T^2} L_{\epsilon}^{rot}(z) = 0.$$

Proposition 5.1. *The equations of motion (5.4) have the dispersionless Lax representation*

$$L^{rot} = \{L^{rot}, M^{rot}\}_1$$

with L^{rot} given by (5.5) and

$$M^{rot}(x; z) = - \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} S_{\alpha} f_{\epsilon}(\alpha, z) \mathbf{e}(\alpha \cdot x), \quad (5.6)$$

where

$$f_{\epsilon}(\alpha, z) = f(\epsilon_1 \alpha_1 + \tau \epsilon_2 \alpha_2, z) = \mathbf{e}(\epsilon_2 \alpha_2) \partial_u \phi(u, z)|_{u=\epsilon_1 \alpha_1 + \tau \epsilon_2 \alpha_2}.$$

The proof of Proposition 5.1 is the same as Proposition 3.1.

Note that operators L^{rot} and M^{rot} satisfy the quasi-periodicity properties,

$$L^{rot}(x_1, x_2; z) = L^{rot}(x_1 + \epsilon_2, x_2; z + 1), \quad (5.7)$$

$$L^{rot}(x_1, x_2; z) = L^{rot}(x_1, x_2 - \epsilon_1; z + \tau), \quad (5.8)$$

$$M^{rot}(x_1, x_2; z) = M^{rot}(x_1 + \epsilon_2, x_2; z + 1), \quad (5.9)$$

and

$$M^{rot}(x_1, x_2; z) - M^{rot}(x_1, x_2 - \epsilon_1; z + \tau) = 2\pi i L^{rot}(x_1, x_2, z). \quad (5.10)$$

Furthermore, it follows from (5.7) and (5.8) that there exists the expansion

$$\int_{T^2} (L^{rot}(x, z))^k = I_{0,k} + \sum_{s=2}^k I_{s,k} \wp^{(s-2)}(z). \quad (5.11)$$

The phase space \mathcal{R}^0 of the elliptic rotator on $SDiff(T^2)$ can be described similarly to the phase space \mathcal{R} for SIN_θ -group, see (4.14):

$$\begin{aligned} \mathcal{R}^0 = \{ \mathbf{S} \in Ham^*(T^2) \mid i) I_{s,j} < \infty, \\ ii) \lim_{j \rightarrow \infty} \frac{I_{s,j+1}(\mathbf{S})}{I_{s,j}(\mathbf{S})} < 1 \text{ for all } s \leq j \}. \end{aligned} \quad (5.12)$$

Define the r -matrix on $T^2 \otimes T^2$ by

$$r_{\epsilon, \tau}(x, y; z) = \sum_{\alpha \in \mathbb{Z}^{(2)}} \varphi_\epsilon(\alpha, z) \mathbf{e}(\alpha \cdot x) \mathbf{e}(-\alpha \cdot y), \quad z \in E_\tau. \quad (5.13)$$

Proposition 5.2. *a) The r -matrix (5.13) satisfies the "dispersionless" YB equation*

$$\begin{aligned} \{r_{\epsilon, \tau}(x, y; z - w), r_{\epsilon, \tau}(x, v; z)\}_1 + \{r_{\epsilon, \tau}(x, y; z - w), r_{\epsilon, \tau}(y, v; w)\}_1 \\ + \{r_{\epsilon, \tau}(x, v; z), r_{\epsilon, \tau}(y, v; w)\}_1 = 0. \end{aligned} \quad (5.14)$$

b) In terms of the Lax operator (5.5) the canonical brackets on T^2 take the form

$$\begin{aligned} \{L^{rot}(x; z), L^{rot}(y; w)\}_1 = \{r_{\epsilon, \tau}(x, y, z - w), L^{rot}(x; z)\}_1 \\ + \{r(x, y, z - w) L^{rot}(y; w)\}_1. \end{aligned} \quad (5.15)$$

Proof. These statements are consequences of the Fay identity (A.21), cf. Proposition 4.2. \square

The form of the brackets (5.15) implies that

$$\left\{ \int_{T^2} (L^{rot}(x, z))^k, \int_{T^2} (L^{rot}(x, z))^j \right\}_1 = 0.$$

In turn, then (5.11) produces the infinite sequence of conservation laws $I_{s,k}$ in involution. They define the hierarchy of commuting flows

$$\partial_{s,k} \mathbf{S} = \{I_{s,k}, \mathbf{S}\}_1, \quad (5.16)$$

which is the dispersionless limit of the hierarchy (4.16). The equations can be represented in the form of the dispersionless Lax equations.

5.2. *Quadratic Poisson algebras, and the bihamiltonian structure on $S\text{Diff}(T^2)$.* To describe the quadratic Poisson brackets, we first pass from the Lie algebraic Lax operator L^{rot} to the operator

$$L_\epsilon = -S_0 + L^{\text{rot}}(x; z). \quad (5.17)$$

Define the quadratic Poisson algebra by the formula

$$\{L_\epsilon(x; z), L_\epsilon(y; w)\}_2 = \{r_{\epsilon, \tau}(x, y; z - w), L_\epsilon(x; z)L_\epsilon(y; w)\}_1. \quad (5.18)$$

Proposition 5.3. *In terms of the Fourier modes, the Poisson algebra (5.18) has the form*

$$\{S_\alpha, S_0\}_2 = \sum_{\gamma \neq \alpha} S_{\alpha-\gamma} S_\gamma (\wp_\epsilon(\gamma) - \wp_\epsilon(\alpha - \gamma))(\alpha \times \gamma), \quad (5.19)$$

$$\begin{aligned} \{S_\alpha, S_\beta\}_2 &= S_0 S_{\alpha+\beta}(\alpha \times \beta) + \sum_{\gamma \neq \alpha, -\beta} S_{\alpha-\gamma} S_{\beta+\gamma} \mathbf{f}_\epsilon(\alpha, \beta, \gamma) \\ &\quad \times (\gamma \times (\alpha - \beta)), \end{aligned} \quad (5.20)$$

where

$$\begin{aligned} \mathbf{f}_\epsilon(\alpha, \beta, \gamma) &= \zeta_\epsilon(\gamma) + \zeta_\epsilon(\beta - \alpha + \gamma) - \zeta_\epsilon(\beta + \gamma) + \zeta_\epsilon(\alpha - \gamma) \\ &= -\frac{\vartheta'(0)\vartheta_\epsilon(\alpha)\vartheta_\epsilon(\beta)\vartheta_\epsilon(\beta - \alpha + 2\gamma)}{\vartheta_\epsilon(\alpha - \gamma)\vartheta_\epsilon(\beta + \gamma)\vartheta_\epsilon(\beta - \alpha + \gamma)\vartheta_\epsilon(\gamma)}, \end{aligned}$$

$$\zeta_\epsilon(\gamma) = \zeta(\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2), \quad \vartheta_\epsilon(\gamma) = \vartheta(\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2).$$

The proof of this proposition is similar to that of Proposition 3.2.

By the same trick as before we can show that the linear and the quadratic brackets are compatible. It allows us to construct the integral of the $\tilde{h}_{s,k}$ type and define the recurrence representation (4.22) for the bihamiltonian structure on the hierarchy. In particular, Eq. (5.4) can be represented as

$$\partial_t \mathbf{S} = \{\mathbf{S}_0, \mathbf{S}\}_2.$$

5.3. Rational limit of the elliptic hydrodynamics.

1. *Description of the limit.* So far we have been dealing with the elliptic curve E_τ parameterized by the two half-periods ω_1, ω_2 , $\tau = \omega_2/\omega_1$. We replace our main functions in the following way:

$$\begin{aligned} \zeta_\epsilon(\gamma) &= \zeta(\epsilon_1\gamma_1\omega_1 + \epsilon_2\gamma_2\omega_2; \omega_1, \omega_2), \\ \wp_\epsilon(\gamma) &= \wp(\epsilon_1\gamma_1\omega_1 + \epsilon_2\gamma_2\omega_2; \omega_1, \omega_2). \end{aligned} \quad (5.21)$$

Now we are going to consider the rational limit of the elliptic curves, i.e., degeneration $\lim \omega_{1,2} \rightarrow \infty$. We look at the double scaling limit,

$$\lim \epsilon_{1,2} \rightarrow 0, \quad \lim \omega_{1,2} \rightarrow \infty, \quad \lim \epsilon_1\omega_1 = 1, \quad \lim \epsilon_2\omega_2 = \tau.$$

Then $\lim \omega_{1,2} \rightarrow \infty$ leads to the rational degeneration of the elliptic functions (5.21),

$$\lim_{\omega_{1,2} \rightarrow \infty} \zeta_\epsilon(\gamma) = \frac{1}{\gamma_\tau}, \quad \lim_{\omega_{1,2} \rightarrow \infty} \wp_\epsilon(\gamma) = \frac{1}{\gamma_\tau^2}, \quad \gamma_\tau = \gamma_1 + \gamma_2\tau.$$

This implies, in particular, that the inverse inertia tensor assumes the form

$$\mathbf{J}(\mathbf{S}) = \bar{\partial}^{-2} \mathbf{S}(x_1, x_2) = \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} \frac{1}{\alpha_\tau^2} S_{-\alpha} T_\alpha, \quad (5.22)$$

where $\alpha_\tau = \alpha_1 + \alpha_2 \tau$. The equations of motion are defined by the corresponding quadratic Hamiltonian,

$$H = -\frac{1}{2} \int_{T^2} \mathbf{S} \bar{\partial}^{-2} \mathbf{S} = -\frac{1}{2} \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} \frac{1}{\alpha_\tau^2} S_\alpha S_{-\alpha}, \quad (5.23)$$

with respect to the linear Poisson bracket on $Ham(T^2)$:

$$\partial_t \mathbf{S} = \{\mathbf{S}, \bar{\partial}^{-2} \mathbf{S}\}_1. \quad (5.24)$$

In the Fourier components the latter becomes

$$\partial_t S_\alpha = \sum_{\gamma \in \tilde{\mathbb{Z}}^{(2)}} S_\gamma \frac{1}{\gamma_\tau^2} S_{\alpha-\gamma}.$$

We call these equations *the modified hydrodynamics* on the torus T^2 .

Remark 5.1. Recall that the standard hydrodynamics of an ideal fluid on the torus T^2 is given by the Euler equation

$$\partial_t \mathbf{S} = \{\mathbf{S}, \Delta^{-1} \mathbf{S}\}_1,$$

on the vorticity function \mathbf{S} . The definition of the Laplace operator uses metric, while to define the modified hydrodynamics (5.24) we need to fix a complex structure on T^2 .

Now consider the Lax representation for the modified hydrodynamics. In the double scaling limit we have

$$\begin{aligned} \lim \varphi_\epsilon(\alpha, z) &= \exp(2\pi i \alpha_1 z) \left(\frac{1}{\alpha_\tau} + \frac{1}{z} \right), \\ \lim f_\epsilon(\alpha, z) &= -\exp(2\pi i \alpha_1 z) \frac{1}{\alpha_\tau^2}. \end{aligned}$$

Therefore, the Lax pair (5.6) in the rational limit is

$$\begin{aligned} L(x_1, x_2; z) &= -\sum_{\alpha} S_\alpha \exp(2\pi i \alpha_1 z) \left(\frac{1}{\alpha_\tau} + \frac{1}{z} \right) T_\alpha, \\ M(x_1, x_2) &= -\sum_{\alpha} S_\alpha \exp(2\pi i \alpha_1 z) \frac{1}{\alpha_\tau^2} T_\alpha. \end{aligned}$$

We can drop the exponents in both operators and come to the following expressions

$$L(x_1, x_2; z) = -\bar{\partial}^{-1} \mathbf{S}(x_1, x_2) - \frac{1}{z} \mathbf{S}(x_1, x_2), \quad (5.25)$$

and

$$M(x_1, x_2) = -\bar{\partial}^{-2} \mathbf{S}(x_1, x_2). \quad (5.26)$$

It is easy to see that the dispersionless Lax equation

$$\partial_t L(x_1, x_2; z) = \{L(x_1, x_2; z), M(x_1, x_2)\}_2$$

is equivalent to Eqs. (5.24).

The infinite set of the integrals of motion comes from the decomposition

$$(-1)^k \int_{T^2} L^k(x_1, x_2; z) dx_1 dx_2 = I_{0,k} + \sum_{s=2}^k I_{m,k} z^{-m}.$$

The corresponding infinite hierarchy of the modified hydrodynamics

$$\partial_{s,t} \mathbf{S} = \{I_{s,k}, \mathbf{S}\}_1$$

has the dispersionless Lax form representation.

2. Bihamiltonian structure. It turns out that the bihamiltonian structure survives in this limit. In fact, we have the following quadratic Poisson algebra $\mathcal{P}_{0,\tau}$ on the commutative torus T^2 :

$$\{S_\alpha, S_0\}_2 = \sum_{\gamma \neq \alpha} S_{\alpha-\gamma} S_\gamma \left(\frac{1}{\gamma_\tau^2} - \frac{1}{(\alpha_\tau - \gamma_\tau)^2} \right) (\alpha \times \gamma), \quad (5.27)$$

$$\begin{aligned} \{S_\alpha, S_\beta\}_2 &= S_0 S_{\alpha+\beta} (\alpha \times \beta) + \sum_{\gamma \neq \alpha, -\beta} S_{\alpha-\gamma} S_{\beta+\gamma} f(\alpha, \beta, \gamma) \\ &\quad \times (\gamma \times (\alpha - \beta)), \end{aligned} \quad (5.28)$$

where

$$f(\alpha, \beta, \gamma) = \left(\frac{1}{\gamma_\tau} + \frac{1}{\beta_\tau - \alpha_\tau + \gamma_\tau} - \frac{1}{\beta_\tau + \gamma_\tau} + \frac{1}{\alpha_\tau - \gamma_\tau} \right), \quad \gamma_\tau = \gamma_1 + \tau \gamma_2.$$

Similarly to the above, we can consider a one-parameter family of the quadratic algebras by replacing $S_0 \rightarrow S_0 + \lambda$. Note that it degenerates to the standard Poisson brackets on T^2 in the limit $\lambda \rightarrow \infty$. This allows us to define the bihamiltonian structure for the hierarchy of the modified hydrodynamics. In particular, Eq. (5.24) can be rewritten as

$$\partial_t \mathbf{S} = \{S_0, \mathbf{S}\}_2.$$

Such a bihamiltonian structure is a curious feature in the modified hydrodynamics, emphasizing its drastic difference from the classical hydrodynamics.

6. Quantum Counterparts

In this section we present two associative algebras “quantizing” the two Poisson algebras discussed above in the case of the NCT. It is easy with the linear bracket. Indeed, the quantization of the Lie-Poisson algebra on \sin_θ^* leads to the universal enveloping algebra of \sin_θ .

Replace the classical variables \mathbf{S} on the NCT \mathcal{A}_θ by non-commuting variables $\hat{\mathbf{S}}$. Now we consider the quantization of the quadratic Poisson algebra $P_{\theta,\epsilon,\tau}$ and construct the associative algebra $\hat{P}_{\theta,\epsilon,\tau,\hbar}$, where $\hbar \in E_\tau$ is the deformation parameter. For this we introduce the quantum R -matrix related to the group SIN_θ . The quantum Yang-Baxter equation defined on $GL(N, \mathbb{C})$ [13, 14] is generalized to SIN_θ in the following way. The quantum R -matrix on $SIN_\theta \otimes SIN_\theta$ assumes the form

$$R(z, w) = \sum_c \varphi_{\theta,\epsilon}(c, z - w|\hbar) T_c \otimes T_{-c},$$

where T_c is the basis of SIN_θ and the main ingredient in our construction is the function

$$\varphi_{\theta,\epsilon}(c, z|\hbar) = \mathbf{e}_\theta(\epsilon_2 c_2 z) \phi((\epsilon_1 c_1 + \tau \epsilon_2 c_2) \theta + \hbar, z), \quad \hbar \in E_\tau.$$

Note that, in contrast with the classical r -matrix (4.11), there is an additional term $\varphi_{\theta,\epsilon}(0, z - w|\hbar) T_0 \otimes T_0$.

Proposition 6.1. *The matrix R satisfies the quantum YB equation*

$$R_{12}(z - w) R_{13}(z) R_{23}(w) = R_{23}(w) R_{13}(z) R_{12}(z - w). \quad (6.1)$$

Proof. Consider the coefficient in front of $T_a \otimes T_b \otimes T_c$ in the l.h.s of (6.1). It vanishes if $a + b + c \neq 0$. Therefore (6.1) implies

$$\begin{aligned} & \mathbf{e}_\theta(a \times b + 2b \times c) \varphi_{\theta,\epsilon}(c, z - w|\hbar) \varphi_{\theta,\epsilon}(a - c, z|\hbar) \varphi_{\theta,\epsilon}(b + c, w|\hbar) \\ & - \mathbf{e}_\theta(a \times b + 2b \times f) \varphi_{\theta,\epsilon}(f, z - w|\hbar) \varphi_{\theta,\epsilon}(a - f, z|\hbar) \varphi_{\theta,\epsilon}(b + f, w|\hbar) = 0 \end{aligned}$$

Here a, b are fixed and c, f are arbitrary elements of the lattice $\mathbb{Z}_{\theta,\epsilon}(\tau)$ (C.16). Choose f in the form $f = a - b - c$ to obtain

$$\begin{aligned} & \mathbf{e}_\theta(a \times b + 2b \times c) (\varphi_{\theta,\epsilon}(c, z - w|\hbar) \varphi_{\theta,\epsilon}(a - c, z|\hbar) \varphi_{\theta,\epsilon}(b + c, w|\hbar) \\ & - \varphi_{\theta,\epsilon}(a - b - c, z - w|\hbar) \varphi_{\theta,\epsilon}(a - c, w|\hbar) \varphi_{\theta,\epsilon}(b + c, z|\hbar)) = 0. \end{aligned}$$

This equality can be transformed to the form

$$\varphi_{\theta,\epsilon}(a + 2\hbar, z|\hbar) \varphi_{\theta,\epsilon}(b, w|\hbar) \mathbf{e}_\theta(a \times b + 2b \times c)$$

$$\times (\zeta_{\theta,\epsilon}(c + \hbar) - \zeta_{\theta,\epsilon}(a - b - c + \hbar) + \zeta_{\theta,\epsilon}(a - c + \hbar) - \zeta_{\theta,\epsilon}(b + c + \hbar)) = 0$$

by means of (A.27). If $b = 0$ then the l.h.s. vanishes and we come to (6.1). Let $b \neq 0$ and consider the shift $c \rightarrow c + jb$, where $j \in \mathbb{Z}$. The shift does not change the exponential factor. Take the sum over the orbit generated by the shifts

$$\sum_{j \in \mathbb{Z}} \zeta_{\theta,\epsilon}(c + jb + \hbar) - \zeta_{\theta,\epsilon}(a - b - jb - c + \hbar) + \zeta_{\theta,\epsilon}(a - c + jb + \hbar) - \zeta_{\theta,\epsilon}(b + c + jb + \hbar).$$

One can see that the neighboring terms in the series vanish and we come to (6.1) for an arbitrary b . \square

This proposition allows us to define the associative algebra $\hat{P}_{\theta,\epsilon,\tau,\hbar}$ by the relation

$$R(z - w) L_1^{\bar{\hbar}}(z) L_2^{\bar{\hbar}}(w) = L_2^{\bar{\hbar}}(w) L_1^{\bar{\hbar}}(z) R(z - w), \quad (6.2)$$

where

$$L^{\bar{\hbar}}(z) = \hat{S}_0 \varphi_{\theta,\epsilon}(0, z|\hbar) T_0 + \sum_{\alpha} \hat{S}_{\alpha} \varphi_{\theta,\epsilon}(\alpha, z|\hbar) T_{\alpha}.$$

In order to obtain the relations in $\hat{P}_{\theta,\epsilon,\tau,\hbar}$ one should exclude the spectral parameters z, w from (6.2).

Lemma 6.1. *The relations in the associative algebra $\hat{P}_{\theta, \tau, \hbar}$ assume the form*

$$\sum_c \hat{S}_{a-c} \hat{S}_{b+c} \mathbf{e}_\theta(c \times (a-b)) \mathbf{f}_{\theta, \epsilon}(a, b, c|\hbar) = 0, \quad \text{for any } a, b \in \mathbb{Z} \oplus \mathbb{Z}, \quad (6.3)$$

where

$$\begin{aligned} \mathbf{f}_{\theta, \epsilon}(a, b, c|\hbar) &= \zeta_{\theta, \epsilon}(c + \hbar) - \zeta_{\theta, \epsilon}(a - b - c + \hbar) - \zeta_{\theta, \epsilon}(b + c + \hbar) \\ &\quad + \zeta_{\theta, \epsilon}(a - c + \hbar) \\ &= - \frac{\vartheta'(0) \vartheta_{\theta, \epsilon}(a + 2\hbar) \vartheta_{\theta, \epsilon}(b) \vartheta_{\theta, \epsilon}(b - a + 2c)}{\vartheta_{\theta, \epsilon}(a - c + \hbar) \vartheta_{\theta, \epsilon}(b + c + \hbar) \vartheta_{\theta, \epsilon}(b - a + c + \hbar) \vartheta_{\theta, \epsilon}(c + \hbar)}, \end{aligned} \quad (6.4)$$

and $\vartheta_{\theta, \epsilon}, \zeta_{\theta, \epsilon}(\cdot)$ are the constants (C.17), (C.18).

Proof. Consider in (6.2) the matrix element $T_a \otimes T_b$. We come to the relation

$$\begin{aligned} \sum_c \hat{S}_{a-c} \hat{S}_{b+c} \{ \varphi_{\theta, \epsilon}(c, z - w|\hbar) \varphi_{\theta, \epsilon}(a - c, z|\hbar) \varphi_{\theta, \epsilon}(b + c, w|\hbar) \\ - \varphi_{\theta, \epsilon}(a - b - c, z - w|\hbar) \varphi_{\theta, \epsilon}(a - c, w|\hbar) \varphi_{\theta, \epsilon}(b + c, z|\hbar) \} \\ \times \mathbf{e}_\theta(c \times (a - b)) = 0. \end{aligned} \quad (6.5)$$

The expression in the brackets $\{ \}$ is equal to

$$\begin{aligned} \mathbf{e}_\theta((a_2 \epsilon_2 z + b_2 \epsilon_2 w) \{ \phi_{\theta, \epsilon}(c, z - w|\hbar) \phi_{\theta, \epsilon}(a - c, z|\hbar) \phi_{\theta, \epsilon}(b + c, w|\hbar) \\ - \phi_{\theta, \epsilon}(a - b - c, z - w|\hbar) \phi_{\theta, \epsilon}(a - c, w|\hbar) \phi_{\theta, \epsilon}(b + c, z|\hbar) \}), \end{aligned}$$

where $\phi_{\theta, \epsilon}(c, z|\hbar) = \phi((\epsilon_1 c_1 + \tau \epsilon_2 c_2) \theta + \hbar, z)$. Finally, by using (A.27) we come from (6.5) to (6.4). \square

We rewrite (6.3) in the form most resembling the original Sklyanin relations [14]. Let $\mathfrak{A}(\alpha, \beta)$ (respectively, $\mathfrak{S}(\alpha, \beta)$) be the operator of antisymmetrization (respectively, symmetrization) with respect to the permutations of two indices (α, β) .

Proposition 6.2. *The relations (6.4) are equivalent to the commutator relations*

$$[\hat{S}_0, \hat{S}_b] = - \sum_{\gamma \neq -b} \hat{S}_{-\gamma} \hat{S}_{b+\gamma} \mathbf{e}_\theta(b \times \gamma) \frac{\mathbf{f}_{\theta, \epsilon}(0, b, \gamma|\hbar)}{\mathbf{f}_{\theta, \epsilon}(0, b, 0|\hbar)}, \quad (6.6)$$

$$\begin{aligned} [\hat{S}_\alpha, \hat{S}_\beta] &= \mathfrak{A}(\alpha, \beta) \left\{ \left(\hat{S}_{\alpha+\beta} \hat{S}_0 \mathbf{e}_\theta(-\beta \times \alpha) - \hat{S}_0 \hat{S}_{\alpha+\beta} \mathbf{e}_\theta(\beta \times \alpha) \right) \frac{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, \alpha|\hbar)}{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, 0|\hbar)} \right. \\ &\quad + \sum_{\gamma \neq \alpha, -\beta} \left(\hat{S}_{\alpha-\gamma} \hat{S}_{\beta+\gamma} \mathbf{e}_\theta(\gamma \times (\alpha - \beta)) - \hat{S}_{\beta+\gamma} \hat{S}_{\alpha-\gamma} \mathbf{e}_\theta(-\gamma \times (\alpha - \beta)) \right) \\ &\quad \times \frac{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, \gamma|\hbar)}{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, 0|\hbar)} \left. \right\}, \end{aligned} \quad (6.7)$$

where $\mathbf{C}_\theta(\alpha, \beta)$ are $\mathfrak{sl}(N, \mathbb{C})$ structure constants (B.8), and

$$\begin{aligned} \mathfrak{S}(\alpha, \beta) \left\{ \left(\hat{S}_{\alpha+\beta} \hat{S}_0 \mathbf{e}_\theta(-\beta \times \alpha) - \hat{S}_0 \hat{S}_{\alpha+\beta} \mathbf{e}_\theta(\beta \times \alpha) \right) \frac{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, \alpha|\hbar)}{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, 0|\hbar)} \right. \\ + \sum_{\gamma \neq \alpha, -\beta} \left(\hat{S}_{\alpha-\gamma} \hat{S}_{\beta+\gamma} \mathbf{e}_\theta(\gamma \times (\alpha - \beta)) - \hat{S}_{\beta+\gamma} \hat{S}_{\alpha-\gamma} \mathbf{e}_\theta(-\gamma \times (\alpha - \beta)) \right) \\ \times \frac{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, \gamma|\hbar)}{\mathbf{f}_{\theta, \epsilon}(\alpha, \beta, 0|\hbar)} \left. \right\} = 0. \end{aligned}$$

Proof. The relation (6.3) can be rewritten in the form

$$\sum_c \left(\hat{S}_{a-c} \hat{S}_{b+c} \mathbf{e}_\theta(c \times (a-b)) - \hat{S}_{b+c} \hat{S}_{a-c} \mathbf{e}_\theta(c \times (a-b)) \right) \mathbf{f}_{\theta,\epsilon}(a, b, c|\hbar) = 0,$$

due to the equality $\mathbf{f}_{\theta,\epsilon}(a, b, c|\hbar) = -\mathbf{f}_{\theta,\epsilon}(a, b, a-b-c|\hbar)$. For the case $a = 0$ this relation assumes the form (6.6). To come to (6.7) we single out the two terms with $c = 0$ and put them in the left-hand side. In the right-hand side we first write down the terms with $c = a$ and $c = -b$ and obtain the required relations. \square

It would be interesting to describe a quantum version of the whole bihamiltonian structure for elliptic rotators on \sin^* or $Ham(T^2)$. Posing the problem more generally, the bihamiltonian recursion procedure of generating the conserved quantities from a Casimir function for a linear family of Poisson structures (see [19]) might have a quantum analog as an expansion of a central element for a linear family of associative algebras. The latter seems to be a very strong requirement on a pair of associative algebras and it would be very interesting to find any non-trivial example of this kind. Namely, one is looking for a pair of associative algebras, such that their mixed associator satisfies some consistency condition. This would provide the most straightforward quantization for a system with a bihamiltonian structure.

Acknowledgements. We are grateful for hospitality to the Max-Planck-Institut für Mathematik in Bonn, where most of this work was done. B.K. is also grateful for hospitality to the University of Nice and IHES in Bures-sur-Yvette. The work of B.K. was partially supported by an NSERC research grant. The work of A.L. and M.O. was supported in part by grants RFBR-03-02-17554, NSCh-1999-2003.2 and CRDF RM1-2545.

Appendices

Appendix A. Elliptic functions. Here we summarize the main formulae for elliptic functions. Consider an elliptic curve

$$E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), \quad q = \mathbf{e}(\tau) = \exp(2\pi i \tau). \quad (\text{A.1})$$

The basic element in our consideration is the theta function:

$$\begin{aligned} \vartheta(z; \tau) &= q^{\frac{1}{8}} \sum_{n \in \mathbb{Z}} (-1)^n e^{\pi i (n(n+1)\tau + 2nz)} \\ &= q^{\frac{1}{8}} e^{-\frac{i\pi}{4}} (e^{i\pi z} - e^{-i\pi z}) \prod_{n=1}^{\infty} (1 - q^n)(1 - q^n e^{2i\pi z}) \\ &\quad \times (1 - q^n e^{-2i\pi z}). \end{aligned} \quad (\text{A.2})$$

1. The Weierstrass functions

$$\sigma(z; \tau) = \exp(\eta z^2) \frac{\vartheta(z; \tau)}{\vartheta'(0; \tau)}, \quad (\text{A.3})$$

where

$$\eta(\tau) = -\frac{1}{6} \frac{\vartheta'''(0; \tau)}{\vartheta'(0; \tau)}, \quad (\text{A.4})$$

$$\zeta(z; \tau) = \partial_z \log \vartheta(z; \tau) + 2\eta(\tau)z, \quad \zeta(z; \tau) \sim \frac{1}{z} + O(z^3), \quad (\text{A.5})$$

$$\wp(z; \tau) = -\partial_z \zeta(z; \tau), \quad (\text{A.6})$$

$$\wp(u; \tau) = \frac{1}{u^2} + \sum'_{j,k} \left(\frac{1}{(j+k\tau+u)^2} - \frac{1}{(j+k\tau)^2} \right). \quad (\text{A.7})$$

2. Function ϕ

$$\phi(u, z) = \frac{\vartheta(u+z)\vartheta'(0)}{\vartheta(u)\vartheta(z)}. \quad (\text{A.8})$$

It has a pole at $z = 0$ and the expansion

$$\phi(u, z) = \frac{1}{z} + \zeta(u|\tau) + 2\eta(\tau)z + \frac{z}{2}((\zeta(u; \tau) + 2\eta(\tau)z)^2 - \wp(u)) + \dots, \quad (\text{A.9})$$

as $z \rightarrow 0$. Furthermore,

$$\phi(u, z)^{-1} \partial_u \phi(u, z) = \zeta(u+z) - \zeta(u) + 2\eta(\tau)z, \quad (\text{A.10})$$

$$\phi(u, z) = \exp(-2\eta uz) \frac{\sigma(u+z)}{\sigma(u)\sigma(z)}, \quad (\text{A.11})$$

$$\phi(u, z)\phi(-u, z) = \wp(z) - \wp(u). \quad (\text{A.12})$$

3. Quasi-periodicity

$$\vartheta(z+1; \tau) = -\vartheta(z; \tau), \quad \vartheta(z+\tau; \tau) = -\mathbf{e}\left(-\frac{1}{2}\tau - z\right)\vartheta(z; \tau), \quad (\text{A.13})$$

$$\zeta(z+1; \tau) = \zeta(z; \tau) - 2\eta, \quad \zeta(z+\tau; \tau) = \zeta(z; \tau) - 2(\pi i + \eta\tau), \quad (\text{A.14})$$

$$\wp(u+1; \tau) = \wp(u+\tau; \tau) = \wp(u; \tau), \quad (\text{A.15})$$

$$\phi(u+1, z) = \phi(u, z), \quad \phi(u+\tau, z) = \mathbf{e}(-z)\phi(u, z). \quad (\text{A.16})$$

4. Parity

$$\vartheta(-z; \tau) = -\vartheta(z; \tau), \quad (\text{A.17})$$

$$\zeta(-z; \tau) = -\zeta(z; \tau), \quad (\text{A.18})$$

$$\wp(-u; \tau) = \wp(u; \tau), \quad (\text{A.19})$$

$$\phi(-u, -z) = -\phi(u, z). \quad (\text{A.20})$$

5. The Fay three-section formula

$$\begin{aligned} & \phi(u_1, z_1)\phi(u_2, z_2) - \phi(u_1 + u_2, z_1)\phi(u_2, z_2 - z_1) \\ & - \phi(u_1 + u_2, z_2)\phi(u_1, z_1 - z_2) = 0. \end{aligned} \quad (\text{A.21})$$

Particular cases of this formula are (A.12), the Calogero functional equation

$$\phi(u, z)\partial_v\phi(v, z) - \phi(v, z)\partial_u\phi(u, z) = (\wp(v) - \wp(u))\phi(u + v, z), \quad (\text{A.22})$$

and

$$\begin{aligned} & \phi(u_1, z)\phi(u_2, z) - \phi(u_1 + u_2, z)(\zeta(u_1) + \zeta(u_2) - 2\eta(\tau)(u_1 + u_2)) \\ & + \partial_z\phi(u_1 + u_2, z) = 0. \end{aligned} \quad (\text{A.23})$$

For $u_1 + u_2 + u_3 = 0$,

$$\begin{aligned} & \phi(u_1, z)\phi(u_2, z)\phi(u_3, z) = [\wp(z) - \wp(u_3)][\zeta(u_1) + \zeta(u_2) \\ & + \zeta(u_3 - z) + \zeta(z)], \end{aligned} \quad (\text{A.24})$$

as follows from (A.10), (A.12), and (A.23). Then

$$\begin{aligned} & \phi(u_1, z)\phi(u_2, z)\phi(u_3, z)|_{z \rightarrow 0} = \frac{1}{z^3} + \frac{1}{z^2} [\zeta(u_1) + \zeta(u_2) + \zeta(u_3)] \\ & - \frac{1}{2}\wp'(u_3) - \wp(u_3) [\zeta(u_1) + \zeta(u_2) + \zeta(u_3)] + O(z). \end{aligned} \quad (\text{A.25})$$

From (A.21) and (A.10) we have

$$\phi(u_1, z)\phi(u - u_1, z) = \phi(u, z)(\zeta(u_1) + \zeta(u - u_1) - \zeta(u + z) + \zeta(z)). \quad (\text{A.26})$$

Another important relation is

$$\begin{aligned} & \phi(v, z - w)\phi(u_1 - v, z)\phi(u_2 + v, w) - \phi(u_1 - u_2 - v, z - w) \\ & \times \phi(u_2 + v, z)\phi(u_1 - v, w) \\ & = \phi(u_1, z)\phi(u_2, w)f(u_1, u_2, v), \end{aligned} \quad (\text{A.27})$$

where

$$f(u_1, u_2, v) = \zeta(v) - \zeta(u_1 - u_2 - v) + \zeta(u_1 - v) - \zeta(u_2 + v). \quad (\text{A.28})$$

One can rewrite the last function as

$$f(u_1, u_2, v) = -\frac{\vartheta'(0)\vartheta(u_1)\vartheta(u_2)\vartheta(u_2 - u_1 + 2v)}{\vartheta(u_1 - v)\vartheta(u_2 + v)\vartheta(u_2 - u_1 + v)\vartheta(v)}. \quad (\text{A.29})$$

To prove (A.27) one should consider the expression

$$\frac{\phi(v, z - w)\phi(u_1 - v, z)\phi(u_2 + v, w) - \phi(u_1 - u_2 - v, z - w)\phi(u_2 + v, z)\phi(u_1 - v, w)}{\phi(u_1, z)\phi(u_2, w)}.$$

It is a doubly periodic function in z and w without poles. Therefore, it is a constant, which depends on u_1, u_2, v . This constant is equal to $f(u_1, u_2, v)$ (A.28). It is easy to check that the elliptic functions (A.28) and (A.29) coincide.

A particular case of (A.27), which corresponds to the case $v = u_1$ (or $v = -u_2$), is the Fay identity (A.21). Another particular case comes from $u_2 = 0$ (or $u_1 = 0$):

$$\begin{aligned} & \phi(v, z - w)\phi(u - v, z)\phi(v, w) - \phi(u - v, z - w)\phi(v, z)\phi(u - v, w) \\ & = \phi(u_1, z)(\wp(v) - \wp(u - v)). \end{aligned} \quad (\text{A.30})$$

Appendix B. Elliptic constants related to $\mathfrak{gl}(N, \mathbb{C})$. Let $1 \leq k < N$ be a coprime number with respect to N and set $\theta = \frac{k}{N}$. Define

$$\mathbf{e}_\theta(z) = \exp(2\pi i \theta z), \quad (\text{B.1})$$

$$Q = \text{diag}(\mathbf{e}_\theta(1), \dots, \mathbf{e}_\theta(m), \dots, 1), \quad (\text{B.2})$$

$$\Lambda = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ 1 & 0 & 0 & \dots & 0 \end{pmatrix}. \quad (\text{B.3})$$

Consider a finite two-dimensional lattice of order N^2 ,

$$\mathbb{Z}_N^{(2)} = \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}/N\mathbb{Z}.$$

The matrices

$$T_a = \frac{1}{2\pi i \theta} \mathbf{e}_\theta \left(\frac{a_1 a_2}{2} \right) Q^{a_1} \Lambda^{a_2}, \quad a = (a_1, a_2) \in \mathbb{Z}^{(2)} \quad (\text{B.4})$$

generate the basis in $\mathfrak{gl}(N, \mathbb{C})$. We use the Greek letters for the elements of the lattice

$$\tilde{\mathbb{Z}}_N^{(2)} = \mathbb{Z}_N^{(2)} \setminus (0, 0). \quad (\text{B.5})$$

Thus, $\{T_\alpha\}$ define a basis in $\mathfrak{sl}(N, \mathbb{C})$. Since

$$T_a T_b = \frac{1}{2\pi \theta} \mathbf{e}_\theta \left(-\frac{a \times b}{2} \right) T_{a+b}, \quad (a \times b = a_1 b_2 - a_2 b_1) \quad (\text{B.6})$$

the commutation relations in this basis assume the form

$$[T_\alpha, T_\beta] = \mathbf{C}_\theta(\alpha, \beta) T_{\alpha+\beta}, \quad (\text{B.7})$$

where

$$\mathbf{C}_\theta(\alpha, \beta) = \frac{1}{\pi \theta} \sin \pi \theta (\alpha \times \beta). \quad (\text{B.8})$$

Let

$$\mathbb{Z}_\theta^{(2)}(\tau) = (\gamma_1 + \gamma_2 \tau) \theta, \quad \gamma \in \tilde{\mathbb{Z}}_N^{(2)} \quad (\text{B.9})$$

be a regular lattice of order $N^2 - 1$ on E_τ . Introduce the following constants on $\mathbb{Z}_\theta^{(2)}(\tau)$:

$$\vartheta_\theta(\gamma) = \vartheta((\gamma_1 + \gamma_2 \tau) \theta), \quad (\text{B.10})$$

$$\zeta_\theta(\gamma) = \zeta((\gamma_1 + \gamma_2 \tau) \theta), \quad \wp_\theta(\gamma) = \wp((\gamma_1 + \gamma_2 \tau) \theta), \quad (\text{B.11})$$

and the quasi-periodic functions on E_τ ,

$$\phi_\theta(\gamma, z) = \phi((\gamma_1 + \gamma_2 \tau) \theta, z), \quad (\text{B.12})$$

$$\varphi_\theta(\gamma, z) = \mathbf{e}_\theta(\gamma_2 z) \phi_\theta(\gamma, z), \quad (\text{B.13})$$

$$f_\theta(\gamma, z) = \mathbf{e}_\theta(\gamma_2 z) \partial_u \phi(u, z)|_{u=(\gamma_1 + \gamma_2 \tau) \theta}. \quad (\text{B.14})$$

It follows from (A.8) that

$$\varphi_\theta(\gamma, z + 1) = \mathbf{e}_\theta(\gamma_2) \varphi_\theta(\gamma, z), \quad \varphi_\theta(\gamma, z + \tau) = \mathbf{e}_\theta(-\gamma_1) \varphi_\theta(\gamma, z). \quad (\text{B.15})$$

Appendix C. Non-commutative torus.

1. *Definition and representation.* The non-commutative torus \mathcal{A}_θ is a unital algebra with the two generators (U_1, U_2) that satisfy the relation

$$U_1 U_2 = \mathbf{e}_\theta^{-1} U_2 U_1, \quad \mathbf{e}_\theta = e^{2\pi i \theta}, \quad \theta \in [0, 1). \quad (\text{C.1})$$

Elements of \mathcal{A}_θ are the double sums

$$\mathcal{A}_\theta = \left\{ X = \sum_{a_1, a_2 \in \mathbb{Z}} c_{a_1, a_2} U_1^{a_1} U_2^{a_2} \mid c_{a_1, a_2} \in \mathbb{C} \right\},$$

where c_{a_1, a_2} are elements of the ring \mathfrak{S} of rapidly decreasing sequences on \mathbb{Z}^2 ,

$$\mathfrak{S} = \{c_{a_1, a_2} \mid \sup_{a_1, a_2 \in \mathbb{Z}} (1 + a_1^2 + a_2^2)^k |c_{a_1, a_2}|^2 < \infty \text{ for all } k \in \mathbb{N}\}. \quad (\text{C.2})$$

The trace functional $\text{tr}(x)$ on \mathcal{A}_θ is defined by

$$\text{tr}(X) = c_{00}. \quad (\text{C.3})$$

It satisfies the evident identities

$$\text{tr}(1) = 1, \quad \text{tr}(XY) = \text{tr}(YX).$$

The dual space to \mathfrak{S} is the space

$$\mathfrak{S}' = \{S_{a_1, a_2} \mid \sum_{a_1, a_2} c_{a_1, a_2} S_{-a_1, -a_2} < \infty, \quad c_{a_1, a_2} \in \mathfrak{S}\}. \quad (\text{C.4})$$

The relation with the commutative algebra of smooth functions on the two-dimensional torus

$$T^2 = \{\mathbb{R}^2 / \mathbb{Z} \oplus \mathbb{Z}\} \sim \{0 < x_1 \leq 1, 0 < x_2 \leq 1\} \quad (\text{C.5})$$

comes from the identification

$$U_1 \rightarrow \mathbf{e}(x), \quad U_2 \rightarrow \mathbf{e}(y), \quad (\text{C.6})$$

while the multiplication on T^2 becomes the Moyal multiplication:

$$(f * g)(x) := fg + \sum_{n=1}^{\infty} \frac{(\pi \theta)^n}{n!} \varepsilon_{r_1 s_1} \dots \varepsilon_{r_n s_n} (\partial_{r_1 \dots r_n}^n f)(\partial_{s_1 \dots s_n}^n g). \quad (\text{C.7})$$

The trace functional (C.3) in the Moyal identification is the integral

$$\text{tr} f = -\frac{1}{4\pi^2} \int_{\mathcal{A}_\theta} f dx_1 dx_2 = f_{00}. \quad (\text{C.8})$$

We can identify U_1, U_2 with matrices from $\text{GL}(\infty)$. Define $\text{GL}(\infty)$ as the associative algebra of infinite matrices $c_{jk} E_{jk}$, where $E_{jk} = ||\delta_{jk}||$, such that

$$\sup_{j, k \in \mathbb{Z}} |c_{jk}|^2 |j - k|^n < \infty \text{ for } n \in \mathbb{N}.$$

Consider the following two matrices from $GL(\infty)$:

$$Q = \text{diag}(\mathbf{e}(j\theta)) \text{ and } \Lambda = \|\delta_{j,j+1}\|, \quad j \in \mathbb{Z}.$$

We have the following identification:

$$U_1 \rightarrow Q, \quad U_2 \rightarrow \Lambda. \quad (\text{C.9})$$

Another useful realization of \mathcal{A}_θ in the Schwartz space on \mathbb{R} by the operators

$$U_1 f(x) = f(x - \theta), \quad U_2 f(x) = \exp(2\pi i x) f(x). \quad (\text{C.10})$$

2. *sin-algebra*. Define the following quadratic combinations of the generators

$$T_\alpha = \frac{i}{2\pi\theta} \mathbf{e}\left(\frac{\alpha_1\alpha_2}{2}\theta\right) U_1^m U_2^n, \quad \alpha \in \tilde{\mathbb{Z}}^{(2)}, \quad (\text{C.11})$$

$$\tilde{\mathbb{Z}}^{(2)} = \{\alpha = (\alpha_1, \alpha_2), \alpha_j \in \mathbb{Z}, \alpha \neq (0, 0)\}.$$

Their commutator has the form of the sin-algebra

$$[T_\alpha, T_\beta] = \mathbf{C}_\theta(\alpha, \beta) T_{\alpha+\beta}, \quad (\text{C.12})$$

where

$$\mathbf{C}_\theta(\alpha, \beta) = \frac{1}{\pi\theta} \sin \pi\theta(\alpha \times \beta). \quad (\text{C.13})$$

We denote by \sin_θ the Lie algebra with the generators T_α over the ring \mathfrak{S} (C.2),

$$\psi = \sum_{\alpha} \psi_{\alpha} T_{\alpha}, \quad \psi_{\alpha} \in \mathfrak{S}, \quad (\text{C.14})$$

and by SIN_θ the group of invertible elements from \mathcal{A}_θ . The coalgebra \sin_θ^* is the linear space

$$\sin_\theta^* = \left\{ \mathbf{S} = \sum_{\alpha \in \tilde{\mathbb{Z}}^{(2)}} s_{\alpha} T_{\alpha}, \quad s_{\alpha} \in \mathfrak{S}' \right\}.$$

In the Moyal representation (C.7) the commutator of \sin_θ has the form

$$[f(x, y), g(x, y)] = \{f, g\}^\theta := \frac{1}{\theta} (f * g - g * f). \quad (\text{C.15})$$

3. *Elliptic constants related to NCT \mathcal{A}_θ* . Introduce two numbers $\epsilon = (\epsilon_1, \epsilon_2)$ such that $\epsilon_a\theta < 1$ and $\epsilon_a\theta$ are irrational. Consider the dense set $\mathbb{Z}_{\theta,\epsilon}(\tau)$ in E_τ :

$$\mathbb{Z}_{\theta,\epsilon}(\tau) = \{(\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2)\theta \in E_\tau \mid (\gamma_1, \gamma_2) \in \tilde{\mathbb{Z}}^{(2)}\}. \quad (\text{C.16})$$

The corresponding elliptic functions with the arguments from $\mathbb{Z}_{\theta,\epsilon}(\tau)$ are as follows:

$$\vartheta_{\theta,\epsilon}(\gamma) = \vartheta((\epsilon_1\gamma_1 + \epsilon_2\gamma_2\tau)\theta), \quad (\text{C.17})$$

$$\zeta_{\theta,\epsilon}(\gamma) = \zeta((\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2)\theta), \quad \wp_{\theta,\epsilon}(\gamma) = \wp((\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2)\theta), \quad (\text{C.18})$$

$$\phi_{\theta,\epsilon}(\gamma, z) = \phi(-(\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2)\theta, z), \quad (\text{C.19})$$

$$\varphi_{\theta,\epsilon}(\gamma) = \mathbf{e}_\theta(\epsilon_2\gamma_2 z) \phi_{\theta,\epsilon}(\gamma, z), \quad (\text{C.20})$$

$$f_{\theta,\epsilon}(\gamma) = \mathbf{e}_\theta(\epsilon_2\gamma_2 z) \partial_u \phi(u, z)|_{u=(\epsilon_1\gamma_1 + \tau\epsilon_2\gamma_2)\theta}. \quad (\text{C.21})$$

4. *Dispersionless limit.* In the limit $\theta \rightarrow 0$ the Lie algebra \sin_θ becomes the Lie algebra of Hamiltonian functions

$$Ham(T^2) \sim C^\infty(T^2)/\mathbb{C} \quad (C.22)$$

equipped with the canonical Poisson brackets. In $Ham(T^2)$ we have the Fourier basis

$$\mathbf{e}(\alpha \cdot x) = \exp(2\pi i(\alpha_1 x_1 + \alpha_2 x_2)) \quad (C.23)$$

instead of the basis (C.11). The commutator (C.12) becomes

$$[\mathbf{e}(\alpha x), \mathbf{e}(\beta x)] = (\alpha \times \beta) \mathbf{e}((\alpha + \beta) \cdot x).$$

The algebra $Ham(T^2)$ (without constant Hamiltonians) is isomorphic to the Lie algebra $SVect_0(T^2)$ of the divergence-free *zero-flux* vector fields on T^2 equipped with the area form $-4\pi^2 dx_1 dx_2$. Let $h(x_1, x_2) \in Ham(T^2)$. Then the Hamiltonian field V_h corresponding to the Hamiltonian function h is

$$V_h = -\frac{1}{4\pi^2}((\partial_2 h)\partial_1 - (\partial_1 h)\partial_2), \quad (C.24)$$

while

$$[V_h, V_{h'}] = V_{\{h, h'\}}. \quad (C.25)$$

For $f(x) = \sum_\alpha f_\alpha \mathbf{e}(\alpha \cdot x)$

$$\int_{T^2} f = -\frac{1}{4\pi^2} f_{00}. \quad (C.26)$$

References

1. Ziglin, S.: Non-integrability of the problem of the motion of four point vortices. Dokl. Akad. Nauk SSSR **250**(6), 1296–1300 (1980)
2. Reyman, A.G., Semenov-Tyan-Shanskii, M.A.: Lie algebras and Lax equations with spectral parameter on elliptic curve. Notes of the LOMI Seminar **150**, 104–118 (1986)
3. Levin, A., Olshanetsky, M., Zotov, A.: *Hitchin Systems: symplectic Hecke correspondence and two-dimensional version*. Commun. Math. Phys. **236**(1), 93–133 (2003)
4. Belavin, A., Drinfeld, V.: Solutions of the classical Yang-Baxter equation for simple Lie algebras. Funct. Anal Appl. **16**(3), 1–29 (1982)
5. Feigin, B., Odesski, A.: Sklyanin's elliptic algebras. Funct. Anal. Appl. **23**(3), 207–214 (1989)
6. Arnold, V.I.: The Hamiltonian nature of the Euler equation in the dynamics of rigid body and of an ideal fluid. Russ. Math. Surv. **24**(3), 225–226 (1969)
7. Arnold, V.I., Khesin, B.A.: *Topological methods in Hydrodynamics*. Applied Mathematical Science **125**, New York: Springer-Verlag, 1998
8. Manakov, S.: A remark on the integration of the Eulerian equations of the dynamics of an n -dimensional rigid body. Funct. Anal. Appl. **10**(4), 93–94 (1976)
9. Ward, R.S.: Infinite-dimensional gauge groups and special nonlinear gravitons. J. Geom. Phys. **8**(1–4), 317–325 (1992)
10. Khesin, B., Ovsienko, V.: The super Korteweg-de Vries equation as an Euler equation. Funct. Anal. Appl. **21**(4), 81–82 (1987)
11. Segal, G.: The geometry of the KdV equation. In: *Topological methods in quantum field theory (Trieste, 1990)*. Internat. J. Modern Phys. **A6**(16), 2859–2869 (1991)
12. Khesin, B., Misiolek, G.: Euler equations on homogeneous spaces and virasoro orbits. Adv. Math. **176**, 116–144 (2003)
13. Belavin, A.: Discrete groups and integrability of quantum systems. Funct. Anal. Appl. **14**, 18–26 (1980)

14. Sklyanin, E.K.: Some algebraic structures connected with the Yang-Baxter equation. *Funct. Anal. Appl.* **16**(4), 27–34 (1982)
15. Olshanetsky, M.: Integrable tops and non-commutative torus. In: *Proc. of Int. Workshop Supersymmetries and Quantum Symmetries*, Sept. 2001, Karpacz, Poland, <http://arxiv.org/abs/nlin.SI/0203003>, 2002
16. Etingof, P.I., Frenkel, I.B.: Central extensions of current groups in two dimensions. *Commun. Math. Phys.* **165**(3), 429–444 (1994)
17. Semenov-Tyan-Shansky, M.: What a classical r -matrix is. *Funct. Anal. Appl.* **17**(4), 17–33 (1983)
18. Braden, H.W., Dolgushev, V.A., Olshanetsky, M.A., Zotov, A.V.: Classical R-Matrices and the Feigin-Odesskii Algebra via Hamiltonian and Poisson Reductions. *J. Phys.* **A36**, 6979–7000 (2003)
19. Magri, F.: A simple model of the integrable Hamiltonian equation. *J. Math. Phys.* **19**, 1156–1162 (1978)

Communicated by L. Takhtajan