

Tutorial 0201 (Thursday, 4-5), Quiz 2, rubric.

It can be shown that on the set $U = \mathbf{R}^2 \setminus \{(x, 0) | x \geq 0\}$, i.e., the plane with the positive x axis and origin removed, the function

$$P(x, y) = 2\sqrt{(x^2 + y^2)^{\frac{1}{2}} - x}$$

is harmonic. The formula

$$Q(x_0, y_0) = \int_{(1,0)}^{(x_0, y_0)} -\frac{\partial P}{\partial y} dx + \frac{\partial P}{\partial x} dy$$

gives a function which is harmonic on U and such that

$$f(x + iy) = P(x, y) + iQ(x, y)$$

is analytic on U . Use this formula to determine Q on the part of the unit circle which lies in U . (The identity

$$\sin \frac{x}{2} = \sqrt{\frac{1 - \cos x}{2}},$$

valid when $\sin \frac{x}{2}$ is nonnegative, may be useful.) What is

$$\lim_{(x,y) \rightarrow (1,0^+)} Q(x, y) - \lim_{(x,y) \rightarrow (1,0^-)} Q(x, y)$$

(where the limits are taken along the unit circle)? What do we call the half-line $\{(x, 0) | x \geq 0\}$ in this case?

First, we note that

$$\frac{\partial P}{\partial x} = \frac{\frac{x}{(x^2+y^2)^{1/2}} - 1}{\sqrt{(x^2 + y^2)^{1/2} - x}}, \quad [1 \text{ mark}] \quad \frac{\partial P}{\partial y} = \frac{y}{(x^2 + y^2)^{1/2} \sqrt{(x^2 + y^2)^{1/2} - x}}; \quad [1 \text{ mark}]$$

we could of course simplify the first of these but we don't really need to for this particular problem.

The unit circle can be parameterised as

$$(x(t), y(t)) = (\cos t, \sin t), \quad [1 \text{ mark}]$$

where the portion of the circle lying in U corresponds to the range $t \in (0, 2\pi)$, and the point $(1, 0)$ corresponds to $t = 0$. (That this value for t is not in the range just given merely means that the function Q is not defined at the point $(1, 0)$; it does not affect any other part of the problem.) Given this, we may evaluate the line integral as follows. Let $\theta \in (0, 2\pi)$; then

$$Q(\cos \theta, \sin \theta) = \int_{(1,0)}^{(\cos \theta, \sin \theta)} -\frac{\partial P}{\partial y} dx + \frac{\partial P}{\partial x} dy.$$

Now at the point $(\cos t, \sin t)$ on the unit circle we have

$$\frac{\partial P}{\partial x} = \frac{\cos t - 1}{\sqrt{1 - \cos t}}, \quad \frac{\partial P}{\partial y} = \frac{\sin t}{\sqrt{1 - \cos t}}, \quad [0.5 \text{ marks each}]$$

so this integral becomes

$$\begin{aligned} Q(\cos \theta, \sin \theta) &= \int_0^\theta -\frac{\sin t}{\sqrt{1 - \cos t}}(-\sin t) + \frac{\cos t - 1}{\sqrt{1 - \cos t}} \cos t dt && [1 \text{ mark for } -\sin t \text{ and } \cos t, \\ &= \int_0^\theta \frac{\sin^2 t + \cos^2 t - \cos t}{\sqrt{1 - \cos t}} dt && 1 \text{ mark for bounds}] \\ &= \int_0^\theta \sqrt{1 - \cos t} dt = \int_0^\theta \sqrt{2} \sin \frac{t}{2} dt && [1 \text{ mark}] \\ &= -2\sqrt{2} \cos \frac{t}{2} \Big|_0^\theta = 2\sqrt{2} \left(1 - \cos \frac{\theta}{2}\right), && [1 \text{ mark}] \end{aligned}$$

the desired answer. (Note that we are allowed to write $\sin \frac{t}{2} = \sqrt{\frac{1-\cos t}{2}}$, taking the positive square root, since $t \in (0, 2\pi)$ implies that $\sin \frac{t}{2} > 0$. It is instructive to think about this last statement geometrically.)

Penultimately, then, we have, taking the limits along the unit circle, [1 mark for $\theta \rightarrow 0^+$, 1 mark for $\theta \rightarrow 2\pi^-$, 1 mark for the final answer]

$$\begin{aligned} \lim_{(x,y) \rightarrow (1,0^+)} Q(x,y) - \lim_{(x,y) \rightarrow (1,0^-)} Q(x,y) &= \lim_{\theta \rightarrow 0^+} Q(\cos \theta, \sin \theta) - \lim_{\theta \rightarrow 2\pi^-} Q(\cos \theta, \sin \theta) \\ &= \lim_{\theta \rightarrow 0^+} 2\sqrt{2} \left(1 - \cos \frac{\theta}{2} \right) - \lim_{\theta \rightarrow 2\pi^-} 2\sqrt{2} \left(1 - \cos \frac{\theta}{2} \right) = 0 - 2\sqrt{2}(1 - (-1)) = -4\sqrt{2}. \end{aligned}$$

In this case, the function P is (up to a constant multiple) the imaginary part of the branch of the square-root function corresponding to the given set, and the half-line $\{(x,0)|x \geq 0\}$ is the associated branch cut. [1 mark for saying something related to branch cut or branch point]