

Tutorial 0401 (Friday, 10-11), Quiz 2, rubric.

It can be shown that on the set $U = \mathbf{R}^2 \setminus \{(x, 0) | x \leq 0\}$, i.e., the plane with the negative x axis and origin removed, the function

$$P(x, y) = 4\sqrt{(x^2 + y^2)^{\frac{1}{2}} + x}$$

is harmonic. The formula

$$Q(x_0, y_0) = \int_{(1,0)}^{(x_0, y_0)} -\frac{\partial P}{\partial y} dx + \frac{\partial P}{\partial x} dy$$

gives a function which is harmonic on U and such that

$$f(x + iy) = P(x, y) + iQ(x, y)$$

is analytic on U . Use this formula to determine Q on the part of the unit circle which lies in U . (The identity

$$\cos \frac{x}{2} = \sqrt{\frac{1 + \cos x}{2}},$$

valid when $\cos \frac{x}{2}$ is positive, may be useful.) What is

$$\lim_{(x,y) \rightarrow (-1,0^+)} Q(x, y) - \lim_{(x,y) \rightarrow (-1,0^-)} Q(x, y)$$

(where the limits are taken along the unit circle)? What do we call the half-line $\{(x, 0) | x \leq 0\}$ in this case?

First, we note that

$$\frac{\partial P}{\partial x} = 2 \frac{\frac{x}{(x^2+y^2)^{1/2}} + 1}{\sqrt{(x^2 + y^2)^{1/2} + x}}, \quad [1 \text{ mark}] \quad \frac{\partial P}{\partial y} = 2 \frac{y}{(x^2 + y^2)^{1/2} \sqrt{(x^2 + y^2)^{1/2} + x}}; \quad [1 \text{ mark}]$$

we could of course simplify the first of these but we don't really need to for this particular problem.

The unit circle can be parameterised as

$$(x(t), y(t)) = (\cos t, \sin t), \quad [1 \text{ mark}]$$

where the portion of the circle lying in U corresponds to the range $t \in (-\pi, \pi)$, and the point $(1, 0)$ corresponds to $t = 0$. Given this, we may evaluate the line integral as follows. Let $\theta \in (-\pi, \pi)$; then

$$Q(\cos \theta, \sin \theta) = \int_{(1,0)}^{(\cos \theta, \sin \theta)} -\frac{\partial P}{\partial y} dx + \frac{\partial P}{\partial x} dy.$$

Now at the point $(\cos t, \sin t)$ on the unit circle we have

$$\frac{\partial P}{\partial x} = 2 \frac{\cos t + 1}{\sqrt{\cos t + 1}}, \quad \frac{\partial P}{\partial y} = 2 \frac{\sin t}{\sqrt{1 + \cos t}}, \quad [0.5 \text{ marks each}]$$

so this integral becomes

$$\begin{aligned} Q(\cos \theta, \sin \theta) &= \int_0^\theta -2 \frac{\sin t}{\sqrt{1 + \cos t}} (-\sin t) + 2 \frac{\cos t + 1}{\sqrt{\cos t + 1}} \cos t dt && [1 \text{ mark for } -\sin t \text{ and } \cos t, \\ &= \int_0^\theta 2 \frac{\sin^2 t + \cos^2 t + \cos t}{\sqrt{1 + \cos t}} dt && 1 \text{ mark for bounds}] \\ &= \int_0^\theta 2\sqrt{1 + \cos t} dt = \int_0^\theta 2\sqrt{2} \cos \frac{t}{2} dt && [1 \text{ mark}] \\ &= 4\sqrt{2} \sin \frac{t}{2} \Big|_0^\theta = 4\sqrt{2} \sin \frac{\theta}{2}, && [1 \text{ mark}] \end{aligned}$$

the desired answer. (Note that we are allowed to write $\cos \frac{t}{2} = \sqrt{\frac{1+\cos t}{2}}$, taking the positive square root, since $t \in (-\pi, \pi)$ implies that $\cos \frac{t}{2} > 0$. It is instructive to think about this last statement geometrically.)

Penultimately, then, we have, taking the limits along the unit circle, [1 mark for π^- , 1 mark for π^+ , 1 mark for the final answer]

$$\begin{aligned} \lim_{(x,y) \rightarrow (-1,0^+)} Q(x,y) - \lim_{(x,y) \rightarrow (-1,0^-)} Q(x,y) &= \lim_{\theta \rightarrow \pi^-} Q(\cos \theta, \sin \theta) - \lim_{\theta \rightarrow -\pi^+} Q(\cos \theta, \sin \theta) \\ &= \lim_{\theta \rightarrow \pi^-} 4\sqrt{2} \sin \frac{\theta}{2} - \lim_{\theta \rightarrow -\pi^+} 4\sqrt{2} \sin \frac{\theta}{2} = 4\sqrt{2} - 4(-\sqrt{2}) = 8\sqrt{2}. \end{aligned}$$

In this case, the function P is (up to a constant multiple) the real part of the branch of the square-root function corresponding to the given set, and the half-line $\{(x,0)|x \leq 0\}$ is the associated branch cut. [1 mark for saying something related to branch cut or branch point]